CORRESPONDENCE

The Emergency Medical Care of Patients With Acute Myocardial Infarction—Results From the First Medical Contact Study in the Context of the Berlin Myocardial Infarction Registry

by PD. Dr. med. Martin Stockburger et al. in issue 29–30/2016

Additional Prehospital Problems

The authors have provided a valuable article on the care of patients with myocardial infarction (1). There is no disputing that correct diagnosis leads to swifter intervention or that training is required in this area. However, for the most part the authors limit themselves to seeking mistakes by emergency physicians, which is not the only factor in prehospital care. I will take the liberty of indicating other problems in prehospital care for STEMI, as the impression given in the article—that a high percentage of emergency physicians on the scene make errors in their work—is inaccurate.

It continues to occur that initial efforts to treat acute coronary syndrome take the form of emergency treatment, calling for an emergency physician to be present. Naturally, allocation of treatment depends on a complex interplay of many factors, but in Germany, if one examines the additional requests for emergency physicians in rural areas in particular, it leads to long delays.

How do emergency hospital admissions/catheterization laboratories work on the ground? Telemetry and discussions between physicians can generally ensure swift intervention as early as during prehospital care. How is it that almost 50% of correctly diagnosed STEMI cases in Berlin are initially admitted via the emergency room? Would the incorrectly diagnosed STEMI cases actually have benefited from a correct diagnosis?

A new profession is currently being developed in Germany: the emergency paramedic. The intention is that, with the aid of algorithms, they will be able to take on some work now done by physicians. Thus in the future much will depend on the composition and routine of emergency paramedic teams on the scene. As Mencl et al. have shown, regular training for paramedics was never successful in improving STEMI diagnosis outcomes beyond a certain amount (2).

DOI: 10.3238/arztebl.2017.0072a

REFERENCES

1. Stockburger M, Maier B, Fröhlich G, et al.: The emergency medical care of patients with acute myocardial infarction—results from the First Medical Contact Study in the context of the Berlin Myocardial Infarction Registry. *Dtsch Arztebl Int* 2016; 113: 497–502.
2. Mencl F, Wilber S, Frey J, et al.: Paramedic ability to recognize ST-segment elevation myocardial infarction on prehospital electrocardiograms. *Prehosp Emerg Care* 2013; 17: 203–10.

PD Dr. med. habil. Udo Meißner
Bamberg, Germany
udo.meissner@outlook.de

In Reply:

Many thanks to Dr. Meißner for his critical comments. We share his view that quality of care and care times for acute myocardial infarction are determined by multiple factors, including some that go beyond correct primary diagnosis of STEMI by emergency physicians. Correct initial allocation of a case by the emergency services central office, with sufficient requesting of emergency physicians, certainly plays a major role in this. A high level of well-integrated collaboration is also required for the interface between prehospital care by emergency physicians and subsequent hospital treatment to function smoothly. This is why the First Medical Contact Study of the Berlin Myocardial Infarction Registry (BHIR, *Berliner Herzinfarktregister*) (1) was conducted jointly by hospital departments and the Berlin Fire Department as suppliers of emergency medical services and care by emergency physicians.

Exploratory analysis of outlier values in BHIR procedure data had once again shown the marked trigger function of ECG diagnosis of STEMI by emergency physicians for subsequent inpatient care. This had already been described elsewhere (2). The First Medical Contact Study was initiated to provide quantitative analysis of this trigger and to identify any potential for improvement. Its findings do indeed indicate that there is potential for optimization. However, this by no means attributes all errors to a single party as Dr. Meißner suspects; unfortunately, representation of an analysis of all issues, from initial diagnosis to catheter intervention, goes beyond the scope of an article in *Deutsches Ärzteblatt*. Like Dr. Meißner, we believe that structured prehospital care with quality ensured by, for example, telemedical transfer is particularly important, also regarding the introduction of emergency paramedics. Detailed planning is currently underway for another, multimodal, collaborative regional project to improve various aspects of prehospital care for myocardial infarction.

DOI: 10.3238/arztebl.2017.0072b

REFERENCES

1. Stockburger M, Maier B, Fröhlich G, et al.: The emergency medical care of patients with acute myocardial infarction—results from the First Medical Contact Study in the context of the Berlin Myocardial Infarction Registry. *Dtsch Arztebl Int* 2016; 113: 497–502.
2. Brown JP, Mahmud E, Dunford JV, Ben-Yehuda O: Effect of prehospital 12-lead electrocardiogram on activation of the cardiac catheterization laboratory and door-to-balloon time in ST-segment elevation acute myocardial infarction. *Am J Cardiol* 2008; 101: 158–61.

PD Dr. med. Martin Stockburger
Medizinische Klinik I, Nauen
Germany
martin.stockburger@havelland-kliniken.de

Conflict of interest statement

The authors of all contributions declare that no conflict of interest exists.