Direkte DNA-Analysen im <i>GCH1</i> -Gen (<i>DYT5</i>), <i>ICD10</i> G24.1V bei Verdacht auf familiäre dopa-responsive Dystonie	Direkte DNA-Analyse im <i>GCH1</i> -Gen (<i>DYT5</i>), <i>ICD10</i> G24.1V bei Verdacht auf familiäre dopa-responsive Dystonie
<p>Patient: [redacted] DNA-Nr.: 110980(20168521)</p> <p>Probeneingang: 30.05.2011; Material: 1 x 8 ml EDTA-Blut Auftrag vom: Siehe Probeneingang</p> <p>Klinische Angaben: Dopa-responsive Dystonie</p> <p>Familienanamnese: keine Angabe</p>	<p>Patient: [redacted] DNA-Nr.: 110980 (20168521)</p> <p>Probeneingang: 30.05.2011; Material: 1 x 8 ml EDTA-Blut Auftrag vom: Siehe Probeneingang</p> <p>Klinische Angaben: Dopa-responsive Dystonie</p> <p>Familienanamnese: keine Angabe</p>
<p>wir möchten Ihnen den Befund der molekulargenetischen Analysen bei [redacted] mitteilen. Hinweise aus der Familienanamnese bzw. dem Erkrankungsverlauf hatten Sie an eine familiäre dopa-responsive Dystonie mit Mutation im <i>GCH1</i>-Gen (<i>DYT5</i>) denken lassen.</p> <p>Aus der eingesandten Probe wurden durch PCR alle kodierenden Abschnitte (Exone 1-6; ENST491895) des <i>GCH1</i>-Gens amplifiziert. Die PCR-Amplifikate wurden anschließend direkt sequenziert (ABI3730XL). Sequenzdaten wurden im SequencePilot (si-medi) auf Sequenzabweichungen hin untersucht. Diese Methode erlaubt, mehr als 99% aller Punktmutationen zu finden. Zusätzlich wurden die Exone 1, 2, 3, 5 & 6 durch MLPA (Kit 099) auf genomische Deletionen getestet. Intrinsische Veränderungen, die sehr selten auch klinisch relevant sein können, werden meistens nicht nachgewiesen. Abrechnungsinformation für ambulante Patienten: 11320 [5x], 11321 [7x], 11322 [7x].</p> <p>Mutationen im <i>GCH1</i>-Gen führen zu einem autosomal-dominant erblichen Dystonie-Syndrom (<i>DYT5</i>). Die Erkrankung wird typischerweise in der Kindheit als fluktuierende dystone Gangstörung mit Generalisierungstendenz und parkinsonartiger Begleitsymptomatik diagnostiziert. Levodopa-Gaben verbessern diese Symptome dramatisch und anhaltend.</p>	<p>wir möchten Ihnen den Befund der molekulargenetischen Analysen bei [redacted] mitteilen. Hinweise aus der Familienanamnese bzw. dem Erkrankungsverlauf hatten Sie an eine familiäre dopa-responsive Dystonie mit Mutation im <i>GCH1</i>-Gen (<i>DYT5</i>) denken lassen.</p> <p>Aus der eingesandten Probe wurden durch PCR alle kodierenden Abschnitte (Exone 1-6; ENST491895) des <i>GCH1</i>-Gens amplifiziert. Die PCR-Amplifikate wurden anschließend direkt sequenziert (ABI3730XL). Sequenzdaten wurden im SequencePilot (si-medi) auf Sequenzabweichungen hin untersucht. Diese Methode erlaubt, mehr als 99% aller Punktmutationen zu finden. Zusätzlich wurden die Exone 1, 2, 3, 5 & 6 durch MLPA (Kit 099) auf genomische Deletionen getestet. Intrinsische Veränderungen, die sehr selten auch klinisch relevant sein können, werden meistens nicht nachgewiesen. Abrechnungsinformation für ambulante Patienten: 11320 [5x], 11321 [7x], 11322 [7x].</p> <p>Mutationen im <i>GCH1</i>-Gen führen zu einem autosomal-dominant erblichen Dystonie-Syndrom (<i>DYT5</i>). Die Erkrankung wird typischerweise in der Kindheit als fluktuierende dystone Gangstörung mit Generalisierungstendenz und parkinsonartiger Begleitsymptomatik diagnostiziert. Levodopa-Gaben verbessern diese Symptome dramatisch und anhaltend.</p>
<p>Bei [redacted] wurde eine <i>GCH1</i>-Mutation nachgewiesen.</p> <p>Damit konnte die klinische Verdachtsdiagnose bisher molekulargenetisch bestätigt werden.</p>	<p>Bei [redacted] wurde keine <i>GCH1</i>-Mutation nachgewiesen.</p> <p>Damit konnte die klinische Verdachtsdiagnose bisher molekulargenetisch nicht bestätigt werden.</p>
<p>Exonische Polymorphismen: Keine</p> <p>Entsprechend der Vorgaben des Gendiagnostik-Gesetzes (§9) sollen Patienten im Rahmen einer genetischen Untersuchung auf die Möglichkeit einer genetischen Beratung hingewiesen werden.</p> <p>Wir möchten uns bei Ihnen für die Überweisung der Diagnostik bedanken. Mit freundlichen Grüßen</p>	<p>Exonische Polymorphismen: Keine</p> <p>Entsprechend der Vorgaben des Gendiagnostik-Gesetzes (§9) sollen Patienten im Rahmen einer genetischen Untersuchung auf die Möglichkeit einer genetischen Beratung hingewiesen werden.</p> <p>Wir möchten uns bei Ihnen für die Überweisung der Diagnostik bedanken. Mit freundlichen Grüßen</p>

Direkte DNA-Analyse im <i>TH</i> -Gen (<i>DYT5</i>), <i>ICD10</i> G24.1V bei Verdacht auf familiäre dopa-responsive Dystonie	Direkte DNA-Analyse im <i>TH</i> -Gen (<i>DYT5</i>), <i>ICD10</i> G24.1V bei Verdacht auf familiäre dopa-responsive Dystonie
<p>Patient: [redacted] DNA-Nr.: 110980(20168521)</p> <p>Probeneingang: 30.05.2011; Material: 1 x 8 ml EDTA-Blut Auftrag vom: Siehe Probeneingang</p> <p>Klinische Angaben: Dopa-responsive Dystonie</p> <p>Familienanamnese: keine Angabe</p>	<p>Patient: [redacted] DNA-Nr.: 110980 (20168521)</p> <p>Probeneingang: 30.05.2011; Material: 1 x 8 ml EDTA-Blut Auftrag vom: Siehe Probeneingang</p> <p>Klinische Angaben: Dopa-responsive Dystonie</p> <p>Familienanamnese: keine Angabe</p>
<p>wir möchten Ihnen den Befund der molekulargenetischen Analysen bei [redacted] mitteilen. Hinweise aus der Familienanamnese bzw. dem Erkrankungsverlauf hatten Sie an eine familiäre dopa-responsive Dystonie mit Mutation im <i>TH</i>-Gen (<i>DYT5</i>) denken lassen.</p> <p>Aus der eingesandten Probe wurden durch PCR alle kodierenden Abschnitte (Exone 1-14; ENST381178) des <i>TH</i>-Gens amplifiziert. Die PCR-Amplifikate wurden anschließend direkt sequenziert (ABI3730XL). Sequenzdaten wurden im SequencePilot (si-medi) auf Sequenzabweichungen hin untersucht. Diese Methode erlaubt, mehr als 99% aller Punktmutationen zu finden. Große genomische Deletion und (selten relevante) intrinsische Veränderungen können meistens nicht nachgewiesen werden. Abrechnungsinformation für ambulante Patienten: 11321 [11x], 11322 [11x].</p> <p>Mutationen im <i>TH</i>-Gen führen zu einem autosomal-rezessiv erblichen Dystonie-Syndrom (<i>DYT5</i>). Die Erkrankung wird typischerweise in der Kindheit als fluktuierende dystone Gangstörung mit Generalisierungstendenz und parkinsonartiger Begleitsymptomatik diagnostiziert. Levodopa-Gaben verbessern diese Symptome dramatisch und anhaltend.</p>	<p>wir möchten Ihnen den Befund der molekulargenetischen Analysen bei [redacted] mitteilen. Hinweise aus der Familienanamnese bzw. dem Erkrankungsverlauf hatten Sie an eine familiäre dopa-responsive Dystonie mit Mutation im <i>TH</i>-Gen (<i>DYT5</i>) denken lassen.</p> <p>Aus der eingesandten Probe wurden durch PCR alle kodierenden Abschnitte (Exone 1-14; ENST381178) des <i>TH</i>-Gens amplifiziert. Die PCR-Amplifikate wurden anschließend direkt sequenziert (ABI3730XL). Sequenzdaten wurden im SequencePilot (si-medi) auf Sequenzabweichungen hin untersucht. Diese Methode erlaubt, mehr als 99% aller Punktmutationen zu finden. Große genomische Deletion und (selten relevante) intrinsische Veränderungen können meistens nicht nachgewiesen werden. Abrechnungsinformation für ambulante Patienten: 11321 [11x], 11322 [11x].</p> <p>Mutationen im <i>TH</i>-Gen führen zu einem autosomal-rezessiv erblichen Dystonie-Syndrom (<i>DYT5</i>). Die Erkrankung wird typischerweise in der Kindheit als fluktuierende dystone Gangstörung mit Generalisierungstendenz und parkinsonartiger Begleitsymptomatik diagnostiziert. Levodopa-Gaben verbessern diese Symptome dramatisch und anhaltend.</p>
<p>Bei [redacted] wurde eine <i>TH</i>-Mutation nachgewiesen.</p> <p>Damit konnte die klinische Verdachtsdiagnose bisher molekulargenetisch bestätigt werden.</p>	<p>Bei [redacted] wurden keine <i>TH</i>-Mutationen nachgewiesen.</p> <p>Damit konnte die klinische Verdachtsdiagnose bisher molekulargenetisch nicht bestätigt werden.</p>
<p>Exonische Polymorphismen: c.813G>A, p.K271K heterozygot (rs6357)</p> <p>Entsprechend der Vorgaben des Gendiagnostik-Gesetzes (§9) sollen Patienten im Rahmen einer genetischen Untersuchung auf die Möglichkeit einer genetischen Beratung hingewiesen werden.</p> <p>Wir möchten uns bei Ihnen für die Überweisung der Diagnostik bedanken. Mit freundlichen Grüßen</p>	<p>Exonische Polymorphismen: c.813G>A, p.K271K heterozygot (rs6357)</p> <p>Entsprechend der Vorgaben des Gendiagnostik-Gesetzes (§9) sollen Patienten im Rahmen einer genetischen Untersuchung auf die Möglichkeit einer genetischen Beratung hingewiesen werden.</p> <p>Wir möchten uns bei Ihnen für die Überweisung der Diagnostik bedanken. Mit freundlichen Grüßen</p>

The manipulated genetic reports (left side) and original reports (right side) of *GCH1* and *TH* gene mutation testing are shown. Personal details of the patient and treating physicians are obscured. The main forged passages of the report are highlighted in red. In the original report it is stated that "keine *GCH1/TH*-Mutation nachgewiesen" ("no *GCH1/TH* mutation was found"). The forged report says "eine *GCH1/TH*-Mutation nachgewiesen" ("a *GCH1/TH* mutation was found"). Of note, the critical difference in German lies in the presence or absence of a single letter. The original report also describes that the clinical diagnosis could not be confirmed ("nicht bestätigt"), in contrast to the forged report, stating that the diagnosis is confirmed ("bestätigt").

Study funding: No targeted funding reported.

Disclosure: The authors report no disclosures relevant to the manuscript. Go to Neurology.org for full disclosures.

Received August 15, 2016. Accepted in final form November 28, 2016.

Correspondence to Dr. Münchau: alexander.muenchau@neuro.uni-luebeck.de

© 2017 American Academy of Neurology

1. Asher R. Münchhausen's syndrome. *Lancet* 1951;1:339-341.

- Anderson KE, Weiner WJ, Lang AE. Advances in Neurology: Behavioral Neurology of Movement Disorders. Philadelphia: Lippincott Williams & Wilkins; 2005.
- Chen WJ, Lin Y, Xiong ZQ, et al. Exome sequencing identifies truncating mutations in PRRT2 that cause paroxysmal kinesigenic dyskinesia. *Nat Genet* 2011;43:1252-1255.
- Chen YZ, Friedman JR, Chen DH, et al. Gain-of-function *ADCY5* mutations in familial dyskinesia with facial myokymia. *Ann Neurol* 2014;75:542-549.

William C. Kreisl, MD
Robin Lawrence
Emily Page
Andrew F. Teich, MD,
PhD
Deepa S. Subramaniam,
MD, MSc
Robert B. Innis, MD, PhD
R. Scott Turner, MD, PhD

¹¹C-PBR28 PET DETECTS TRANSLOCATOR PROTEIN IN A PATIENT WITH ASTROCYTOMA AND ALZHEIMER DISEASE

A 66-year-old woman with probable Alzheimer disease (AD) enrolled in National Institute of Mental Health protocols 09-M-0198 and 08-M-0066. The protocols were approved by the Combined Neurosciences Institutional Review Board, and the patient gave informed consent before entering the study.

Brian MRI showed diffuse atrophy and a small T2 hyperintensity in the right posterior temporal region (0.4 × 1.4 cm) interpreted as nonspecific gliosis (figure). ¹⁸Fluorodeoxyglucose PET revealed bilateral temporo-parietal hypometabolism (with no hypermetabolism in the area of the T2 hyperintensity) and ¹¹C-Pittsburgh compound B PET imaging showed fibrillar β-amyloid retention, both consistent with AD pathophysiology. PET imaging with

^{11}C -PBR28, a radioligand for the 18 kDa translocator protein (TSPO), was performed to measure microglial activation. Diffuse ^{11}C -PBR28 binding was found in gray matter, with a small focus of increased binding colocalized to the T2 hyperintensity. ^{11}C -PR28 binding was 19% greater in the T2 hyperintensity than in an identical volume in contralateral brain (figure).

Over the following year, the patient's dementia progressed. Noncontrast brain MRI revealed increase of the T2 hyperintensity (0.7×1.6 cm), again interpreted as nonspecific gliosis; neoplasm was considered less likely. Serial noncontrast brain MRIs obtained every 3–6 months revealed stable T2 hyperintensity in this region.

Two years after the first ^{11}C -PR28 PET scan, noncontrast brain MRI revealed increased size of the T2 hyperintensity ($2.2 \times 1.2 \times 1.7$ cm; figure). Repeat ^{11}C -PBR28 PET imaging again showed diffuse binding, with enlargement of the focal area that colocalized to the T2 hyperintensity. ^{11}C -PR28 binding was 26% greater in the T2 hyperintensity than in contralateral brain (figure). MRI performed with contrast revealed no gadolinium enhancement. Neuro-oncology consultation was obtained and low-grade astrocytoma was considered the most likely diagnosis. Five months later, MRI revealed a ring-enhancing cortical mass ($1.3 \times 2.0 \times 1.7$ cm) in the right posterior superior

Figure Imaging and histopathologic findings in a patient with Alzheimer disease (AD) later diagnosed with glioblastoma multiforme

(A) Results of longitudinal imaging of a patient with AD who was later diagnosed with glioblastoma multiforme. Baseline fluid-attenuated inversion recovery (FLAIR) image shows a small area of hyperintensity in the right temporal region and ^{11}C -PBR28 PET image shows a small area of increased binding in the same region (white arrows). Pittsburgh compound B (PIB) PET image shows diffuse neocortical binding typical of patients with AD. FDG PET scan shows bilateral temporal hypometabolism (yellow arrows). Follow-up FLAIR image shows increased size of the right temporal hyperintensity, with corresponding increase in size of focal binding on follow-up ^{11}C -PBR28 PET. Gadolinium-enhanced T1 image performed at the time of follow-up ^{11}C -PBR28 PET does not demonstrate any evidence of blood-brain barrier disruption. However, 5 months later, gadolinium enhancement is appreciated on repeat T1 MRI, suggesting progression of the underlying glioma. Standardized uptake value ratio (SUVR) ^{11}C -PBR28 images were created by averaging images from 60 to 90 minutes postinjection and then normalizing to binding in cerebellar gray matter. Baseline and follow-up ^{11}C -PBR28 PET images were smoothed and windowed to identical minimum and maximum SUVR for presentation. (B) Amount of ^{11}C -PBR28 binding in the region of T2 hyperintensity (black bars) and in a region with the same volume on the contralateral side of the brain (white bars), at baseline and at follow-up. (C) Hematoxylin & eosin stain of surgical tissue shows hypercellularity, pseudopalisades, and necrosis consistent with glioblastoma multiforme (40× magnification). (D) Anti-CD68 stains ramified microglia but not the infiltrating glioma cells (60×). (E) In the same field, anti-TSPO stains microglia, but also the nuclear rim of the glioma cells (red arrows). (F) Hematoxylin & eosin stain in a field distant from tumor burden shows normal gray matter cytoarchitecture with neurons and astrocytes (40×). (G) Anti- β -amyloid immunostain in same field shows 2 dense core plaques typical of AD (60×). (H) Anti-tau immunostain in same field shows neurofibrillary tangles are also present. In the same field, anti-CD68 immunostaining shows ramified microglia (I), which are also positive on anti-TSPO staining (J).

and middle temporal gyri with surrounding edema and mass effect, consistent with malignant transformation to high-grade astrocytoma.

Right temporoparietal craniotomy and tumor resection was performed. Pathologic examination revealed glioblastoma multiforme (GBM) WHO grade IV. Neural tissue was densely infiltrated with astrocytoma cells with pleomorphic nuclei and mitotic figure (figure). Tumor cells were positive for glial fibrillary acid protein and p53, with proliferative index at 30% estimated by KI67 immunostaining. CD68+ microglia with activated morphology surrounded tumor cells, and both tumor cell nuclei and microglia expressed TSPO (figure). Molecular profiling showed unmethylated methylguanine methyltransferase promoter, no mutations in tumor suppressor genes including *TP53*, no loss of PTEN expression, and variant transcript for EGFRvIII. Gray matter distant from tumor burden showed relatively preserved cytoarchitecture (figure). In this region, amyloid and tau staining showed dense core plaques and neurofibrillary tangles (figure). In tissue with low tumor burden and high plaque and tangle burden, TSPO was mostly expressed by CD68+-activated microglia (figure).

Discussion. Both astrocytoma and AD are associated with overexpression of TSPO in brain. In astrocytoma, TSPO is overexpressed by tumor cells and activated microglia,¹ while in AD, TSPO is primarily expressed by microglia.² PET imaging can measure TSPO in vivo³; however, off-target binding of TSPO radioligands and paucity of pathologic confirmation precludes conclusive interpretation of most human PET studies. This case allows a unique opportunity, in 2 different disease states, to (1) evaluate the ability of ¹¹C-PBR28 to detect abnormal expression of TSPO and (2) confirm which cells express TSPO in each disease. In this patient, focal ¹¹C-PBR28 binding in the right temporal lobe corresponded to a nonenhancing astrocytoma that progressed to pathologically confirmed GBM with TSPO-expressing tumor cells. This focal binding was present before hypermetabolism on FDG or gadolinium enhancement on MRI. The clinical course suggested transformation to secondary GBM while molecular profiling could suggest primary GBM. In either case, ¹¹C-PBR28 binding signaled malignant behavior prior to other imaging modalities. The patient also had diffuse binding of ¹¹C-PBR28 in neocortex at baseline and at follow-up. On pathologic examination of gray matter where cytoarchitecture was relatively spared of tumor burden, β -amyloid plaques and neurofibrillary tangles were associated with TSPO-expressing activated microglia. These results suggest that ¹¹C-PBR28 also detected AD-associated microgliosis.

TSPO appears to be a marker of proliferation in astrocytoma, with expression increasing with tumor grade.⁴ In AD, whether dysfunctional microglial activity plays a primary pathophysiologic role remains unclear. However, TSPO binding increases with AD severity and progression of disease.^{5,6} This report provides evidence that ¹¹C-PBR28 binding on PET corresponds to TSPO density—expressed by tumor cells in astrocytoma and by activated microglia in AD—and supports a role of ¹¹C-PBR28 in quantifying progression of both diseases.

From the Molecular Imaging Branch (W.C.K., E.P., R.B.I.), National Institute of Mental Health, Bethesda, MD; Taub Institute (W.C.K.), Columbia University Medical Center (A.F.T.), New York, NY; and Memory Disorders Program (R.L., R.S.T.) and Lombardi Comprehensive Cancer Center (D.S.S.), Georgetown University, Washington, DC.

Author contributions: William C. Kreisl: study concept and design, acquisition of data, analysis and interpretation of data, drafting the manuscript. Robin Lawrence: acquisition of data, drafting the manuscript. Emily Page: acquisition of data, critical revision of manuscript for intellectual content. Andrew Teich: analysis and interpretation of data, critical revision of manuscript for intellectual content. Deepa S. Subramaniam: acquisition of data, interpretation of data, drafting the manuscript. Robert B. Innis: acquisition of data, study concept and design, critical revision of manuscript for intellectual content, study supervision. R. Scott Turner: study concept and design, critical revision of manuscript for intellectual content, study supervision.

Acknowledgment: The authors thank the patient for her consent to publish this report. Yi Zhang (NIMH) provided assistance in the production of radioligands. PET scans were performed at the NIH PET department. Brent Harris, MD, PhD, performed neuropathologic evaluation.

Study funding: Funded by the National Institute of Mental Health project numbers ZIAMH002852 and ZIAMH0022793 under clinicaltrials.gov identifiers NCT00955422 and NCT00613119 (protocols 08-M-0066 and 09-M-0198).

Disclosure: The authors report no disclosures relevant to the manuscript. Go to Neurology.org for full disclosures.

Received August 17, 2016. Accepted in final form December 6, 2016.

Correspondence to Dr. Kreisl: wck2107@cumc.columbia.edu

© 2017 American Academy of Neurology

1. Su Z, Roncaroli F, Durrenberger PF, et al. The 18-kDa mitochondrial translocator protein in human gliomas: an ¹¹C-(R)PK11195 PET imaging and neuropathology study. *J Nucl Med* 2015;56:512–517.
2. Cosenza-Nashat M, Zhao ML, Suh HS, et al. Expression of the translocator protein of 18 kDa by microglia, macrophages and astrocytes based on immunohistochemical localization in abnormal human brain. *Neuropathol Appl Neurobiol* 2009;35:306–328.
3. Owen DR, Matthews PM. Imaging brain microglial activation using positron emission tomography and translocator protein-specific radioligands. *Int Rev Neurobiol* 2011;101:19–39.
4. Miettinen H, Kononen J, Haapasalo H, et al. Expression of peripheral-type benzodiazepine receptor and diazepam binding inhibitor in human astrocytomas: relationship to cell proliferation. *Cancer Res* 1995;55:2691–2695.

5. Kreisl WC, Lyoo CH, McGwier M, et al. In vivo radioligand binding to translocator protein correlates with severity of Alzheimer's disease. *Brain* 2013;136:2228–2238.
6. Kreisl WC, Lyoo CH, Liow JS, et al. 11C-PBR28 binding to translocator protein increases with progression of Alzheimer's disease. *Neurobiol Aging* 2016;44:53–61.

Joshua Barton, MD
 Todd A. Hardy, MD, PhD
 Sean Riminton, MD, PhD
 Stephen W. Reddel, MD, PhD
 Yael Barnett, MD
 Alasdair Coles, MD, PhD
 Michael H. Barnett, MD, PhD

TUMEFACTIVE DEMYELINATION FOLLOWING TREATMENT FOR RELAPSING MULTIPLE SCLEROSIS WITH ALEMTUZUMAB

Alemtuzumab, a humanized anti-CD52 monoclonal antibody that results in profound depletion and repopulation of mature lymphocytes, is a highly efficacious therapy for active, relapsing-remitting multiple sclerosis (RRMS). In 2 phase 3 clinical trials of alemtuzumab, Comparison of Alemtuzumab and Rebif Efficacy in Multiple Sclerosis (CARE-MS) I and II, 14% and 23% of patients treated with alemtuzumab had a clinical relapse within the first 12 months, respectively.^{1,2}

We report a patient with typical RRMS who developed a tumefactive demyelinating lesion (TDL) 4 months after treatment with alemtuzumab. We hypothesize that dynamic alteration in lymphocyte populations may underpin early disease activity in occasional patients treated with alemtuzumab.

Case report. A 39-year-old woman with longstanding mild neutropenia was diagnosed with RRMS at age 24 years following an episode of myelitis on a background of left facial numbness at age 21 years. Brain MRI

showed typical demyelinating lesions. Glatiramer acetate was started at age 26 following right-sided optic neuritis and switched to fingolimod at age 34 after left-sided optic neuritis occurred. Fingolimod was ceased after 15 months due to radiologic disease progression and profound lymphopenia (lymphocyte nadir $0.13 \times 10^9/L$). Natalizumab was commenced and the patient remained stable for 22 months before treatment was switched to dimethyl fumarate (DMF) at age 36 years due to positive serum JC polyomavirus (JCV) antibody testing (StratifyJCV; Unilabs, Denmark). Recurrent clinical and radiologic disease activity after 8 months prompted escalation of treatment to alemtuzumab (administered over 5 consecutive days according to protocol) following a 4-week DMF washout. The pretreatment lymphocyte count was $1.0 \times 10^9/L$ and the neutrophil count was $1.9 \times 10^9/L$.

Four months after treatment with alemtuzumab, the patient developed burning dysesthesia in the right arm that progressed over 1 month to involve the entire right leg, associated with intermittent gait ataxia. Brain MRI (figure) demonstrated a left hemisphere TDL associated with mild edema and an

Figure MRI of tumefactive demyelinating lesion

Brain MRI at presentation shows a large ($3 \times 2.5 \times 1.7$ cm), hyperintense lesion in the left hemisphere (A; fluid-attenuated inversion recovery [FLAIR]) that exhibits an incomplete ring of gadolinium enhancement (D; T1 with contrast) and restricted diffusion (B; diffusion-weighted imaging, E; apparent diffusion coefficient map). Progress imaging performed 3 months later shows substantial involution of the lesion (C; FLAIR) and resolution of gadolinium enhancement (F; T1 with contrast).