Published in final edited form as:

JAMA Ophthalmol. 2016 June 01; 134(6): 672–673. doi:10.1001/jamaophthalmol.2016.0562.

Commentary on “The effect of multi-spot laser panretinal photocoagulation on retinal sensitivity and driving eligibility in diabetic retinopathy”

Cynthia Owsley, PhD, MSPH and

Department of Ophthalmology, School of Medicine, University of Alabama at Birmingham

Gerald McGwin Jr., MS, PhD

Department of Epidemiology, School of Public Health, University of Alabama at Birmingham

In a study performed in the United Kingdom (UK), the authors report that the vast majority of study patients who underwent bilateral multi-spot pan-retinal laser photocoagulation (PRP) for proliferative diabetic retinopathy met the UK’s visual field standard for holding a car or motorcycle license six months following treatment. As the authors point out, although PRP has been shown to reduce the risk of severe vision loss from proliferative diabetic retinopathy, the treatment can come at a cost to peripheral vision, namely, loss of peripheral light sensitivity. In this study, the authors document some loss of peripheral vision; however, 35 of 37 (92%) of the patients at six months still met the UK’s visual field requirement for licensure six months following treatment.

Driving is a primary mode of personal travel in the UK and many other countries, and is linked to health and well-being. Previous research has demonstrated that driving cessation and transportation challenges are associated with increased likelihood of depression, placement into long-term care, and poor health.^{1,2} Thus, from a quality of life perspective, it is an important finding, and provides some encouragement to patients who want to be drivers; peripheral visual loss caused by the PRP procedure under study did not make the vast majority of patients ineligible for driving licensure based on the visual field standard. A cautionary note is that laser scar can expand beyond six months creating the potential for more extensive peripheral vision loss, as the authors acknowledge. Thus, longer term follow-up of these patients is necessary before strong conclusions can be made on the impact of multi-spot PRP, as performed in this study, on maintenance of eligibility for licensure beyond six months.

Being able to pass the UK visual field standard does not address the safety or performance of these drivers on the road. The visual field screening approach used by the UK does not have proven efficacy with respect to driver safety; that is, we do not know if the screening approach and the cut-point for passing is associated with a lower rate of motor vehicle collisions, as compared to those who fail. Policies for vision standards (visual acuity or visual field) for driver licensure in jurisdictions around the world, including the US, are largely not based on safety data, and thus, the UK is not alone in this regard. In the US, the

situation is further exacerbated by the fact that even though most states have visual field requirements for licensure, some states do not screen for visual field at license application or renewal, rendering the policy unenforceable.³

It also remains to be determined whether actual on-road performance by drivers post PRP is negatively impacted by the peripheral vision reductions caused by PRP. There are a variety of on-road research methodologies that could address this question,⁴ but no published research studies to date on the issue. Visual field loss due to other eye conditions can cause problems in lane control,⁵ thus raising the question of whether lane control might also be the case in patients with PRP-associated peripheral vision loss. Furthermore, visual field impairment in older drivers, regardless of etiology, has been associated with increased motor vehicle collision risk.⁶

Before widespread generalization of these findings, it is important to acknowledge that the study was based on a small sample of 37 patients at the six-month follow-up. Furthermore, the study excluded patients with other ophthalmic conditions, which also reduces generalization, especially since ocular co-morbidities, in addition to diabetic retinopathy, are common in persons with diabetes. It is important to point out that diabetes and its other complications can impact eligibility for licensure, and can also impact driving safety and performance.⁷ In addition, less than half of the drivers (46.5%) in the study were currently driving at baseline, and the article does not indicate whether the enrollees actually wanted to return to driving.

In summary, this study presents encouraging information for UK patients with proliferative diabetic retinopathy who undergo multi-spot laser panretinal photocoagulation; the results suggest that for some patients, eligibility for driving licensure based on the UK visual field standard is likely to be intact six months later. Future research should not only address this issue in larger samples allowing for ocular co-morbidities with longer follow-up time, but should also examine the driver safety and on-road performance of patients who undergo PRP. It is important to establish not just their eligibility for licensure, but the implications of PRP for collision risk and on-road performance.

Acknowledgments

Preparation of this commentary was made possible by grants from the National Eye Institute (R01EY18966), the National Institute on Aging (P30AG22838), Research to Prevent Blindness, and the EyeSight Foundation of Alabama.

References

1. Ragland DR, Satariano WA, MacLeod KE. Driving cessation and increased depressive symptoms. *J Gerontol A Biol Med Sci*. 2005; 60a:399–403.
2. Freeman EE, Gange SJ, Munoz B, West SK. Driving status and risk of entry into long-term care in adults. *Am J Public Health*. 2006; 96:1254–1259.
3. Prevent Blindness. State Vision Screening and Standards for License to Drive. 2003. <http://lowvision.preventblindness.org/daily-living-2/state-vision-screening-and-standards-for-license-to-drive> Accessed February 1, 2016
4. Owsley C, Wood J, McGwin G Jr. A roadmap for interpreting the literature on vision and driving. *Sur Ophthalmol*. 2015; 60:250–262.

5. Wood JM, McGwin G Jr, Elgin J, et al. On-road driving performance by persons with hemianopia and quadrantanopia. *Invest Ophthalmol Vis Sci*. 2009; 50:577–585. [PubMed: 18936138]
6. Huisinigh C, McGwin G Jr, Wood J, Owsley C. The driving visual field and a history of motor vehicle collision involvement in older drivers: A population-based examination. *Invest Ophthalmol Vis Sci*. 2015; 56:132–138.
7. American Diabetes Association. Diabetes and driving. *Diabetes Care*. 2012; 35(Supplement):581–586. [PubMed: 22301117]