Published in final edited form as:

Psychiatry Res. 2011 August 15; 188(3): 320–326. doi:10.1016/j.psychres.2011.05.037.

Neuropsychological performance in patients with schizophrenia and controls as a function of cigarette smoking status

Victoria C. Wing^{a,*}, Ingrid Bacher^a, Kristi A. Sacco^b, and Tony P. George^{a,b}

^aCentre for Addiction and Mental Health (CAMH), Schizophrenia Program; Division of Addiction Psychiatry, Department of Psychiatry, University of Toronto, Toronto, ON, Canada

^bProgram for Research in Smokers with Mental Illness (PRISM); Division of Substance Abuse, Department of Psychiatry, Yale University School of Medicine, New Haven, CT, USA

Abstract

Schizophrenia is associated with many neurocognitive deficits, some of which are improved by nicotine and cigarette smoking. To better understand the relationship between smoking and cognitive function in schizophrenia, cross-sectional assessment of neuropsychological performance as a function of smoking status (smoker or non-smoker) and smoking history (current, former or never-smoker) in clinically-stable outpatients with schizophrenia and controls was evaluated. Subjects ($N=140$) were divided into subgroups on the basis of self-report and biochemical verification of smoking history. Current smokers with schizophrenia ($n=38$), former smokers with schizophrenia ($n=17$), never-smokers with schizophrenia ($n=12$), control smokers ($n=31$), control former smokers ($n=16$), and control never-smokers ($n=26$) were administered a comprehensive neuropsychological battery. Smokers were studied under non-deprivation conditions.

Comparison of neuropsychological performance in schizophrenia and control subjects revealed significant main effects of diagnosis. Analysis of the data as a function of smoking history demonstrated that never-smokers with schizophrenia performed the poorest on measures of sustained attention, processing speed and response inhibition, when compared to the other schizophrenia subgroups. Cigarette smoking did not alter neuropsychological performance in controls.

Our findings suggest that smoking status and history differentially alters neuropsychological outcomes in schizophrenia compared to non-psychiatric controls, and that never-smokers may present with more severe neurocognitive impairments.

* **Address for Correspondence:** Centre for Addiction and Mental Health 33 Russell St., room 1910A Toronto, Ontario, Canada, M5S2S1 Tel: (416) 535-8501 ext.4882 Fax: (416) 979 4676 vicky_wing@camh.net.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Declarations: Drs. Wing, Bacher and Sacco have no conflicts to report. Dr. George reports that he has received grant support from Pfizer, Inc., and has received consulting fees from Pfizer, Evotec, Eli Lilly, Janssen-Ortho and Astra-Zeneca in the past two years.

Keywords

Schizophrenia; Cognition; Cigarette Smoking; Never-smoker; Sustained Attention; Processing Speed

1. Introduction

It has been suggested that the higher rates of tobacco smoking in schizophrenia compared to the general population may relate, at least in part, to pro-cognitive effects of nicotine (Adler et al., 1993; Levin et al., 1996; Dalack et al., 1998; Depatie et al., 2002; George et al., 2002; Smith et al., 2002; Sacco et al., 2005; Smith et al., 2006; Barr et al., 2008). In fact, patients with schizophrenia may selectively derive cognitive enhancement by cigarette smoking in the areas of attention, working memory and information processing (Adler et al., 1993; Sacco et al., 2005). Such putative amelioration of neurocognitive deficits associated with schizophrenia may also contribute to the reduced likelihood of smoking cessation in this population which has been observed both in epidemiological (Lasser et al., 2000; Etter et al., 2004) and clinical studies (Addington et al., 1998; George et al., 2000; Evins et al., 2001; George et al., 2002; Dolan et al., 2004; Evins et al., 2005; Culhane et al., 2008; Moss et al., 2009). However, little is known about the effects of smoking status on neurocognitive deficits in schizophrenia, or of clinical characteristics that may differ between smokers and non-smokers with schizophrenia.

Neurocognitive deficits are a core feature of schizophrenia (Park and Holzman, 1992; Keefe et al., 1995) but since it is estimated from clinical samples that approximately 58-88% of patients with schizophrenia are smokers (Ziedonis and Williams, 2003; Kalman et al., 2005), the nature of the cognitive deficits which exist in non-smoking patients with this disorder is unclear. Preliminary data suggests that visuospatial working memory is worse in non-smokers with schizophrenia compared to smokers with schizophrenia but this effect did not reach statistical significance as the sample sizes were low ($n=8$ and $n=23$, respectively) (George et al., 2002). In addition, the classification of a non-smoking sample into never- and former smokers, in comparison to current smokers, would facilitate the parsing of state-versus trait-dependent effects, which may be of particular relevance to the neurocognitive endophenotypes associated with schizophrenia, as well as their relationships to treatment outcomes.

Accordingly, baseline data from studies examining the effect of smoking on cognitive function in schizophrenia was used to compare the effects of smoking status on neuropsychological performance in schizophrenia in comparison to controls without a psychiatric illness. Current (non-deprived), former (quit for at least 6 months) and never- (less than 100 cigarettes in lifetime (Hughes et al., 2000)) smokers were studied using a comprehensive battery of neuropsychological tests.

2. Methods

2.1 Subjects

The majority of the data (~70% of sample) were derived from a study of the effects of cigarette smoking on cognitive function in schizophrenia, and the role of nicotinic acetylcholine receptors, conducted at Yale University (Sacco et al., 2005; Sacco et al., 2006); the remainder of the data (~30% of the sample) were derived from a cross-sectional study of cognitive function in smokers and non-smokers with schizophrenia at the Centre for Addiction and Mental Health (CAMH), Toronto. Patients with schizophrenia were recruited from clinician referrals and flyers; controls were recruited from the community using newspaper and internet advertisements. Written informed consent was obtained from all subjects seeking participation as approved by the Human Investigation Committee at Yale University and the CAMH Research Ethics Board. All subjects were screened by trained staff members using the Structured Clinical Interview for DSM-IV (SCID) for Axis I disorders (First, 1994). Subjects with schizophrenia or schizoaffective disorder were outpatients at the time of the interview, judged to be psychiatrically stable, scored < 70 on the PANSS (Kay et al., 1987) and were on a stable dose of antipsychotic medication for at least three months prior to study enrollment. Control (CON) subjects did not meet criteria for any current Axis I disorder on the SCID. Participants who met SCID criteria for alcohol or substance abuse or dependence in the last three months, other than nicotine dependence for those assigned to the current smoker groups, were not enrolled. All enrolled subjects tested negative for illicit substances using urine toxicology (Medtox®; Wilmington, NC) and were not using nicotine replacement therapy.

Subjects ($N=140$) were divided into six groups: current smokers with schizophrenia ($n=38$), former smokers with schizophrenia ($n=17$), never-smokers with schizophrenia ($n=12$), control smokers ($n=31$), control former smokers ($n=16$) and control never-smokers ($n=26$). Smokers reported smoking 10 cigarettes per day, had expired breath carbon monoxide (CO) levels > 10 ppm, and scored 5 on the Fagerstrom Test of Nicotine Dependence (FTND) (Heatherton et al., 1991; Weinberger et al., 2007). Non-smoking participants were either never-smokers, reporting smoking fewer than 100 cigarettes in their lifetime (Hughes et al., 2000), or former smokers who had quit smoking at least six months prior to participation in this study. Non-smoking status was established by self-report, and biochemically verified by expired breath CO levels less than 10 ppm

2.2 Study Design

Data from participants was derived from a single baseline neuropsychological assessment. To minimize tobacco deprivation, all smoking subjects were given frequent smoke breaks, to ensure that deprivation from cigarette smoking did not exceed 30 minutes. Details of the neuropsychological test battery have been previously described (Sacco et al., 2005), but are given briefly in the sections below. The battery of neuropsychological tasks took approximately 2.5-3.0 hours to complete.

2.3 Neuropsychological Tests

2.3.1 Conners' Continuous Performance Test (CPT)—The Conners' CPT-X is designed to measure sustained attention, concentration, response inhibition and impulsivity (Conners, 1995). Subjects press the space bar as quickly as possible after each of a series of similar visual stimuli except when presented with the letter "X". Commonly reported outcome measures include: percentage of hits, percentage of commission errors, reaction time for hits (in milliseconds), a measure of overall attentiveness (d'), and hit rate reaction time standard error variability. Schizophrenia is associated with attention and response inhibition deficits when compared to non-psychiatric population (Cornblatt and Keilp, 1994; Enticott et al., 2008).

2.3.2 Stroop Color Word Test (SCWT)—The SCWT measures participants' ability to shift their perceptual set to conform to changing conditions requiring mental control, response inhibition, response flexibility and selective attention with the occurrence of perceptual interference (Stroop, 1935). Participants report the color ink in which the names of colors are printed. The difference in response time (milliseconds) when the ink is a different color than the color name compared to a response time when the ink is the color named, is called "Stroop Interference" (SI). Facilitation occurs when the word is the same as the print color. Other common outcome measurements are neutral, congruent, and incongruent reaction times. Performance of the SCWT is known to activate the anterior cingulate cortex (Peterson et al., 1999; Swick and Jovanovic, 2002). SI is impaired in many neuropsychiatric disorders including schizophrenia (Hepp et al., 1996; Henik and Salo, 2004).

2.3.3 Wisconsin Card Sorting Test (WCST)—Participants were administered the computerized WCST (Heaton et al., 1993). The WCST assesses executive functions including cognitive flexibility in response to feedback which is thought to relate to dorsolateral prefrontal cortex function (Goldberg and Weinberger, 1988). A total of 128 cards are presented and the test requires participants to sort the cards on the basis of the color, shape, or number of figures. The only feedback provided to the subject is whether responses are correct or incorrect. Common outcome measures are categories completed, % total errors, % perseverative errors and % non-perseverative errors. Performance on this task is known to be impaired in schizophrenia (Berman et al., 1986).

2.4 Statistical Analysis

There were three primary experimental questions: **1)** Are there baseline differences in demographic and clinical features between current, former and never-smokers in the schizophrenia or control groups? We used 2 (diagnosis) $\times$ 3 (smoking history i.e., current, former and never-smokers) ANOVAs and Chi square tests to analyze data across all study groups, and 1-way ANOVA and Chi square within diagnostic groups. **2)** Are there differences between smoking and non-smoking (combination of former and never-smokers) schizophrenia or control subjects on neuropsychological outcome measures? 2 (diagnosis) $\times$ 2 (smoking status i.e., smoker versus nonsmoker) were used to analyze data across these groups, and independent samples t-tests were used to for effects of smoking status within each diagnostic group. **3)** Are there differences amongst current, former and never-smokers

in each diagnostic group on neuropsychological outcomes? 2 (diagnosis) $\times$ 3 (smoking history i.e., current, former and never-smokers) ANOVAs were performed to examine interactive effects of these two between-subject factors. One-way ANOVAs followed by Bonferroni post hoc tests corrected for multiple comparisons were conducted to identify differences between smoking histories within each diagnostic group. ANCOVAs were employed to control for differences in IQ across smoking subgroups. All statistical analyses were conducted using SPSS version 15.0 and statistical significance was set at $p < 0.05$.

3. Results

3.1 Demographic and Clinical Characteristics

The clinical and demographic variables of the schizophrenia and control current, former and never-smoker subgroups are shown in Table 1. The six groups were similar in terms of sex but there were significant differences in race within the smoking history subgroups ($p = 0.04$). There was a significant difference in age in the control, but not schizophrenia, subgroups ($p < 0.01$); never-smokers were significantly younger than former and current smokers (p 's < 0.05). One-way ANOVAs found differences in IQ across smoking history subgroups in both the schizophrenia ($p = 0.01$) and control ($p < 0.01$) groups; IQ scores in the control current smoker control were significantly lower than the never-smoker control group and in the schizophrenia participants current smokers had a significantly lower IQ than the former smokers (p 's < 0.01). Years of education did not differ across the schizophrenia groups but did in the control group ($p < 0.01$); control current smokers had significantly less years of education than control former smokers ($p < 0.01$).

Smoking measures including cigarettes per day, expired breath CO and level of nicotine dependence measured by FTND scores did not significantly differ between schizophrenia and control smokers. Current, former and never-smokers in the schizophrenia group demonstrated similar levels of clinical symptoms as measured by the PANSS and among the subgroups there were no significant differences in the proportion of subjects with schizophrenia versus schizoaffective disorder.

3.2 Comparison of Neuropsychological Performance in Schizophrenia and Control Groups as a Function of Current Smoking Status

We compared neuropsychological task performance in control smokers ($n = 31$), control nonsmokers ($n = 42$), smokers with schizophrenia ($n = 38$) and nonsmokers with schizophrenia ($n = 29$). The data were analyzed using a two-way ANOVA for diagnosis and current smoking status. There was a main effect of diagnosis on most neuropsychological measures with poorer performance in the schizophrenia group compared controls on CPT Hit Rate (%), Hit Rate Reaction Time and Variability Index; SCWT reactions times in the incongruent, congruent and neutral conditions; and errors, perseverative errors, non-perseverative responses and categories completed on the WCST (p 's > 0.03). Smoking main effects and the diagnosis $\times$ smoking status interactions were non-significant.

t -tests were conducted to compare smokers and non-smokers within the psychiatric groups. Within the patients with schizophrenia, non-smokers had significantly slower SCWT

reaction times in the Congruent condition ($t(38)=2.24, p=0.03$). No significant differences were found between control smokers and non-smokers.

3.3 Comparison of Neuropsychological Performance in Schizophrenia and Control Groups as a Function of Smoking History

We compared neuropsychological task performance in control current smokers ($n=31$), control former smokers ($n=16$), control never-smokers ($n=26$), smokers with schizophrenia ($n=38$), former smokers with schizophrenia ($n=27$) and never-smokers with schizophrenia ($n=12$) (Table 2). The data were analyzed using a two-way ANOVA for diagnosis and smoking history. Performance was significantly impaired in the schizophrenia participants on the majority of measures compared to controls: CPT Hit Rate ($F(1,134)=12.0, p=0.001$), Commission Rate ($F(1,134)=4.94, p=0.028$), Hit Rate Reaction Time ($F(1,134)=8.35, p=0.005$) and Variability Index ($F(1,134)=17.8, p<0.001$); SCWT reaction times in the incongruent ($F(1,80)=5.39, p=0.023$), congruent ($F(1,80)=14.8, p<0.001$), and neutral conditions ($F(2,80)=9.16, p=0.003$); WCST errors ($F(1,129)=6.13, p=0.015$), perseverative errors ($F(1,129)=4.20, p=0.042$), nonperseverative responses ($F(1,129)=5.43, p=0.021$) and categories completed ($F(1,129)=6.44, p=0.012$). A significant effect of smoking history was observed on SCWT reaction times in the incongruent ($F(2,80)=3.27, p=0.043$), congruent ($F(2,80)=5.44, p=0.006$) and neutral ($F(2,80)=5.14, p=0.008$) conditions. A significant interaction between smoking history and diagnosis was found on CPT Hit Rate ($F(2,134)=6.30, p=0.002$), Commission Rate ($F(2,134)=3.42, p=0.036$), Variability Index ($F(2,134)=3.71, p=0.045$) and Attentional Index ($F(2,134)=4.95, p=0.008$); and SCWT reaction times in the Incongruent ($F(2,80)=3.51, p=0.035$) and Congruent ($F(2,80)=6.52, p=0.002$) conditions.

Inspection of Table 2 suggests that in general task performance in patients with schizophrenia was the most impaired in the never-smoker group compared to former and current smokers. However, one-way ANOVAs for the effect of smoking history on performance in the schizophrenia participants, only identified significant effects on measures of sustained attention (CPT Hit Rate: $F(2,66)=3.76, p=0.03$), processing speed (SCWT Congruent: $F(2,39)=9.04, p=0.001$; SCWT Neutral: $F(2,39)=8.92, p=0.001$) and response inhibition (SCWT Incongruent condition: $F(2,39)=4.69, p=0.02$). Bonferroni post hoc comparisons (with $\alpha=0.0167$ to correct for multiple comparisons) revealed that never-smokers had poorer performance than both current and former smokers on SCWT reaction times in the Neutral (p 's <0.01), Congruent (p 's <0.01) and Incongruent ($p=0.02$ and $p=0.04$ respectively) conditions, lower CPT Hit Rates than former smokers ($p=0.02$). Because of observed differences in IQ scores across these three schizophrenia subgroups, ANCOVAs of neuropsychological performance on these tasks were performed in order to co-vary for IQ level; all significant effects persisted after this adjustment.

Analysis of neuropsychological performance across the three smoking history subgroups in controls identified no differences in performance on neuropsychological tests employed.

4. Discussion

There has been considerable study of neuropsychological deficits in patients with schizophrenia, but given the high prevalence of cigarette smoking in schizophrenia (Hughes et al., 1986; Kalman et al., 2005) there has been surprisingly little investigation of cross-sectional differences in neurocognitive function in smoking versus non-smoking patients with schizophrenia. We recently demonstrated that smoking status modifies pre-pulse inhibition (PPI) deficits, a neurophysiological measure of information processing, in schizophrenia; PPI deficits were present in nonsmokers with schizophrenia but in smokers with schizophrenia, studied under satiated conditions, PPI levels were elevated to that of non-psychiatric control smokers (Woznica et al., 2009). To our knowledge, the present study is the first study to report on cross-sectional differences in neuropsychological performance in schizophrenia as a function of smoking status or smoking history. Taken together, our results suggest smoking status (i.e., smoker versus non-smoker) only has a significant effect on neuropsychological deficits in patients with schizophrenia in the area of processing speed (SCWT Congruent and Neutral reaction times). However, when non-smokers were subdivided into former and never-smokers, a deficit in task performance in never-smokers with schizophrenia was identified across a range of measures. Selected neuropsychological measures such as sustained attention (CPT Hit Rate), processing speed (SCWT Congruent and Neutral reaction times) and response inhibition (SCWT Incongruent reaction times) were significantly impaired in never-smokers compared to current and former smokers, even after adjusting for IQ differences across schizophrenia subgroups. Such differences in neuropsychological performance as a function of smoking status and smoking history were not observed in the non-psychiatric controls.

Our results may suggest that never-smokers with schizophrenia have the most severe deficits in neuropsychological performance, and that some of these impairments, are likely to be independent of smoking exposure given our historical and biochemical verifications of non-smoking status. Therefore, we suggest that patients with schizophrenia who do not initiate (and maintain) smoking behavior may constitute a subgroup of patients with more severe impairment of neuropsychological task performance. Accordingly, this group of never-smoking patients with schizophrenia may thus represent a specific subtype of the illness with more severe neuropsychological impairment. This has implications for our understanding of cigarette smoking effects on endophenotypes associated with this disorder, and suggest that the presence or absence of cigarette smoking may have prognostic implications for the severity of neuropsychological deficits, and possibly treatment and rehabilitative outcomes in patients with schizophrenia. Whether these subjects would obtain cognitive enhancement from nicotine or nicotine-like pharmacotherapies is an open question. Studies in non-smokers with schizophrenia (including former and never-smokers) have reported that the cognitive deficits in these patients are insensitive to treatment with the nicotinic acetylcholine receptor (nAChR) antagonist mecamylamine (Sacco et al., 2006) but that nicotine replacement therapy can improve some aspects of cognitive performance (CPT, SCWT and episodic memory) (Barr et al., 2008; Jubelt et al., 2008). The cholinesterase inhibitor galantamine (Dyer et al., 2008; Sacco et al., 2008) and DMXB-A, a partial $\alpha 7$ -

nAChR agonist, had limited pro-cognitive effects in non-smokers with schizophrenia (Olincy et al., 2006; Freedman et al., 2008)

As is commonly observed, IQ and years of education were lower in the schizophrenia group compared to controls (Kondel et al., 2003). Within the control group, current smokers had both the lowest level of education and the lowest IQ, consistent with previous reports in the literature on the interaction between cigarette smoking, IQ and education (Hemmingsson et al., 2008; Weiser et al., 2010). It is noteworthy that the IQ of the control smokers (ca. 100) falls directly within expectation of a naturally occurring control group, and that all control groups fall within one standard deviation of the average. Lower IQ scores were also observed in current smokers with schizophrenia as compared to the former smokers with schizophrenia. This observation argues against there being a contribution of IQ to the observed group differences in task performance as never-smokers, not current smokers, generally had the greatest deficiencies in task performance of the three schizophrenia subgroups. Furthermore, ANCOVAs controlling for group differences in IQ across these schizophrenia subgroups ruled out a contribution of IQ disparities between subgroups to the observed differences in neuropsychological performance. Interestingly, we and other laboratories have observed that patients who quit smoking have better prefrontal cortical neuropsychological performance (assessed by the CPT, visuospatial working memory task, WCST, and Trail Making Test B) than those who did not quit smoking (Dolan et al., 2004; Culhane et al., 2008; Moss et al., 2009). While our moderately small sample size in the schizophrenia former groups ($n=17$) may not have allowed for detection of significant differences versus the current smokers, task performance on several CPT and SCWT measures tended to be better in former smoker groups than in the other two groups.

In summary, we compared patients with schizophrenia and controls as a function of smoking status (smokers and non-smokers) and smoking history (current, former and never-smokers) on a broad array of neuropsychological outcome measures. While there were clear deficits in all subgroups of patients with schizophrenia compared to controls, the deficits in performance, predominantly in the areas of sustained attention, processing speed and response inhibition, were the most pronounced in non-smokers with schizophrenia and in particular in the never-smoking subgroup. Our findings may suggest that never-smoking patients with schizophrenia may constitute a subtype of schizophrenia associated with more severe neurocognitive dysfunction. Since treatment and rehabilitative outcomes in schizophrenia have been associated with severity of baseline neurocognitive impairment (Green, 1996), smoking status should be considered as a potential predictor of treatment outcome in future studies in schizophrenia. However, given our small subgroup sample sizes and that the data are derived from a secondary analysis of baseline data from larger prospective laboratory studies of the effects of smoking and cognition in schizophrenia (Sacco et al., 2005; Sacco et al., 2006) our findings should be considered preliminary and in need of replication using larger samples.

Acknowledgements

This work was supported in part by grants K02-DA-16611 and R01-DA-14039 from the National Institute on Drug Abuse (to Dr. George), Young Investigator (to Dr. Sacco) and Independent Investigator Awards from the National Alliance for Research on Schizophrenia and Depression (to Dr. George), an Endowed Chair in Addiction Psychiatry

from University of Toronto (to Dr. George) and an Ontario Mental Health Foundation operating grant (to Drs. George, Wing and Bacher).

We thank Angelo Termine, Jennifer Vessicchio, LCSW, Melissa Dudas, Aisha Seyal, Taryn Allen, Erin Reutenauer and Becky Wu for their assistance with study procedures.

References

- Addington J, El-Guebaly N, Campbell W, Hodgins DC, Addington D. Smoking cessation treatment for patients with schizophrenia. *American Journal of Psychiatry*. 1998; 155:974–976. [PubMed: 9659869]
- Adler LE, Hoffer LD, Wiser A, Freedman R. Normalization of auditory physiology by cigarette smoking in schizophrenic patients. *American Journal of Psychiatry*. 1993; 150:1856–1861. [PubMed: 8238642]
- Barr RS, Culhane MA, Jubelt LE, Mufti RS, Dyer MA, Weiss AP, Deckersbach T, Kelly JF, Freudenreich O, Goff DC, Evins AE. The effects of transdermal nicotine on cognition in nonsmokers with schizophrenia and nonpsychiatric controls. *Neuropsychopharmacology*. 2008; 33:480–490. [PubMed: 17443126]
- Berman KF, Zec RF, Weinberger DR. Physiological dysfunction of the dorsolateral prefrontal cortex in schizophrenia. II. Role of neuroleptic treatment, attention and mental effort. *Archives of General Psychiatry*. 1986; 43:126–135. [PubMed: 2868701]
- Connors, CK. The Continuous Performance Test (CPT). Psychological Assessment Resources, Inc.; Odessa, FL: 1995.
- Cornblatt BA, Keilp JG. Impaired attention, genetics, and the pathophysiology of schizophrenia. *Schizophrenia Bulletin*. 1994; 20:31–46. [PubMed: 8197420]
- Culhane MA, Schoenfeld DA, Barr RS, Cather C, Deckersbach T, Freudenreich O, Goff DC, Rigotti NA, Evins AE. Predictors of early abstinence in smokers with schizophrenia. *Journal of Clinical Psychiatry*. 2008; 69:1743–1750. [PubMed: 19026259]
- Dalack GW, Healy DJ, Meador-Woodruff JH. Nicotine dependence and schizophrenia: clinical phenomenon and laboratory findings. *American Journal of Psychiatry*. 1998; 155:1490–1501. [PubMed: 9812108]
- Depatie L, Driscoll GA, Holahan A-LV, Atkinson V, Thavundayil JX, Kin N. N. Ying, Samarthji L. Nicotine and behavioral markers of risk for schizophrenia: a double-blind, placebo-controlled, cross-over study. *Neuropsychopharmacology*. 2002; 27:1056–1070. [PubMed: 12464463]
- Dolan SL, Sacco KA, Termine A, Seyal AA, Dudas MM, Vessicchio JC, Wexler BE, George TP. Neuropsychological deficits are associated with smoking cessation treatment failure in patients with schizophrenia. *Schizophrenia Research*. 2004; 70:263–275. [PubMed: 15329302]
- Dyer MA, Freudenreich O, Culhane MA, Pachas GN, Deckersbach T, Murphy E, Goff DC, Evins AE. High-dose galantamine augmentation inferior to placebo on attention, inhibitory control and working memory performance in nonsmokers with schizophrenia. *Schizophrenia Research*. 2008; 102:88–95. [PubMed: 18325740]
- Enticott PG, Ogloff JR, Bradshaw JL. Response inhibition and impulsivity in schizophrenia. *Psychiatry Research*. 2008; 157:251–254. [PubMed: 17916385]
- Etter M, Mohr S, Garin C, Etter JF. Stages of change in smokers with schizophrenia or schizoaffective disorder and in the general population. *Schizophrenia Bulletin*. 2004; 30:459–468. [PubMed: 15279060]
- Evins AE, Cather C, Deckerbach T, Freudenreich O, Culhane MA, Olm-Shipman CM, Henderson DC, Schoenfeld DA, Goff DC, Rigotti NA. A double-blind placebo-controlled trial of bupropion sustained-release for smoking cessation in schizophrenia. *Journal of Clinical Psychopharmacology*. 2005; 25:218–225. [PubMed: 15876899]
- Evins AE, Mays VK, Rigotti NA, Tisdale T, Cather C, Goff DC. A pilot trial of bupropion added to cognitive behavioral therapy for smoking cessation in schizophrenia. *Nicotine & Tobacco Research*. 2001; 3:397–403. [PubMed: 11694208]
- First, MB. *Diagnostic and Statistical Manual of Mental Disorders*. 4th edition. American Psychiatric Association; Washington, D.C.: 1994. (DSM-IV)

- Freedman R, Olincy A, Buchanan RW, Harris JG, Gold JM, Johnson L, Allensworth D, Guzman-Bonilla A, Clement B, Ball MP, Kutnick J, Pender V, Martin LF, Stevens KE, Wagner BD, Zerbe GO, Soti F, Kem WR. Initial phase 2 trial of a nicotinic agonist in schizophrenia. *American Journal of Psychiatry*. 2008; 165:931–936. [PubMed: 18676593]
- George TP, Vessicchio JC, Termine A, Bregartner TA, Feingold A, Rounsaville BJ, Kosten TR. A placebo-controlled study of bupropion for smoking cessation in schizophrenia. *Biological Psychiatry*. 2002; 52:53–61. [PubMed: 12079730]
- George TP, Vessicchio JC, Termine A, Sahady DM, Head CA, Pepper WT, Kosten TR, Wexler BE. Effects of smoking abstinence on visuospatial working memory function in schizophrenia. *Neuropsychopharmacology*. 2002; 26:75–85. [PubMed: 11751034]
- George TP, Zeidonis DM, Feingold A, Pepper WT, Satterburg CA, Winkel J, Rounsaville BJ, Kosten TR. Nicotine transdermal patch and atypical antipsychotic medications for smoking cessation in schizophrenia. *American Journal of Psychiatry*. 2000; 157:1835–1842. [PubMed: 11058482]
- Goldberg TF, Weinberger DR. Probing prefrontal function in schizophrenia with neuropsychological paradigms. *Schizophrenia Research*. 1988; 14:179–183.
- Green MF. What are the functional consequences of neurocognitive deficits in schizophrenia? *American Journal of Psychiatry*. 1996; 153:321–330.
- Heatherton TF, Kozlowski LT, Frecker RC, Fagerstrom KO. The Fagerstrom Test for Nicotine Dependence: a revision of the Fagerstrom Tolerance Questionnaire. *British Journal of Addiction*. 1991; 86:1119–1127. [PubMed: 1932883]
- Heaton, RK., Chelune, GJ., Talley, JL., Kay, GG., Curtiss, G. Wisconsin Card Sorting Test. Manual. Psychological Assessment Resources; Odessa, FL: 1993.
- Hemmingson T, Kriebel D, Melin B, Allebeck P, Lundberg I. How does IQ affect onset of smoking and cessation of smoking—linking the Swedish 1969 conscription cohort to the Swedish survey of living conditions. *Psychosomatic Medicine*. 2008; 70:805–810. [PubMed: 18606723]
- Henik A, Salo R. Schizophrenia and the stroop effect. *Behavioural and Cognitive Neuroscience Reviews*. 2004; 3:42–59.
- Hepp H, Maier S, Hermle L, Spitzer M. The stroop effect in schizophrenic patients. *Schizophrenia Research*. 1996; 22:187–195. [PubMed: 9000316]
- Hughes JR, Hatsukami DK, Mitchell JE, Dahlgreen LA. Prevalence of smoking among psychiatric outpatients. *American Journal of Psychiatry*. 1986; 143:993–997. [PubMed: 3487983]
- Hughes JR, Rose GL, Callas PW. Do former smokers respond to nicotine differently from never smokers? A pilot study. *Nicotine & Tobacco Research*. 2000; 2:255–262. [PubMed: 11082826]
- Jubelt LE, Barr RS, Goff DC, Logvinenko T, Weiss AP, Evins AE. Effects of transdermal nicotine on episodic memory in non-smokers with and without schizophrenia. *Psychopharmacology*. 2008; 199:89–98. [PubMed: 18548234]
- Kalman D, Morrisette SB, George TP. Co-morbidity of smoking with psychiatric and substance use disorders. *American Journal on Addictions*. 2005; 14:106–123. [PubMed: 16019961]
- Kay SR, Fiszbein A, Opler L. The positive and negative syndrome scale for schizophrenia. *Schizophrenia Bulletin*. 1987; 13:261–276. [PubMed: 3616518]
- Keefe RSE, Roitman S. E. Lees, Harvey PD, Blum CS, DuPre RL, Prieto RM, Davidson M, Davis KL. A pen-and-paper human analogue of a monkey prefrontal cortex activation task: spatial working memory in patients with schizophrenia. *Schizophrenia Research*. 1995; 17:25–33. [PubMed: 8541247]
- Kondel TK, Mortimer AM, Leeson VC, Laws KR, Hirsch SR. Intellectual differences between schizophrenic patients and normal controls across the adult lifespan. *Journal of Clinical and Experimental Neuropsychology*. 2003; 25:1045–1056. [PubMed: 14566578]
- Lasser K, Boyd JW, Woolhandler S, Himmelstein DU, McCormick D, Bor DH. Smoking and mental illness: a population-based prevalence study. *Journal of the American Medical Association*. 2000; 284:2606–2610. [PubMed: 11086367]
- Levin ED, Wilson W, Rose J, McEvoy J. Nicotine-haloperidol interactions and cognitive performance in schizophrenics. *Neuropsychopharmacology*. 1996:15.

- Moss TG, Sacco KA, Allen TM, Weinberger AH, Vessicchio JC, George TP. Prefrontal cognitive dysfunction is associated with tobacco dependence treatment failure in smokers with schizophrenia. *Drug & Alcohol Dependence*. 2009; 104:94–99. [PubMed: 19447570]
- Olinicy A, Harris JG, Johnson LL, Pender V, Kongs S, Allensworth D, Ellis J, Zerbe GO, Leonard S, Stevens KE, Stevens JO, Martin L, Adler LE, Soti F, Kem WR, Freedman R. Proof-of-Concept Trial of an alpha-7 Nicotinic Agonist in Schizophrenia. *Archives of General Psychiatry*. 2006; 63:630–638. [PubMed: 16754836]
- Park S, Holzman PS. Schizophrenics show spatial working memory deficits. *Archives of General Psychiatry*. 1992; 49:975–982. [PubMed: 1449384]
- Peterson BS, Skudlarski P, Gatenby JC, Zhang H, Anderson AW, Gore JC. An fMRI study of Stroop word-color interference: evidence for cingulate subregions subserving multiple distributed attentional systems. *Biological Psychiatry*. 1999; 45:1237–1258. [PubMed: 10349031]
- Sacco KA, Creeden CL, Reutenauer EL, George TP. Effects of galantamine on neurocognitive deficits in smokers and non-smokers with schizophrenia. *Schizophrenia Research*. 2008; 103:326–327. [PubMed: 18571377]
- Sacco KA, Termine A, Dudas MM, Seyal AA, Allen TM, Vessicchio JC, Wexler BE, George TP. Neuropsychological Deficits in Non-Smokers with Schizophrenia: Effect of a Nicotinic Antagonist. *Schizophrenia Research*. 2006; 85:213–221. [PubMed: 16678390]
- Sacco KA, Termine A, Seyal AA, Dudas MM, Vessicchio JC, Krishnan-Sarin S, Jatlow PI, Wexler BE, George TP. Effects of cigarette smoking function on spatial working memory and attentional function in schizophrenia: Involvement of nicotinic receptor mechanisms. *Archives of General Psychiatry*. 2005; 62:649–659. [PubMed: 15939842]
- Smith RC, Singh A, Infante M, Khandat A, Kloos A. Effects of cigarette smoking and nicotine nasal spray on psychiatric symptoms and cognition in schizophrenia. *Neuropsychopharmacology*. 2002; 27:479–497. [PubMed: 12225705]
- Smith RC, Warner-Cohen J, Matute M, Butler E, Kelly E, Vaidhyathanaswamy S, Khan A. Effects of nicotine nasal spray on cognitive function in schizophrenia. *Neuropsychopharmacology*. 2006; 31:637–643. [PubMed: 16160711]
- Stroop JR. Studies in interference in serial verbal reaction. *Journal of Experimental Psychology*. 1935; 18:643–662.
- Swick D, Jovanovic J. Anterior cingulate cortex and the Stroop task: neuropsychological evidence for topographic specificity. *Neuropsychologia*. 2002; 40:1240–1253. [PubMed: 11931927]
- Weinberger AH, Reutenauer EL, Allen TM, Termine A, Vessicchio JC, Sacco KA, Easton CJ, McKee SA, George TP. The Reliability and Internal Consistency of the Fagerstrom Test for Nicotine Dependence, Minnesota Nicotine Withdrawal Scale and Tiffany Questionnaire for Smoking Urges in Cigarette Smokers with and without Schizophrenia. *Drug & Alcohol Dependence*. 2007; 86:278–282. [PubMed: 16876968]
- Weiser M, Zarka S, Werbeloff N, Kravitz E, Lubin G. Cognitive test scores in male adolescent cigarette smokers compared to non-smokers: a population-based study. *Addiction*. 2010; 105:358–363. [PubMed: 19919595]
- Woznica AA, Sacco KA, George TP. Prepulse inhibition in patients with schizophrenia and controls: Effects of smoking status. *Schizophrenia Research*. 2009; 112:86–90. [PubMed: 19423298]
- Ziedonis DM, Williams JM. Management of smoking in people with psychiatric disorders. *Current Opinion in Psychiatry*. 2003; 16:305–315.

Table 1

Demographic and Clinical Data of Current, Former, and Never-Smokers as a Function of Psychiatric Diagnosis.

	SZ Current Smokers (n=38)	SZ Former Smokers (n=17)	SZ Never- Smokers (n=12)	Smoking History Effect*		CON Current Smokers (n=31)	CON Former Smokers (n=16)	CON Never- Smokers (n=26)	Smoking History Effect*		Diagnosis Effect*		Smoking History Effect*		Diagnosis × Smoking Interaction*	
				F or χ^2 (df)	p				F or χ^2 (df)	p	F, χ^2 or t (df)	p	F or χ^2 (df)	p	F (df)	P
Age	41.9±9.4	42.6±10.2	42.5±9.1	0.03 (2,63)	0.97	42.2±9.8	45.7±12.9	34.0±12.7 ^{b,c}	5.8 (2,67)	<0.01	0.7 (1,130)	0.40	2.6 (2,130)	0.08	2.7 (2,130)	0.07
Sex																
Male	24	9	8	0.7 (2)	0.70	14	7	15	1.1 (2)	0.56	2.0 (1)	0.16	1.0 (2)	0.60	-	-
Female	14	8	4			17	9	11								
Race																
Caucasian	19	14	6	6.8 (6)	0.34	21	14	14	8.7 (6)	0.19	8.0 (3)	0.11	13.2 (6)	0.04	-	-
African-American	14	3	5			7	1	4								
Hispanic	2	0	0			1	0	2								
Other	3	0	1			2	1	6								
Education (years)	12.2±2.6	13.6±4.3	12.8±1.9	1.1 (2,63)	0.33	14.0±0.5 ^c	16.4±0.6	15.7±0.5	5.2 (2,69)	<0.01	23.1 (1,132)	<0.01	5.2 (2,132)	0.01	0.6 (2,132)	0.56
Estimated IQ	85.4±12.12 ^a	99.8±13.4	91.0±12.2	7.6 (2,61)	<0.01	99.6±13.0 ^d	105.7±13.7	111.7±6.8	7.6 (2,65)	<0.01	37.3 (1,126)	<0.01	10.3 (2,126)	0.01	3.1 (2,136)	0.05
Cigarettes per day	20.7±10.5	-	-	-	-	17.7±7.6	-	-	-	-	1.3 (62)	0.20	-	-	-	-
Expired CO (ppm)	25.5±12.6	1.6±1.6	1.2±0.8	-	-	20.2±7.8	0.9±1.0	0.8±0.8	-	-	2.0 (67)	0.05	-	-	-	-
FTND	6.8±1.4	-	-	-	-	7.1±2.0	-	-	-	-	0.5 (27)	0.61	-	-	-	-
PANSS Positive	13.4±2.7	13.6±3.1	13.9±2.4	0.3 (2,64)	0.74	-	-	-	-	-	-	-	-	-	-	-
PANSS Negative	14.0±3.1	14.2±2.9	13.9±2.2	.05 (2,64)	0.95	-	-	-	-	-	-	-	-	-	-	-
PANSS General	26.7±5.4	29.8±4.7	27.7 ±2.1	2.7 (2,64)	0.07	-	-	-	-	-	-	-	-	-	-	-
PANSS Total	54.1±9.1	57.6±9.4	55.2±3.7	1.2 (2,64)	0.30	-	-	-	-	-	-	-	-	-	-	-
Diagnosis																
Schizophrenia	26	8	5	3.2 (2)	0.20	-	-	-	-	-	-	-	-	-	-	-
Schizoaffective	12	9	6													

Psychiatry Res. Author manuscript; available in PMC 2017 March 1.

Abbreviations: CO, carbon monoxide; FTND, Fagerstrom Test of Nicotine Dependence; PANSS, Positive and Negative Symptom Scale; SZ, schizophrenia; CON, control.

* Analysis of Variance, Independent t-tests and Chi-square Tests were used for continuous and categorical data respectively.

Significant differences in performance compared to other subgroups (Bonferroni post hoc tests corrected for multiple comparisons, $p<0.05$) are represented by:

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

^a compared to the SZ former smokers;
^b compared to the CON current smokers;
^c compared to CON former smokers;
^d compared to CON never-smokers

Table 2

Neuropsychological Performance of Current, Former and Never-Smokers as a Function of Psychiatric Diagnosis.

Neuropsychological Outcome Measure	SZ Current Smokers (n=38)	SZ Former Smokers (n=17)	SZ Never-Smokers (n=12)	Smoking History Effect (1-way ANOVA)		CON Current Smokers (n=31)	CON Former Smokers (n=16)	CON Never-Smokers (n=26)	Smoking History Effect (1-way ANOVA)		Diagnosis Effect (2-way ANOVA)		Smoking History Effect (2-way ANOVA)		Diagnosis × Smoking History Interaction	
				F (df)	p				F (df)	p	F (df)	p	F (df)	p	F (df)	p
CPT Hit Rate (%)	98.0±2.6	^a 99.0±1.0	96.1±4.6	3.8 (2,64)	0.03	99.1±1.0	98.6±2.1	99.4±0.8	2.1 (2,70)	0.13	12.0 (1,134)	.001	2.2 (2,134)	0.12	6.3 (2,134)	<0.01
CPT Commission Rate (%)	28.6±14.6	25.8±18.1	39.1±27.2	2.0 (2,64)	0.14	26.2±18.7	26.4±15.7	19.8±13.1	1.3 (2,70)	0.27	4.9 (1,132)	0.03	0.3 (2,132)	0.73	3.4 (2,132)	0.04
CPT Hit Rate Reaction Time (ms)	422.4±92.3	441.8±90.2	441.8±74.2	0.4 (2,64)	0.68	373.2±82.5	424.7±84.8	371.4±84.4	2.5 (2,70)	0.09	8.4 (1,134)	<0.01	1.9 (2,134)	0.15	0.8 (2,134)	0.45
CPT Variability Index	13.7±10.6	10.6±5.2	19.2±17.6	2.1 (2,64)	0.13	8.3±5.9	8.5±4.6	6.6±5.7	0.8 (2,70)	0.45	17.8 (1,134)	<0.01	1.3 (2,134)	0.29	3.2 (2,134)	0.05
CPT Attentional Index	2.1±1.1	2.8±1.1	1.9±1.5	2.4 (2,64)	0.10	3.0±1.3	1.9±1.6	2.6±1.4	2.8 (2,70)	0.07	1.1 (1,134)	0.29	0.6 (2,134)	0.57	4.9 (2,134)	<0.01
SCWT Incongruent (ms)	^a 1179±468	^a 1040±106	1855±1050	4.7 (2,37)	0.02	1054±450	1006±300	1027±603	0.02 (2,43)	0.97	5.4 (1,80)	0.02	3.3 (2,80)	0.04	5.4 (2,80)	0.04
SCWT Congruent (ms)	^a 894±249	^a 859±78	1392±512	9.0 (2,37)	<0.01	780±203	739±155	750±396	0.09 (2,43)	0.92	14.8 (1,80)	<0.01	5.4 (2,80)	<0.01	6.5 (2,80)	<0.01
SCWT Neutral (ms)	^a 914±190	^a 865±133	1314±422	8.9 (2,36)	<0.01	744±234	765±147	843±530	0.3 (2,44)	0.68	9.2 (1,80)	<0.01	5.1 (2,80)	0.01	2.0 (2,80)	0.14
SCWT Interference (ms)	286±374	130±114	448±869	0.8 (2,37)	0.46	306±347	162±128	236±423	0.4 (2,41)	0.75	0.2 (1,80)	0.64	0.8 (2,80)	0.46	0.6 (2,80)	0.57
SCWT Facilitation	7.6±185	6.6±124	22.0±294	0.01 (2,36)	0.98	39.6±249	128±222	55.6±207	0.02 (2,41)	0.69	1.1 (1,80)	0.29	0.2 (2,80)	0.83	0.2 (2,80)	0.82
WCST Errors (%)	24.9±15.5	30.2±17.5	30.3±18.2	0.8 (2,60)	0.46	20.5±17.1	25.0±20.7	17.3±10.0	1.1 (2,69)	0.33	6.1 (1,129)	0.02	0.9 (2,129)	0.40	0.8 (2,129)	0.45
WCST Perseverative Errors (%)	14.4±11.2	16.1±10.1	18.8±17.3	0.6 (2,60)	0.55	11.4±11.0	15.2±16.0	9.2±5.4	1.6 (2,69)	0.24	4.2 (1,129)	0.04	0.6 (2,129)	0.57	0.3 (2,129)	0.28
WCST Perseverative Responses	13.9±9.9	18.1±12.3	22.6±23.6	1.8 (2,60)	0.17	12.3±14.5	17.6±20.3	9.5±5.9	1.5 (2,67)	0.21	3.6 (1,125)	0.06	1.3 (2,125)	0.28	2.1 (2,125)	0.12
WCST Nonperseverative Responses (%)	10.5±6.7	14.3±8.8	11.3±5.7	1.4 (2,60)	0.25	9.0±8.2	9.9±6.5	8.0±5.5	0.4 (2,69)	0.69	5.4 (1,129)	0.02	1.3 (2,129)	0.27	0.5 (1,129)	0.63
WCST Categories Completed	4.7±2.2	4.0±2.7	4.3±2.2	0.5 (2,60)	0.60	5.2±1.7	4.9±2.1	5.8±1.0	1.9 (2,69)	0.16	6.4 (1,129)	0.01	0.9 (2,129)	0.40	0.8 (2,129)	0.47

Abbreviations: CPT, Continuous Performance Test; SCWT, Stroop Color Word Test; WCST, Wisconsin Card Sorting Task; SZ, Schizophrenia; CON, Control.

^a represents a significant difference in performance compared to the SZ never-smoker subgroup (Bonferroni post hoc tests corrected for multiple comparisons, $p < 0.04$)