Published in final edited form as:

J Phys Chem Lett. 2015 June 18; 6(12): 2388–2392. doi:10.1021/acs.jpcclett.5b00913.

Radical reactivity in the condensed phase: Intermolecular versus intramolecular reactions of alkoxy radicals

Anthony J. Carrasquillo¹, Kelly E. Daumit¹, and Jesse H. Kroll^{*,1,2}

¹Department of Civil and Environmental Engineering, Massachusetts Institute of Technology, Cambridge, MA

²Department of Chemical Engineering, Massachusetts Institute of Technology, Cambridge, MA

Abstract

Condensed-phase alkoxy (RO) radicals can undergo unimolecular (e.g., intramolecular H-atom abstraction) reactions as well as bimolecular (intermolecular H-atom abstraction) reactions, though the competition between these two channels is not well constrained. Here we examine this branching, by generating RO radicals from the photolysis of a large alkyl nitrite ($C_{20}H_{41}ONO$) in hexanes, and nebulizing the mixture into an aerosol mass spectrometer for analysis. Product ions associated with unimolecular (isomerization) reactions were observed to increase upon photolysis. However, no formation of the C_{20} alcohol ($C_{20}H_{41}OH$, the expected product from $RO + RH$ reactions) was observed, suggesting that bimolecular reactions are at most a minor channel for this condensed-phase system (involving saturated hydrocarbons). This result, combined with previous studies of condensed-phase RO radicals carried out at higher concentrations, suggests that when 1,5-H-atom abstraction reactions are facile (i.e., in which a 1,5-H-atom shift from a secondary or tertiary carbon can occur), this channel will dominate over bimolecular reactions.

Graphical abstract

*Corresponding Author: Jesse H. Kroll. jhkroll@mit.edu.

Supporting Information. High-mass AMS spectrum of the ROH alcohol?, ion time series for octyl-2-dodecyl-nitrite photolysis in CCl_4 and eicosanyl-1-nitrite, hexacosanyl-2-nitrite, and octacosanyl-10-nitrite photolysis in hexanes, the protocol for RONO preparation, and the method and results from GC-MS analysis of the reaction mixture are given in the Supporting Information. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Keywords

Alkyl nitrites; unimolecular; bimolecular; organic aerosol; combustion

Condensed-phase organic radicals are ubiquitous in both natural and engineered chemical systems, and play central roles in key processes such as the degradation of pollutants,¹ weathering of materials,² combustion reactions,³ and the oxidative evolution of atmospheric organic aerosol particles.⁴ Such organic radicals, which include alkyl (R), alkoxy (RO) and alkylperoxy (RO₂) species, are highly reactive and often can undergo several different reactions. Most mechanistic studies of radical reactions have focused on the gas phase, which (under most conditions) limits the available reactions to unimolecular processes, or bimolecular reactions with major inorganic species (O₂, NO_x, etc.).^{3,5,6} However, the condensed phase represents a more complex molecular environment, since the high concentrations may facilitate bimolecular reactions with other organic species (molecules or radicals).^{7–12} Condensed-phase radicals can therefore exhibit fundamentally different reactivity than gas-phase radicals; however, these generally have received far less study, leading to major uncertainties in the radical chemistry of important multiphase systems (e.g., atmospheric aerosol and cloud droplets).

Here we utilize a novel approach for studying the reactions of large organic radicals in the condensed phase, via nebulization of the reaction mixture into an aerosol mass spectrometer. This technique, used previously for the study of aqueous-phase oxidation reactions, allows for the sensitive, real-time measurement of radical reaction kinetics and products, and avoids many of the analytical challenges associated with collecting aliquots of the mixture for offline analysis.^{13,14} Our specific focus is the chemistry of alkoxy radicals (RO), key intermediates in the atmospheric oxidation of organic compounds, as well as combustion processes.^{3,13–16} RO radicals can form within the condensed phase (e.g., in atmospheric particles or droplets, fuel mixtures, etc.) by several channels: self- and cross-reactions of RO₂ radicals, the reaction of RO₂ with NO, and the direct photolysis or thermal decomposition of peroxide (ROOH, ROOR) species. In the gas phase, RO radicals react via one of three channels (Scheme 1): (**r**₁) isomerization, a 1,5-intramolecular H-atom abstraction reaction that increases the degree of functionalization; (**r**₂) reaction with O₂ to form a carbonyl and HO₂; or (**r**₃) C-C bond scission (fragmentation) to form two smaller species (a carbonyl and an alkyl radical).¹⁸ In the condensed organic phase, an additional channel (**r**₄) is available, the intermolecular (bimolecular) abstraction of H atoms from other organic species (RO + RH), forming an alcohol (ROH) and an alkyl radical.

Bimolecular reactions of RO (**r**₄) have shown to be an important channel for small (C₁–C₄) RO radicals in organic solvents,¹⁹ and recently have been invoked to explain observations of alcohol formation in heterogeneous oxidation experiments of organic aerosol particles.^{8–10,20–22} On the other hand, studies of larger condensed-phase radicals indicate that reaction **r**₄ becomes less important once RO radicals reach the size at which unimolecular isomerization is facile (i.e., in which a 1,5 H-atom shift from a secondary or tertiary carbon can occur), with decreases in observed ROH yields and increases in the yields of isomerization products.²³ While such studies suggest the dominance of reaction **r**₁

over **r**₄ for large RO radicals, they were carried out under oxygen-free conditions, with high concentrations of radicals, and so results may not be representative of radical reactivity in the atmosphere. Thus the relative importance of unimolecular and bimolecular reactions of large RO radicals is poorly constrained under relevant atmospheric conditions. Indeed, literature recommendations for the relative rates of unimolecular and bimolecular alkoxy radical reactions vary dramatically; for example one compilation of radical reaction rates indicates a fast rate of bimolecular reaction,²⁴ whereas structure-activity relationships predict it to be orders of magnitude slower than unimolecular reactions.²⁵

In our previous work, we examined the gas-phase unimolecular chemistry of alkoxy radicals, specifically the branching between isomerization (**r**₁) and fragmentation (**r**₃), using the photolysis of a series of gas-phase C₁₀ alkyl nitrites (RONO) to generate RO radicals.²⁶ Here we extend this technique to the study of the chemistry of RO radicals in the condensed phase. Large alkyl nitrites were photolyzed in hexanes (0.01-0.1% v/v) to generate condensed-phase RO radicals, and reaction kinetics and products were monitored via nebulization into a high-resolution aerosol mass spectrometer (AMS).²⁷ Described here are results for octyl-2-dodecyl-1-nitrite (C₂₀H₄₁ONO), which forms an octyl-2-dodecyl-1-oxy (C₂₀H₄₁O) radical upon photolysis. Other large alkyl nitrites (eicosanyl-1-nitrite, C₂₀H₄₁ONO, hexacosanyl-2-nitrite, C₂₆H₅₃ONO, and octacosanyl-10-nitrite, C₂₈H₅₇ONO) were also studied; results were similar to those of octyl-2-dodecyl-1-nitrite and thus are shown only in the Supplementary Information.

Upon UV irradiation of the RONO mixture, there is a dramatic change in the chemical composition of the solution mixture. The output of the atomizer (aerosol loading as measured by the AMS) increased by as much as a factor of 10 over the course of an hour. While the reason for this remains unclear, this growth likely reflects changes in the behavior of the atomized solution over the course of the experiment. In a blank experiment involving the irradiation of C₂₀H₄₁OH, (the synthetic precursor of the nitrite, as described in the Experimental Section), no chemical changes were observed, confirming that the observed chemistry arises from RONO photolysis.

Figure 1 shows the aerosol mass spectra (*m/z* 260 – 320) for a 0.025% C₂₀H₄₁ONO/hexane mixture before and after irradiation. The major ion associated with the precursor C₂₀H₄₁ONO is C₁₉H₃₉⁺ [M – CH₂ONO]⁺ at *m/z* 267.305; as expected, the intensity of this ion drops dramatically with irradiation. A second tracer ion for RONO might be C₂₀H₄₀⁺ [M – HONO]⁺ at *m/z* 280.313. However this is also a major ion ([M – H₂O]⁺) in the mass spectrum of the corresponding alcohol, ROH (C₂₀H₄₁OH) (Figure S2). ROH may be formed photochemically (**r**₄, Scheme 1) or it may instead simply be unreacted precursor from RONO synthesis; these two possibilities are discussed below.

Several large C₂₀ oxygen-containing ions increase upon irradiation, indicating that they are reaction products. The three most intense ones are C₂₀H₃₈O⁺, C₂₀H₃₉O⁺, and C₂₀H₃₉O₂⁺; similar product ions have been previously observed in studies of gas-phase alkane oxidation by Lim and Ziemann¹⁶ and in our earlier work on SOA from photolysis of gas-phase RONO species.²⁶ Based on that work, these ions correspond to the hydroxycarbonyl (C₂₀H₃₈O⁺, [M – H₂O]⁺), its cyclization product the cyclic hemiacetal (C₂₀H₃₉O⁺, [M – OH]⁺), and acetal

or hemiacetal dimers ($C_{20}H_{39}O_2^+$, $[M - C_{20}H_{41}O_2]^+$), respectively, all first-generation products of alkane oxidation. The source of a fourth major product ion, an N-containing fragment at m/z 277.276 ($C_{19}H_{35}N^+$), is unknown; it may result from fragmentation of a nitrate species on the heater or during ionization, but further study is needed.

Typical time series of the key ions in the mass spectrum, normalized to the total organic signal to account for changes in atomizer output, are shown in Figure 2. Repeat experiments gave similar results, though with some differences (up to 50%) in the relative abundances of the large oxygen-containing ions ($C_{20}H_{38}O^+$ and $C_{20}H_{39}O^+$); these may be due to small variations in the AMS vaporizer temperature but do not affect the overall conclusions of this work. Also shown are the traces for NO^+ and NO_2^+ , fragments of the RONO reactant as well as organic nitrate (RONO₂) products. Upon initiation of photolysis, the abundance of $C_{19}H_{39}^+$ decays exponentially ($k_{\text{phot}} = 3.6 \times 10^{-4} \text{ s}^{-1}$), as expected for the nitrite reactant. By contrast, $C_{20}H_{40}^+$ drops slightly and then remains fairly constant; the difference between these two traces suggests contributions by ROH, which is discussed in detail below. The NO^+ ion (a minor fragment in the RONO spectrum) also decreases, though not as rapidly as the RONO marker peak ($C_{19}H_{39}^+$), whereas the NO_2^+ ion rapidly increases. After 60 minutes the NO^+/NO_2^+ ratio levels off at ~ 15 , suggesting the formation of organic nitrates.^{28–30}

A mechanism consistent with the above observations is given in Scheme 2. Following formation, the RO radical will either fragment (**r₃**) to two smaller molecules or isomerize (**r₁**) followed by O₂ addition to form an RO₂ radical (**r₂**, reaction with O₂, is expected to be negligible based on analogous gas-phase observations).¹⁸ Products from fragmentation reactions will include species with 19 carbon atoms; these may be too volatile to remain in the aerosol phase for analysis, and/or difficult to distinguish from the C₂₀ fragments within the AMS, and so cannot be easily identified. RO₂ can react with either NO or another RO₂ to form an additional RO radical. The newly formed RO radical will “back-isomerize” and form the hydroxycarbonyl, the cyclic hemiacetal, and acetal or hemiacetal dimers.³¹ Ions corresponding to products from “forward isomerization” reactions (i.e. dihydroxycarbonyls and hydroxycyclic hemiacetals) were not observed, consistent with our previous studies of gas-phase β -substituted RO radicals.²⁶ The observation of ions corresponding to the expected products of RO isomerization – nitrates, hydroxycarbonyls, and (cyclic) hemiacetals – provides strong evidence that unimolecular (intramolecular H-atom abstraction) reactions are important reaction pathways for condensed-phase $C_{20}H_{41}O$ radicals.

The evidence for bimolecular reactions (intermolecular H-atom abstraction reactions) with hexane solvent molecules (**r₄**) is less clear. This reaction yields ROH and a C₆ species; however, only ROH is likely to be of low enough volatility to be measured by the AMS. AMS measurements of a pure ROH standard (Figure S2) confirm that the dominant high molecular weight ion is $C_{20}H_{40}^+$ ($[M - H_2O]^+$), formed from dehydration on the vaporizer or within the ionization region. This ion is present prior to photolysis, suggesting that it is also a tracer ion for the RONO, and/or that some residual ROH from the RONO synthesis was present in the reaction mixture. Its abundance decreases more slowly than that of $C_{19}H_{39}^+$ (the major RONO tracer ion), indicating at least some contribution by ROH, though

this still leaves open the possibility of ROH formation from $\mathbf{r}_4$ (which counterbalances the decrease from loss of RONO). In order to investigate the role of $\mathbf{r}_4$ as a source of ROH, the experiment was repeated using CCl_4 as a solvent instead of hexanes, precluding any possible bimolecular RO + RH reactions. Results were similar to those using hexane solvent (Figure S3), with only a slight drop in $\text{C}_{20}\text{H}_{40}^+$ ion signal upon photolysis, implying little influence of $\mathbf{r}_4$ in either solvent. Furthermore, GC-MS analysis of derivatized aliquots of the irradiated RONO/hexane system (see Supporting Information, Figures S4 and S5) showed no increase in the total abundance of the alcohol. Both results strongly suggest that any ROH present in the reaction mixture was an artifact from RONO synthesis rather than a product of the bimolecular reactions ($\mathbf{r}_4$). The lack of formation of the ROH in these experiments indicates that bimolecular H-atom abstraction reactions (RO + RH) are not competitive with the available unimolecular isomerization pathways in this system.

The present results are consistent with previous studies by Kabasakalian and co-workers of chemistry of condensed-phase long-chain alkoxy radicals, which as discussed above found products of unimolecular 1,5-H-abstraction reactions to dominate over those from bimolecular

RO + RH reactions.^{23,32} Studies that instead showed the dominance of bimolecular reactions focused almost entirely on small ($\text{C}_1 - \text{C}_4$) radicals,¹⁹ for which the 1,5-isomerization cannot occur, and for which fragmentation is known to be slow. Further, most previous studies were conducted under high concentrations (1 – 10%) of radical precursors, which may promote alkoxy disproportionation (RO + RO) reactions that form ROH and carbonyl species. Such reactions likely explain the roughly equal yields of carbonyls and ketones from such studies,²³ which are not expected from the RO + RH reaction. While some evidence for ROH formation and minor reaction products from H-atom abstraction from solvent molecules were also reported,³² their formation may have been influenced by the reaction conditions (i.e. the oxygen-free conditions and high radical concentrations in that study). The results from this work thus suggests that, under more atmospherically relevant conditions (low radical concentrations, the presence of O_2 , and room temperature), isomerization will dominate over bimolecular reactions, at least when saturated hydrocarbon molecules are present. This result is also consistent with the high entropic penalties associated with bimolecular reactions, since they involve the loss of translational rather than vibrational degrees of freedom.³³ This inferred dominance of unimolecular abstraction reactions over bimolecular ones likely holds for RO_2 autoxidation chemistry as well, recently suggested as a potentially important oxidation channel within organic aerosol.³⁴

We note that this work focuses on only one chemical system, the reactions of unfunctionalized RO radicals within a simple liquid alkane solvent, and thus our results do not necessarily imply that condensed-phase bimolecular H-atom abstraction reactions cannot compete with unimolecular reactions in all cases. In particular, when diffusion is slow (e.g., in solid matrices or “glassy” organic aerosol)⁹, or in which the surrounding molecules have readily-abstractable hydrogen atoms (e.g., oxygenates), bimolecular H-atom abstractions may be of increased importance. This is an important area of future research, and may help explain why recent laboratory studies have observed the formation of alcohols, which may have been formed from bimolecular reactions, in contrast to the present results.^{8–10,20–22}

However, alcohol species can also be formed by reactions other than $\text{RO} + \text{RH}$, such as $\text{RO}_2 + \text{RO}_2$ and $\text{RO} + \text{RO}$ disproportionation reactions. Moreover, in many cases (such as in most atmospheric organic aerosol) a given condensed-phase radical will likely be chemically similar to its surrounding molecules. There, any enhancement in bimolecular rates (for example from more readily-abstractable H atoms) will also involve a coincident enhancement in unimolecular rates. In such systems, the reactivity of alkoxy radicals in the condensed (liquid) phase is likely to be dominated by unimolecular chemistry, similar to alkoxy radical reactivity in the gas phase.

Experimental

RONO species were prepared in the laboratory via O- nitrosation of the corresponding alcohols³⁵ using the same protocol used in our previous study²⁶ but with slight modifications to the synthesis protocol for solid-phase alcohol precursors (see the Supporting Information). RONO photolysis experiments were conducted by adding 0.01 – 0.1% (v/v) $\text{C}_{20}\text{H}_{41}\text{ONO}$ in hexanes (Sigma Aldrich, 99% anhydrous) or CCl_4 (Sigma Aldrich, 99%) to a borosilicate bottle attached to a constant output atomizer (TSI, Inc.). The output from the atomizer was passed through a charcoal denuder to remove hexane solvent, resulting in particles of average volume-weighted diameter of ~ 250 nm; the flow was split to a scanning mobility particle sizer (TSI, Inc.) to measure particle size and an Aerodyne High-Resolution Aerosol Mass Spectrometer (AMS) to measure particle composition.²⁷ The AMS was run in high-resolution “W-mode” for exact mass confirmation, and time series for individual ions were constructed using the lower-resolution “V-mode” for greater sensitivity to high-MW ions. After ~ 10 minutes of atomization of the unreacted mixture in the dark, a single 40W blacklight (centered at 350 nm), positioned 2-3 cm adjacent to the reactor, was illuminated. Irradiation continued until the major high molecular weight signals stopped changing (typically within one hour). For several experiments, perfluoroheptacosane (PFH, $\text{C}_{27}\text{F}_{56}$) particles were added for improved mass calibration at $m/z > 250$. PFH particles were generated by passing air over a heated (~ 80 °C) PFH reservoir, and then into a cooling area, forming pure PFH particles that were then added to the atomizer aerosol stream and sent into the AMS.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by the National Science Foundation under grant number CHE-1012809. We thank MIT's Center for Environmental Health Sciences (supported by NIH-NIEHS Center Grant P30-ES002109) at MIT for use of their GC-MS for off-line analysis of the reaction mixture and Dr. Koli Taghizadeh for advice on GC method development.

References

1. Schwarzenbach, R., Gschwend, P., Imboden, D. Environmental Organic Chemistry. 2nd. John Wiley & Sons, Inc.; Hoboken, NJ, USA: 2003.

2. Kutz, M. Handbook of Environmental Degradation of Materials. 2nd. Elsevier Inc.; Waltham, MA, USA: 2012.
3. Curran HJ, Gaffuri P, Pitz WJ, Westbrook CK. A Comprehensive Modeling Study of n-Heptane Oxidation. *Combust Flame*. 1998; 114:149–177.
4. Kroll JH, Seinfeld JH. Chemistry of Secondary Organic Aerosol: Formation and Evolution of Low-Volatility Organics in the Atmosphere. *Atmos Environ*. 2008; 42:3593–3624.
5. Ziemann PJ, Atkinson R. Kinetics, Products, and Mechanisms of Secondary Organic Aerosol Formation. *Chem Soc Rev*. 2012; 41:6582–6605. [PubMed: 22940672]
6. Calvert, JG., Derwent, RG., Orlando, JJ., Tyndall, GS., Wallington, TJ. Mechanisms of the Atmospheric Oxidation of the Alkanes. Oxford University Press; USA: New York: 2008.
7. Smith JD, Kroll JH, Cappa CD, Che DL, Liu CL, Ahmed M, Leone SR, Worsnop DR, Wilson KR. The Heterogeneous Reaction of Hydroxyl Radicals with Sub-Micron Squalane Particles: A Model System for Understanding the Oxidative Aging of Ambient Aerosols. *Atmos Chem Phys*. 2009; 9:3209–3222.
8. Liu CL, Smith JD, Che DL, Ahmed M, Leone SR, Wilson KR. The Direct Observation of Secondary Radical Chain Chemistry in the Heterogeneous Reaction of Chlorine Atoms with Submicron Squalane Droplets. *Phys Chem Chem Phys*. 2011; 13:8993–9007. [PubMed: 21455529]
9. Ruehl CR, Nah T, Isaacman G, Worton DR, Chan AWH, Kolesar KR, Cappa CD, Goldstein AH, Wilson KR. The Influence of Molecular Structure and Aerosol Phase on the Heterogeneous Oxidation of Normal and Branched Alkanes by OH. *J Phys Chem A*. 2013; 117:3990–4000. [PubMed: 23611149]
10. Nah T, Zhang H, Worton DR, Ruehl CR, Kirk BB, Goldstein AH, Leone SR, Wilson KR. Isomeric Product Detection in the Heterogeneous Reaction of Hydroxyl Radicals with Aerosol Composed of Branched and Linear Unsaturated Organic Molecules. *J Phys Chem A*. 2014; 118:11555–11571. [PubMed: 25388092]
11. George IJ, Vlasenko A, Slowik JG, Broekhuizen K, Abbatt JPD. Heterogeneous Oxidation of Saturated Organic Aerosols by Hydroxyl Radicals: Uptake Kinetics, Condensed-Phase Products, and Particle Size Change. *Atmos Chem Phys*. 2007; 7:4187–4201.
12. George IJ, Abbatt JPD. Heterogeneous Oxidation of Atmospheric Aerosol Particles by Gas-Phase Radicals. *Nat Chem*. 2010; 2:713–722. [PubMed: 20729889]
13. Lee AKY, Herckes P, Leaitch WR, Mac Donald aM, Abbatt JPD. Aqueous OH Oxidation of Ambient Organic Aerosol and Cloud Water Organics: Formation of Highly Oxidized Products. *Geophys Res Lett*. 2011; 38:2–6.
14. Daumit KE, Carrasquillo aJ, Hunter JF, Kroll JH. Laboratory Studies of the Aqueous-Phase Oxidation of Polyols: Submicron Particles vs. Bulk Aqueous Solution. *Atmos Chem Phys Discuss*. 2014; 14:13649–13680.
15. Lim, Y Bin, Ziemann, PJ. Products and Mechanism of Secondary Organic Aerosol Formation from Reactions of n-Alkanes with OH Radicals in the Presence of NOx. *Environ Sci Technol*. 2005; 39:9229–9236. [PubMed: 16382947]
16. Lim, Y Bin, Ziemann, PJ. Chemistry of Secondary Organic Aerosol Formation from OH Radical-Initiated Reactions of Linear, Branched, and Cyclic Alkanes in the Presence of NOx. *Aerosol Sci Technol*. 2009; 43:604–619.
17. Lim YB, Ziemann PJ. Effects of Molecular Structure on Aerosol Yields from OH Radical-Initiated Reactions of Linear, Branched, and Cyclic Alkanes in the Presence of NOx. *Environ Sci Technol*. 2009; 43:2328–2334. [PubMed: 19452882]
18. Atkinson R. Rate Constants for the Atmospheric Reactions of Alkoxy Radicals: An Updated Estimation Method. *Atmos Environ*. 2007; 41:8468–8485.
19. Gray P, Williams A. The Thermochemistry And Reactivity Of Alkoxy Radicals. *Chem Rev*. 1959; 59:239–328.
20. Kolesar KR, Buffaloe G, Wilson KR, Cappa CD. OH-Initiated Heterogeneous Oxidation of Internally-Mixed Squalane and Secondary Organic Aerosol. *Environ Sci Technol*. 2014; 48:3196–3202. [PubMed: 24555558]
21. Zhang H, Ruehl CR, Chan AWH, Nah T, Worton DR, Isaacman G, Goldstein AH, Wilson KR. OH-Initiated Heterogeneous Oxidation of Cholestane: A Model System for Understanding the

- Photochemical Aging of Cyclic Alkane Aerosols. *J Phys Chem A*. 2013; 117:12449–12458. [PubMed: 24152093]
22. Chan MN, Zhang H, Goldstein AH, Wilson KR. Role of Water and Phase in the Heterogeneous Oxidation of Solid and Aqueous Succinic Acid Aerosol by Hydroxyl Radicals. *J Phys Chem C*. 2014; 118:28978–28992.
23. Kabasakalian P, Townley ER, Yudis MD. Photolysis of Nitrite Esters in Solution. III. Photochemistry of Primary, Secondary and Tertiary Alkyl Nitrites. *J Am Chem Soc*. 1962; 84:2718–2722.
24. Denisov, E., Afanas'ev, I. *Oxidation and Antioxidants in Organic Chemistry and Biology*. Taylor & Francis; Boca Raton, FL, USA: 2005.
25. Hendry DG, Mill T, Piszkiwicz L, Howard JA, Eigenmann HK. A Critical Review of H-Atom Transfer in the Liquid Phase: Chlorine Atom, Alkyl, Trichloromethyl, Alkoxy, and Alkylperoxy Radicals. *J Phys Chem Ref Data*. 1974; 3:937.
26. Carrasquillo AJ, Hunter JF, Daumit KE, Kroll JH. Secondary Organic Aerosol Formation via the Isolation of Individual Reactive Intermediates: Role of Alkoxy Radical Structure. *J Phys Chem A*. 2014; 118:8807–8816. [PubMed: 25148415]
27. DeCarlo PF, Kimmel JR, Trimborn A, Northway MJ, Jayne JT, Aiken AC, Gonin M, Fuhrer K, Horvath T, Docherty KS, et al. Field-Deployable, High-Resolution, Time-of-Flight Aerosol Mass Spectrometer. *Anal Chem*. 2006; 78:8281–8289. [PubMed: 17165817]
28. Bruns EA, Perraud V, Zelenyuk A, Ezell MJ, Johnson SN, Yu Y, Imre D, Finlayson-Pitts BJ, Alexander ML. Comparison of FTIR and Particle Mass Spectrometry for the Measurement of Particulate Organic Nitrates. *Environ Sci Technol*. 2010; 44:1056–1061. [PubMed: 20058917]
29. Fry JL, Kiendler-Scharr A, Rollins AW, Wooldridge PJ, Brown SS, Fuchs H, Dubé W, Mensah A, dal Maso M, Tillmann R, et al. Organic Nitrate and Secondary Organic Aerosol Yield from NO₃ Oxidation of β -Pinene Evaluated Using a Gas-Phase Kinetics/aerosol Partitioning Model. *Atmos Chem Phys*. 2009; 9:1431–1449.
30. Rollins AW, Kiendler-Scharr A, Fry JL, Brauers T, Brown SS, Dorn HP, Dubé WP, Fuchs H, Mensah A, Mentel TF, et al. Isoprene Oxidation by Nitrate Radical: Alkyl Nitrate and Secondary Organic Aerosol Yields. *Atmos Chem Phys*. 2009; 9:6685–6703.
31. Aimanant S, Ziemann PJ. Chemical Mechanisms of Aging of Aerosol Formed from the Reaction of n-Pentadecane with OH Radicals in the Presence of NO_x. *Aerosol Sci Technol*. 2013; 47:979–990.
32. Kabasakalian P, Townley ER. Photolysis of Nitrite Esters in Solution. I. Photochemistry of N - Octyl Nitrite. *J Am Chem Soc*. 1962; 84:2711–2716.
33. Page MI. The Energetics of Neighbouring Group Participation. *Chem Soc Rev*. 1973; 2:295.
34. Crounse JD, Nielsen LB, Jørgensen S, Kjaergaard HG, Wennberg PO. Autoxidation of Organic Compounds in the Atmosphere. *J Phys Chem Lett*. 2013; 4:3513–3520.
35. Noyes WA. Explanation of the Formation of Alkyl Nitrites in Dilute Solutions; Butyl and Amyl Nitrites. *J Am Chem Soc*. 1933; 55:3888–3889.

Figure 1. Unit mass resolution AMS spectrum (m/z 260 – 320) of $C_{20}H_{41}ONO$ in hexanes in the dark (black) and after 20 min irradiation (red, positive x-axis offset). Ion formulae were identified by high-resolution analysis, and molecular assignments are based on Lim and Ziemann.¹⁶

Figure 2.

Ion time series for a typical $C_{20}H_{41}ONO$ photolysis experiment in hexanes. All ions are normalized to total organic loading to account for changes in the atomizer output. UV lights were turned on at $t=0$ (vertical black line), after which point there is a decrease in the abundance of NO^+ and the major RONO ion ($C_{19}H_{39}^+$), and an increase in NO_2^+ and oxygenated ions ($C_{20}H_{38}O^+$, $C_{20}H_{39}O^+$, and $C_{20}H_{39}O_2^+$).

Scheme 1.

Possible reactions of gas- and condensed-phase alkoxy radicals.

Scheme 2.

Mechanism, reaction products, and ion assignments for a single C_{20} alkoxy radical in the condensed phase.