France passes "right to die" law

Fourteen months after a high profile euthanasia case led to a parliamentary commission on the subject, France has passed a law allowing terminally ill or severely injured patients the right to die.

The law, which was passed by the National Assembly on 30 November but has to be ratified by the Senate, stops short of legalising active euthanasia. The compromise follows the widely reported case of active euthanasia in which Vincent Humbert, aged 22, was seriously injured in a road accident and helped to die by his mother and a doctor on 26 September 2003 (*BMJ* 2003;327:1068).

Under the new law, terminally ill but conscious patients may refuse treatment to prolong their lives. Seriously disabled patients who are not terminally ill may also request an end to treatment. Palliative treatment must continue, however, and the doctor may give increasingly strong doses of painkillers, even if it risks shortening the patient's life.

If the patient is unconscious, the doctor must consult with other doctors and the patient's family. *Brad Spurgeon Paris*

Tighter controls are needed to root out bogus patient groups

Patient organisations need to be more tightly regulated to protect the public from bogus groups and undue influence from the pharmaceutical industry and other funding bodies, MPs heard last week.

Setting standards on transparency and accountability would help to distinguish legitimate patient groups from those set up by drug companies to promote awareness of a specific condition and drugs to treat it, said witnesses at the fourth public hearing of the health committee's inquiry into the influence of the pharmaceutical industry.

There are more than 200 national patient groups in the

United Kingdom, and many rely on funding from the pharmaceutical industry, said Melinda Letts, chair of the committee of medicines working group on patient information.

"The answer is not to ban funding from the pharmaceutical industry but to develop guidelines for what types of funding patient groups should accept," she told the committee.

Funding from any source, whether from government, charitable bodies, or drug companies, has strings attached, said MP Paul Flynn. "Knowing that they can't advertise to the public, pharmaceutical companies use patient organisations to put their case," said Mr Flynn.

Zosia Kmietowicz London

Mother accused of killing son in "fabricated illness" case

A mother fatally poisoned her ailing 9 year old son to death by putting salt in his feed while he was being treated in hospital, the prosecuting counsel told an Old Bailey jury last week.

The prosecutor, Mr Nicholas Hilliard, alleged that Petrina Stocker, aged 42, had a rare syndrome whereby a person fabricates symptoms of illness in another. He alleged that Mrs Stocker also contaminated her son's urine samples with blood. Mrs Stocker denies putting the fatal amount of salt in her son David's feed.

The jury heard that between February and August 2001 her son was treated first in Oldchurch Hospital in Romford and then the specialist children's hospital Great Ormond Street, in London.

Towards the end of his treatment, David was being fed with a specially prepared milk feed. On 18 August 2001 his health deteriorated rapidly.

"Investigations revealed extraordinarily high levels of salt in his blood," said Mr Hilliard. "Efforts were made to reduce the salt levels, but he suffered from severe swelling to the brain and died on August 20. The trial continues.

Tim Wood London

Depression affects elderly people's lives more than physical illnesses

Janice Hopkins Tanne New York

Severe depression impairs elderly patients' lives more than serious medical illnesses, and treatment of depression "has the potential to improve patients' lives in spite of other medical comorbidities," says a multicentre randomised controlled US study published in the *Annals of Family Medicine* (2004;2:555-562).

Depression is often viewed as resulting from medical illness, but it is also related to psychosocial factors, such as the death of a spouse, retirement, social isolation, and diminished income, says the study.

Treating depression in primary care can improve patients' quality of life despite their other illnesses, the authors say. "Indeed, it may be that treatment for depression can lead to more dramatic improvements in functional status, disability, and quality of life than interventions for other chronic illnesses in this age group."

The study enrolled 1801 patients aged 60 and older with major depression or dysthymia from eight US healthcare organisations. The aim was to develop a sample of depressed older patients who were to be treated under real-world conditions, comparing usual care with collaborative disease management. This study reports on the effects of depression and medical illnesses in these patients.

Lead author Dr Polly Hitchcock Noel, of the South Texas Veterans Health Care System in San Antonio, told the *BMJ* that 45% of patients who received collaborative treatment for 12 months had a 50% or better reduction in

depressive symptoms compared with only 19% of patients receiving usual care. Collaborative care included treatment by a depression care manager (a nurse or psychologist) supervised by a psychiatrist, and treatment with antidepressants or short psychotherapy. Usual care patients could use any primary care or specialty mental care services available to them.

Chronic depression affected 83% of patients, 10.6% had post-traumatic stress disorder, 21.7% had panic disorder, and 35.4% had mild cognitive impairment.

Medical problems included chronic lung disease (23.3%), hypertension (57.9%), diabetes (23.2%), arthritis (55.6%), sensory deficit (55.2%), cancer (excluding skin cancer) (10.9%), neurological disease (8.4%), heart disease (27.6%), chronic pain (56.8%), gastrointestinal disease (20.9%), and urinary or prostate disease (38.7%). Patients had a mean of 3.79 chronic diseases (standard error 0.046).

"Controlling for all other variables, depression severity was the only psychological or medical variable that was significantly associated with all four outcomes [quality of life, physical functioning, mental functioning, and disability]... As depression severity increased, quality of life and physical and mental functioning declined, while disability increased," the authors wrote, commenting that depression has a "devastating impact... on both emotional and physical functioning in older adults." □

Depression can result from psychosocial factors, such as the death of a spouse, reduced income, or social isolation