Published in final edited form as:

Prev Med. 2016 April ; 85: 60–68. doi:10.1016/j.ypmed.2015.12.017.

Effect of Appreciation for Indigenous Cultures and Exposure to Racial Insults on Alcohol and Drug use Initiation among Multiethnic Argentinean Youth

Ethel Alderete, MPH, Dr.PH.¹, Steven E. Gregorich, PhD², Madalena Monteban, PhD³, Celia P. Kaplan, Dr. PH, MA^{2,4}, Raul Mejia, MD, PhD^{5,6}, Jennifer Livaudais-Toman, PhD², and Eliseo J. Pérez-Stable, MD^{2,5}

¹Consejo Nacional de Investigaciones Científicas y Técnicas, Universidad Nacional de Jujuy, and Instituto de Ciencia y Tecnología Regional (ICTER), Argentina

²Division of General Internal Medicine, Medical Effectiveness Research Center for Diverse Populations, Department of Medicine, University of California San Francisco; Dr. Pérez-Stable is currently at National Institute of Minority Health and Health Disparities, Bethesda, MD

³Department of Anthropology, University of Georgia, Athens, GA, USA

⁴Helen Diller Family Comprehensive Cancer Center, UCSF

⁵Centro de Estudios de Estado y Sociedad, Buenos Aires, Argentina

⁶Universidad de Buenos Aires, Buenos Aires, Argentina

Abstract

This study evaluated the effect of factors reflecting appreciation of Indigenous culture and racial insults on alcohol and drug use initiation among multi-ethnic youth in Jujuy, Argentina. Students were surveyed from 27 secondary schools that were randomly selected to represent the province. A total of 3,040 eligible students in 10th grade, age 14 to 18 years were surveyed in 2006 and 2,660 of these same students completed surveys in 11th grade in 2007. Multivariate logistic regression models assessed the effect of appreciation for Indigenous cultures and reported exposure to racial insults in 10th grade on incident current alcohol drinking in previous 30 days, binge drinking (≥ 5 drinks at one sitting), and lifetime drug use (marijuana, inhalants or cocaine) in 11th grade among students not reporting these behaviors in 2006. In 2006, 63% of respondents reported high appreciation for Indigenous cultures and 39% had ever experienced racial insults. In 2007, incident current drinking was 24.4%, binge drinking 14.8%, and any drug use initiation was 4.1%. Exposure to racial insults increased the likelihood of binge drinking (OR=1.6; 95% CI 1.2–2.1) but was not significant for any drug use. Appreciation for Indigenous cultures reduced the risk of any drug use initiation (OR=0.5, 95% CI 0.3–0.7) but had no effect for alcohol drinking

Correspondence to: Eliseo J Pérez-Stable, MD, 6707 Democracy Boulevard, Suite 800, Bethesda, MD 20892-5465; Telephone: (415) 502-5601; Fax: (415) 502-8291; eliseo.perez-stable@nih.gov.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

outcomes. These effects were independent of Indigenous ethnicity. Enhancing appreciation for Indigenous cultures and decreasing racial insults are achievable goals that can be incorporated into programs to prevent youth substance use.

Keywords

culture; racism; substance use; youth; Indigenous; Latin America

Introduction

Alcohol and drug use and misuse significantly contribute to the global burden of disease (1, 2). Alcohol is linked to intentional and unintentional injuries and chronic diseases, and it accounts for about the same amount of the global burden of disability-adjusted life-years as tobacco (3). Drug use is linked to suicide, mental health disorders and cognitive impairments (2, 4). Substance use among adolescents merits special attention since this is a vulnerable period for addiction, and it may lead to social problems such as poor school performance and confronting personal and institutional violence (5–7).

Research to understand the psychosocial etiology of adolescent alcohol and drug use has a long history in North America but research in South America is rare. Within this region, Argentina has highest rates of alcohol drinking and cocaine use and shows a raising trend in substance use in general (8). Between 2001 and 2011 binge drinking increased from 29.7% to 63.4%, marijuana use from 4.6% to 13.9%, inhalants use from 0.9% to 4.5%, and cocaine use from 1.4% to 4.6%. The data indicates that current prevention efforts are insufficient, failing to address the complex phenomena that underlie youth's use of psychotropic substances. An under studied topic in a culturally diverse setting like South America (9), is the relationship between cultural factors and substance use (9–12). The province of Jujuy in Northwest Argentina is a unique environment for examining these relationships. The majority of the population is of Indigenous background of Andean and Amazonian ethnic groups, with a minority of European descent and an important segment of mixed-origin individuals (Indigenous and European). In this context a hierarchical social structure persists, with discrimination and racism against Indigenous Peoples (10); (11). At the same time, contemporary social change processes have fostered revalorization of Indigenous cultures. This is also a border area with intense trafficking of cocaine from contiguous producing countries.

We conducted a longitudinal analysis of how cultural factors influence substance use among youth in a social setting for which there has been little if any research apart from the authors' prior work. We examined the effect of two posited protective and risk factors -- appreciation for Indigenous cultures and exposure to racial insults--, on alcohol, marijuana, inhalants and cocaine use. This research is relevant in view of increasing consumption rates and the need to design effective prevention programs. It is also grounded in a new social setting for examining the web of substance use causation.

Longitudinal Studies of cultural factors and youth substance use

A number of studies in North America have assessed the relationship between youth substance use and risk and protective cultural factors, using different indicators across ethnic groups of varied socio-historical trajectories and yielding mixed results (12–19). Few studies have used prospective designs. A prospective study among Mexican heritage adolescents showed that linguistic acculturation was a risk factor for marijuana initiation for boys, was protective for inhalant use initiation, and had no effect on alcohol use (20). Another study showed that Latina/o youth experiencing discrimination reported more use on a combined measure of alcohol, cigarettes and marijuana (21). A study of American Indian adolescents found that perceived discrimination influenced early and problem alcohol drinking (22). In the same population, discrimination was not found to be a predictor of marijuana use (23). Among Navajo adolescents, discrimination was linked to boys' substance use, using a combined measure of alcohol, marijuana, hallucinogens, stimulants and inhalants. Navajo culture and in some cases, connection to White American culture, buffered the negative effect of discrimination (24). Among African American adolescents, perceived discrimination was associated with an increase in a combined measure of cigarettes, marijuana and alcohol use in adulthood (25). Similarly, early experiences of discrimination among African American adolescents were correlated with a combined measure of alcohol and marijuana use (26). The literature for Latin American countries is scarce and limited to adults. In Chile among adult primary health care patients, perceived discrimination was correlated with alcohol and illegal drug use (27). A longitudinal study among adults of the Amazonian Tsimane in Bolivia showed that adherence to shared cultural norms was negatively associated with alcohol drinking (28).

Cultural Appreciation, Racism and Substance Use

Cultural identity indicates the degree to which individuals perceive to be included in an ethnic group and it has been a central focus of health research (29–31). Affiliation to a cultural group however, does not imply that an individual holds a positive affect towards the group. At the cognitive and emotional levels, a range of positive or negative appraisals may emerge as individuals ascribe significance, meaning and value to a culture (32, 33) (34). A positive appraisal of one's culture is thought to contribute to well-being by conferring feelings of self-worth, connectedness, and purpose (35), and there is indication that in educational settings, promoting a positive appraisal for the culture of others improves the learning environment. We hypothesized that appreciation for historically undervalued Indigenous cultures will be protective across ethnic groups in a multicultural context. To the contrary, we hypothesized that exposure to racial insults will have a negative effect across ethnic groups. In previous analyses, exposure to racial insults referring to Indigenous stereotype characteristics was reported by all ethnic groups in this research site. (36) Racism refers to an organized system that assigns hierarchical status to specific groups and uses this ranking to preferentially allocate societal goods and resources to those who are regarded as inherently superior. Stress elicited by exposure to racial insults, even if the receptor does not identify with the posited group, may induce or contribute to alcohol or drug use as a coping strategy (37).

The ecological public health model (38) provides an overall framework for model building in this study. It is understood that substance use behavior is influenced by a wide array of individual, interpersonal and contextual variables. The social norms (39) and tension reduction theories (40, 41) are used to posit plausible explanations of the relations found among variables.

Methods

Sampling and Study Procedures

The study was conducted between 2006 and 2007 in the Province of Jujuy, in Northwest Argentina. Procedures were described in a previous publication (42). The 27 participating schools included 3690 8th grade students age 12 to 16 in 2004. Surveys were self-administered in class and students who were not located in the school at follow-up were surveyed at home. For this report we used data from the 3,040 students between ages 14 and 18 years and completed surveys in 2006. In 2007, 2,660 of 3,040 students (87.5% response rate) completed surveys at time 2 for the one-year cohort. The institutional Committee on Human Research and an NIH-certified human subjects research board in Buenos Aires based at *Centro de Educación Médica e Investigaciones Clínicas* (CEMIC) approved the research protocol. Passive consent was requested from caretakers and students signed an active consent. The questionnaire consisted of translated items from surveys of adolescents in the U.S. (43, 44), and questions developed through qualitative research in the target population (36, 45).

Exposure Variables: Appreciation for Indigenous Cultures and Racial Insults

The variable assessing appreciation for Indigenous cultures was developed through in depth qualitative interviews and was measured with the following items: 1) “It is important to teach children about traditions like *Pachamama* or *Pin Pin*”; 2) “It is important to continue with the traditions and customs of the elders”; and 3) “It is important to learn traditional languages like Quechua, Aymara, Guarani”. The items have a disagreement-agreement response set with a range of 1 to 5. For this analysis we coarsened to a binary indicator of high versus low appreciation (3.6 to 5 vs. 1 to 3.5, respectfully).

Questions on experiences of racial insults were also developed through qualitative research (36). We created two items to assess whether respondents had ever been called any of the following names in a way that was intended to insult or to bother them: 1) “*coya tonto*”, “*chaguanco*”, or “*indio cabeza dura*”; or 2) “*boliviano* or *bolita*”. The items were assessed with a Yes or No response and we constructed a binary variable indicating ever versus never being exposed to one of these two sets racial insults.

Demographics and Family Characteristics

Respondents reported their sex, age, religion, and school attendance. They selected their ethnic identity from the following list: Indigenous, mixed Indigenous and European, and European (42). A few respondents endorsing a category of “other ethnicity” including groups of more privileged social status locally (e.g., Arabs) were grouped with Europeans.

Respondents who self-identified as Indigenous, were also asked to select their identity from a list of Andean and Amazonian Indigenous groups.

The use of an Indigenous language in the family was ascertained as an indicator of family cultural context. Students reported on education and employment status of the primary provider. If the provider had not completed elementary school, was unemployed or on government welfare, youth were categorized as having low socioeconomic status. Religion was categorized as Catholic, Christian or Evangelical, and other low frequency religions.

Psychosocial Factors

Depressive symptoms were ascertained by asking whether the respondent in the past year felt sad and/or could not carry on his/her normal activities or obligations for at least two weeks (46). Thrill seeking orientation was measured with three items using a five-point disagreement-agreement response set: “I do not mind getting in trouble as long as I have fun”; “I like to do dangerous things”; “I like to do things that people say should not be done” (47). We defined scores of 3 to 5 as high and less than 3 as low. The Cronbach’s alpha for the items was 0.71.

Positive expectations for the future were measured with the following two questions: “I think that when I am older I will be able to have all that I need” and “I think that when I am older I will be able to work in what I like”. A list of role models was presented to respondents. Identification with “athletes” was used as an indicator of a conforming role model. “*Villeros*”, a stereotyped young shantytown dweller with a rebellious attitude towards established norms was used as an indicator of a defiant role model. To assess concern for body image we asked whether respondents were trying to lose/gain, maintain their weight, or were not concerned about their weight.

Outcome Variables

Participants reported on the consumption of any alcoholic beverages in the previous month (defined current drinking), and the consumption of five or more drinks on one occasion in the past month (episode of binge drinking). Respondents were asked if they had ever used marijuana, inhalants or solvents (e.g. glue, gasoline), cocaine, or other drugs (e.g. ecstasy, heroin). We also constructed a variable of lifetime use of any of these drugs.

Data Analysis

The sampling design was incorporated into all models by specifying geographic areas as strata and schools as clusters, as well as including weights to adjust for disproportionate stratification. A finite population correction was applied to adjust for the relatively large proportion of available schools sampled within each geographic area. Standard errors and confidence intervals were estimated via the Taylor expansion approximation (Stata version 11.2). We conducted descriptive analyses examining the distribution of demographic and family characteristics and of cultural and psychosocial factors by sex. Chi-square tests and *p* values were calculated. We estimated the total and sex-specific prevalence rates of alcohol drinking and drug use variables in 2006 or at Time-1 when students were in 10th grade, and the respective incidence rates in 2007 or at Time-2 when students were in 11th grade.

Multivariate logistic regression models included cultural appreciation and racial insults as simultaneous independent variables. The models assessed the effect of the independent variables at Time-1 net of one another and of the control variables, on alcohol and drug use at Time-2. Models with the “other drug” outcome (e.g. ecstasy, heroin) yielded unstable results because they were rarely endorsed. These drugs were only included in the any drug use outcome models. For each outcome the models were fit to data from the subsample of respondents who were negative for the outcome at Time-1 to capture incident alcohol and drug use at Time-2. Covariates were selected on the basis of those related to substance use in the literature or in our previously published studies (6, 36, 48). We estimated adjusted odds ratios and 95% confidence intervals. For each outcome we tested interactions of cultural appreciation by sex and by ethnicity, of exposure to racial insults by sex and by ethnicity, and of cultural appreciation levels by exposure to racial insults.

Results

Demographic Characteristics

Among the 3,040 participants who were in the selected age range of 14 to 18 years in 2006, 63.4% reported high levels of cultural appreciation (Table 1) and the proportion did not differ by ethnicity. A smaller proportion of girls than boys reported ever exposure to racial insults (34.5% vs. 44.1%), but the proportions did not differ by ethnicity.

Nearly 20% of youth were not attending school. A larger proportion of girls were classified as low SES (25.8% vs. 20.6%). Girls reported lower levels of parental support (62.4% vs. 72.8%) and thrill seeking orientation (10.4% vs. 16.9%), and were less likely to identify with a conforming (athletic) role model (59.3% vs. 79.8%). On the other hand a greater proportion of girls reported having depressive symptoms (44.7% vs. 29.2%) and trying to lose or gain weight (39.2% vs. 24.5%).

Prevalence and Incidence Rates

Current alcohol drinking prevalence rates were higher among boys (40% vs. 33%) and their prevalence of binge drinking was more than double that of girls (27% vs. 13%; Table 2a). At time 2, boys and girls without current drinking in 2006 had similar rates of current drinking in the past 30 days (27% vs. 23%) but the rate of incident binge drinking was higher among boys (19% vs. 12%; Table 2b).

Lifetime drug use prevalence rates were higher among boys for marijuana (5% vs. 2%) and any drug use (7% vs. 4%), but similar across both sexes for inhalants (2.7%), cocaine (0.9%) and other drug use (0.7%) at time 1 (Table 2a). Incidence rates were similar across boys and girls for the use of marijuana (3.3%), inhalants (1.6%) and other drugs (0.5%). Rates were higher among girls for cocaine (1.3% vs. 0.4%) and higher among boys for any drug use (5.1% vs. 3.3%; Table 2b). To evaluate reliability of responses, we found that 2.6% of respondent answered “yes” to using any drugs in 2006 but responded “no” in 2007.

Predictors of Alcohol Drinking

In multivariate logistic regression models exposure to racial insults increased rate of binge drinking in the previous 30 days (OR=1.6; 95% CI = 1.2–2.1; Table 3). Appreciation for Indigenous cultures did not have a significant effect on the alcohol drinking outcomes. Among the covariates, being a boy (OR=1.7; 95% CI = 1.3–2.3), reporting high levels of thrill seeking orientation (OR=1.9; 95% CI = 1.3–2.8) and identifying with a defiant role model (OR=1.6; 95% CI = 1.2–1.9) increased the likelihood of binge drinking in past 30 days. Reporting high levels of thrill seeking orientation (OR=2.1; 95% CI=1.4–3.1) and identifying with a defiant role model (OR=1.4; 95% CI = 1.1–1.9) increased the rate of current alcohol drinking among those not reporting this behavior the previous year. European youth had increased likelihood of current drinking in previous 30 days at time 2 compared with Indigenous Andean youth (OR=1.6; 95% CI = 1.2–2.1) (data not shown).

Predictors of Drug Use

High levels of appreciation for Indigenous cultures reduced by half or less the odds of drug use initiation in multivariate models. However, exposure to racial insults was not a significant risk factor for drug use. Boys were more likely to initiate marijuana (OR=2.1; 95% CI = 1.2–3.8) or any drug use (OR=1.7; 95% CI = 1.1–2.7), but the likelihood of inhalant use initiation was similar for boys and girls, and the risk of cocaine initiation was lower for boys (OR=0.2; 95% CI = 0.1–0.7). Youth who self-identified as being Indigenous had lower odds of inhalant use initiation compared with Europeans (OR=0.3; 95% CI = 0.1–0.7). When examining disaggregated Indigenous categories European (OR=3.7; 95% CI = 1.5–9.3) and Amazonian youth (OR=4.3; 95% CI = 1.1–17.0) had higher risk of inhalant use compared with Andeans (data not shown).

The perception of having high parental support was protective for incident marijuana use (OR=0.3; 95% CI = 0.1–0.5) and any drug use initiation (OR=0.4; 95% CI = 0.2–0.7). Thrill seeking orientation increased the risk of incident inhalant use (OR=5.0; 95% CI = 2.4–10.4) and cocaine initiation (OR=5.7; 95% CI = 2.1–15.0). Reduced positive expectations for the future was a risk factor only for cocaine use initiation (OR=3.6; 95% CI = 1.3–9.9). Identification with defiant role models increased the risk of incident inhalant use (OR=4.9; 95% CI = 2.4–10.1) and any drug use initiation (OR=1.9; 95% CI = 1.1–3.5). Identification with a conforming role model was protective only for cocaine (OR=0.3; 95% CI = 0.2–0.7).

Interaction effects

The protective effect of appreciation for Indigenous cultures on marijuana (OR=0.2; 95% CI = 0.1–0.3) was significant only among girls. The effects of the exposure variables did not vary across groups by ethnic identity and the effect of appreciation for Indigenous cultures was similar for youth with and without exposure to racial insults.

Discussion

This is the first study in Latin America to assess the effect of cultural factors on youth's substance use behavior. Appreciation for Indigenous cultures resulted in beneficial effects and reporting racist insults had detrimental effects, across the different ethnic groups in the

study. Significant effects were found after controlling for a wide set of co-variables. Among longitudinal studies reporting effects on single substance use, our results are in agreement with those showing that discrimination influences alcohol use among North American Indian youth (22) but does not predict marijuana use (23, 24), with studies showing that discrimination influences alcohol drinking among African American youth (26, 49), and with a study showing no relationship between a cultural identity indicator and alcohol drinking among Mexican American youth (20).

Different effect patterns of cultural appreciation on alcohol and drug use initiation may be explained by the social norms associated with each substance type (50–54). Argentina is a permissive society and underage drinking commonly occurs under adult supervision (55). However, Indigenous communities in the region maintain comparatively strong social sanctions against drug use (56).

The tension reduction approach provides a basis for explaining the effect of racism on alcohol drinking initiation (40, 41), a salient finding, given the high prevalence of alcohol drinking in this sample. Alcohol drinking may occur in response to negative emotional states (17)(15, 37). In Argentina alcoholic beverages are easily available to youth. The lack of effect of exposure to racial insults on drug use merits further investigation. One possible explanation may be related to greater difficulties in accessing illegal substances. It is also possible that aspects of racism not assessed in this study may play a role in drug use initiation.

The protective effect of cultural appreciation on marijuana use was significant only among girls. Several studies have shown that the weakening of cultural connectedness has a greater impact on adolescent girls than boys (57, 58). More lax social trends regarding marijuana use in Argentina (59) may be influencing boys to a greater extent than girls.

The effects of covariates varied across substances, highlighting the importance of evaluating substances separately. Indicators of externalizing behavior (60) like thrill-seeking orientation and identification with a defiant role model were in general associated with increased odds of alcohol and drug use initiation, but not for marijuana. This may be related to the increasing social acceptance of this substance). For cocaine initiation, low expectations for the future were a risk factor and identification with a conforming role model was protective. Parental support was protective for marijuana and for any drug use initiation.

The two main independent variables of appreciation of Indigenous culture and exposure to racial insults may be susceptible to potential confounding with the youths' ethnic identification. The question raised is whether European youth's responses to these questions have the same meaning as those of Indigenous or mixed identity and thus affect interpretation of these results. For example, asking youth whether it is important to learn traditional Indigenous languages would carry a different meaning depending on whether those are the languages of their own cultures (as is the case for the Indigenous and mixed youth) or the languages of an ethnic culture that is different from one's own (as is the case for the European youth). Youth of European origin who express appreciation for Indigenous

cultures may have a greater degree of social integration and higher esteem for the social context in which they live, factors that contribute to substance use risk reduction.

We found similar results in re-analysis of the data excluding the European youth. However, it is possible that in a geographical context where Indigenous and Mixed youth do not make up the vast majority of the sample (as would be the case in other provinces of Argentina), analyses by racial group may differ from these findings. Similarly, the detrimental effect of racism was not restricted to Indigenous youth. Racial insults affect the human dignity of Indigenous individuals and may also affect the non-Indigenous by eliciting the perception of possessing socially devalued characteristics. To our knowledge this is the first study to assess cultural variables across all ethnic groups, indicating that having a positive evaluation of a subaltern social sector may contribute to collective wellbeing in a multicultural context (61). Future studies are granted to further elucidate this issue.

Our findings are subject to several limitations. Self-reports from youth who might under-or over-report their substance use, may bias the results. Furthermore reliability may vary due to inattentiveness, misrepresentation or other reasons. Regardless, this is the standard methodology used in population-based surveys (62), and the lack of perfect reliability of outcome variables does not bias regression parameter estimates as reported. Although the data are not current, as the sociocultural environment remains stable and increasing rates persist, results remain relevant. Cultural factors were measured with indicators encompassing few dimensions of complex psychosocial phenomena. Therefore, lack of significance with the outcomes of interest should be interpreted with caution. On the other hand, the longitudinal study design is a strength over cross-sectional studies. We interviewed school dropouts, broadening the representativeness of results, and we used separate measures of substance use.

These results may have implications for framing public health policies that incorporate a vision of equity and human rights. Enhancing appreciation for Indigenous cultures and decreasing racist acts are potentially modifiable factors in multi-ethnic communities that can be incorporated into programs to prevent youth substance use in Latin America.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This research was funded by Grant no. TW05935 from the Tobacco Research Network Program, Fogarty International Center, National Cancer Institute, National Institute of Drug Abuse, National Institutes of Health and by Grant no. 001726-037 from Research on International Tobacco Control, International Development Research Center, Canada. The administrative and research support of Cecilia Populus-Eudave and the research staff in Jujuy made this study possible.

References

1. Ezzati, M., Lopez, AD., Rodgers, A., Murray, CJL. Organization WH. Global and Regional Burden of Disease Attributable to Selected Major Risk Factors. Geneva: World Health Organization; 2004.

2. Rehm, J., Room, R. The global burden of disease attributable to alcohol, tobacco and illicit drugs. Stockwell, T.Gruenewald, PJ.Toumbourou, JW., Loxley, W., editors. London: John Wiley & Sons; 2005.
3. Murray, CJL., Lopez, AD. The Global Burden of Disease. In: Lopez, Ma, editor. A comprehensive assessment of mortality and disability from disease, injuries and risk factors in 1990 and projected to 2020. Boston: Harvard School of Public Health; 1996. p. 43
4. Cairney S, Maruff P, Burns C, Currie B. The neurobehavioural consequences of petrol (gasoline) sniffing. *Neuroscience and biobehavioral reviews*. 2002 Jan; 26(1):81–9. Epub 2002/02/12. eng. [PubMed: 11835986]
5. Crews F, He J, Hodge C. Adolescent cortical development: a critical period of vulnerability for addiction. *Pharmacology, biochemistry, and behavior*. 2007 Feb; 86(2):189–99. Epub 2007/01/16. eng.
6. Alderete E, Kaplan CP, Nah G, Perez-Stable EJ. Problems related to alcohol drinking among youth in Jujuy, Argentina. *Salud publica de Mexico*. 2008 Jul-Aug;50(4):300–7. Epub 2008/08/02. Problemas relacionados con el consumo de alcohol en jovenes de la provincia de Jujuy, Argentina. spa. [PubMed: 18670721]
7. Pierobon M, Barak M, Hazrati S, Jacobsen KH. Alcohol consumption and violence among Argentine adolescents. *Jornal de pediatria*. 2013 Jan-Feb;89(1):100–7. Epub 2013/04/03. eng. [PubMed: 23544817]
8. CICAD. Situación Actual Del Uso De Drogas En Las Américas Y Desafíos Futuros. Washington: Comision Interamericana Para El Control Del Abuso De Drogas; 2011.
9. Gonzalez Burchard E, Borrell LN, Choudhry S, Naqvi M, Tsai HJ, Rodriguez-Santana JR, et al. Latino populations: a unique opportunity for the study of race, genetics, and social environment in epidemiological research. *American journal of public health*. 2005 Dec; 95(12):2161–8. Epub 2005/11/01. eng. [PubMed: 16257940]
10. Organization of American States. Report on Drug Use in the Americas. Washington, D.C: Inter-American Drug Abuse Control Commission, Inter-American Observatory on Drugs; 2015.
11. Torres-Parodi C, Bolis M. The evolution of the ethnicity/race concept and its impact on framing policies that promote equity. *Revista panamericana de salud publica = Pan American journal of public health*. 2007 Dec; 22(6):405–16. Epub 2008/02/23. Evolucion del concepto etnia/raza y su impacto en la formulacion de politicas para la equidad. spa. [PubMed: 18291060]
12. Kam JA, Cleveland MJ. Perceived discrimination as a risk factor for Latina/o youth's substance use: do parent- and peer-based communication and relationship resources act as protective factors? *Health communication*. 2011 Mar; 26(2):111–24. Epub 2011/01/29. eng. [PubMed: 21271420]
13. Whitbeck LB, Hoyt DR, McMorris BJ, Chen X, Stubben JD. Perceived discrimination and early substance abuse among American Indian children. *Journal of health and social behavior*. 2001 Dec; 42(4):405–24. Epub 2002/02/08. eng. [PubMed: 11831140]
14. Whitbeck LB, Chen X, Hoyt DR, Adams GW. Discrimination, historical loss and enculturation: culturally specific risk and resiliency factors for alcohol abuse among American Indians. *Journal of studies on alcohol*. 2004 Jul; 65(4):409–18. Epub 2004/09/21. eng. [PubMed: 15376814]
15. Terrell F, Miller AR, Foster K, Watkins CE Jr. Racial discrimination-induced anger and alcohol use among black adolescents. *Adolescence*. 2006 Fall;41(163):485–92. Epub 2007/01/18. eng. [PubMed: 17225663]
16. Umana-Taylor AJ, Alfaro EC, Bamaca MY, Guimond AB. The central role of family socialization in Latino adolescents' cultural orientation. *Journal of Marriage and Family*. 2009; 71:46–60.
17. Martin JK, Tuch SA, Roman PM. Problem drinking patterns among African Americans: the impacts of reports of discrimination, perceptions of prejudice, and "risky" coping strategies. *Journal of health and social behavior*. 2003 Sep; 44(3):408–25. Epub 2003/10/30. eng. [PubMed: 14582316]
18. Nasim A, Belgrave FZ, Jagers RJ, Wilson KD, Owens K. The moderating effects of culture on peer deviance and alcohol use among high-risk African-American Adolescents. *Journal of drug education*. 2007; 37(3):335–63. Epub 2007/12/01. eng. [PubMed: 18047186]

19. German M, Gonzales NA, Dumka L. Familism Values as a Protective Factor for Mexican-origin Adolescents Exposed to Deviant Peers. *The Journal of early adolescence*. 2009 Feb; 29(1):16–42. Epub 2009/02/01. Eng. [PubMed: 21776180]
20. Marsiglia FF, Yabiku ST, Kulis S, Nieri T, Parsai M, Becerra D. The Influence of Linguistic Acculturation and Gender on the Initiation of Substance Use Among Mexican Heritage Preadolescents in the Borderlands. *The Journal of early adolescence*. 2011 Apr; 31(2):271–99. Epub 2011/06/11. Eng. [PubMed: 21660121]
21. Kam JA, Cleveland MJ, Hecht ML. Applying general strain theory to examine perceived discrimination's indirect relation to Mexican-heritage youth's alcohol, cigarette, and marijuana use. *Prevention science : the official journal of the Society for Prevention Research*. 2010 Dec; 11(4):397–410. Epub 2010/05/22. eng. [PubMed: 20490921]
22. Cheadle JE, Whitbeck LB. Alcohol use trajectories and problem drinking over the course of adolescence: a study of north american indigenous youth and their caretakers. *Journal of health and social behavior*. 2011 Jun; 52(2):228–45. Epub 2011/05/12. eng. [PubMed: 21558489]
23. Cheadle JE, Sittner Hartshorn KJ. Marijuana use development over the course of adolescence among North American Indigenous youth. *Social science research*. 2012 Sep; 41(5):1227–40. Epub 2012/09/29. eng. [PubMed: 23017929]
24. Galliher RV, Jones MD, Dahl A. Concurrent and longitudinal effects of ethnic identity and experiences of discrimination on psychosocial adjustment of Navajo adolescents. *Developmental psychology*. 2011 Mar; 47(2):509–26. Epub 2010/12/15. eng. [PubMed: 21142373]
25. Brodish AB, Cogburn CD, Fuller-Rowell TE, Peck S, Malanchuk O, Eccles JS. Perceived Racial Discrimination as a Predictor of Health Behaviors: the Moderating Role of Gender. *Race and social problems*. 2011 Oct 1; 3(3):160–9. Epub 2012/07/31. Eng. [PubMed: 22844386]
26. Gibbons FX, Gerrard M, Cleveland MJ, Wills TA, Brody G. Perceived discrimination and substance use in African American parents and their children: a panel study. *Journal of personality and social psychology*. 2004 Apr; 86(4):517–29. Epub 2004/04/01. eng. [PubMed: 15053703]
27. Capezza NM, Zlotnick C, Kohn R, Vicente B, Saldivia S. Perceived discrimination is a potential contributing factor to substance use and mental health problems among primary care patients in Chile. *Journal of addiction medicine*. 2012 Dec; 6(4):297–303. Epub 2012/09/19. eng. [PubMed: 22987021]
28. Reyes-Garcia V, Gravlee CC, McDade TW, Huanca T, Leonard WR, Tanner S. Cultural consonance and psychological well-being. Estimates using longitudinal data from an Amazonian society. *Culture, medicine and psychiatry*. 2010 Mar; 34(1):186–203. Epub 2009/12/04. eng.
29. Oetting ER, Beauvais F. Orthogonal cultural identification theory: the cultural identification of minority adolescents. *The International journal of the addictions*. 1991; 25(5A-6A):655–85. Epub 1990/01/01. eng.
30. Cokley K. Critical issues in the measurement of ethnic and racial identity: A referendum on the state of the field. *Journal of Counseling Psychology*. 2007; 54:224–34.
31. Phinney JS. Stages of ethnic identity development in minority group adolescents. *Journal of Early Adolescence*. 1989; 9:34–49.
32. Sellers RM, Smith MA, Shelton JN, Rowley SA, Chavous TM. Multidimensional model of racial identity: a reconceptualization of African American racial identity. *Personality and social psychology review : an official journal of the Society for Personality and Social Psychology, Inc*. 1998; 2(1):18–39. Epub 2005/01/14. eng.
33. Epplé S, Thubauville S. Cultural diversity in Ethiopia between Appreciation and Suppression. *Paideuma*. 2012; 58:153–65.
34. Morgan DL. Appropriation, Appreciation, Accommodation: Indigenous Wisdoms and Knowledges in Higher Education. *International Review of Education*. 2003; 49(1–2):35–49.
35. Kiang L, Yip T, Gonzales-Backen M, Witkow M, Fuligni AJ. Ethnic identity and the daily psychological well-being of adolescents from Mexican and Chinese backgrounds. *Child development*. 2006 Sep-Oct; 77(5):1338–50. Epub 2006/09/27. eng. [PubMed: 16999802]
36. Alderete E, Monteban M, Gregorich S, Kaplan CP, Mejia R, Perez-Stable EJ. Smoking and exposure to racial insults among multiethnic youth in Jujuy, Argentina. *Cancer causes & control : CCC*. 2012 Mar; 23(Suppl 1):37–44. Epub 2012/02/22. eng. [PubMed: 22350863]

37. Gerrard M, Stock ML, Roberts ME, Gibbons FX, O'Hara RE, Weng CY, et al. Coping with racial discrimination: the role of substance use. *Psychology of addictive behaviors : journal of the Society of Psychologists in Addictive Behaviors*. 2012 Sep; 26(3):550–60. Epub 2012/05/02. eng. [PubMed: 22545585]
38. Green LW, Richard L, Potvin L. Ecological foundations of health promotion. *American journal of health promotion : AJHP*. 1996 Mar-Apr;10(4):270–81. Epub 1996/02/07. eng. [PubMed: 10159708]
39. Berkowitz, AD. Applications of Social Norms Theory to Other Health and Social Justice Issues. In: Perkins, HW., editor. *The Social Norms Approach to Preventing School and College Age Substance Abuse: A Handbook for Educators, Counselors, Clinicians*. Vol. Chapter 16. San Francisco: Jossey-Bass; 2003.
40. Goldman, MS., Del Boca, FK., Darkes, J. Alcohol expectancy theory: The application of cognitive neuroscience. In: Blane, KELHT., editor. *Psychological theories of drinking and alcoholism*. 2. New York: Guilford Press; 1999.
41. Greely, J., Oei, TP. Alcohol and Tension Reduction. In: Leonard, KE., Blane, HT., editors. *Psychological Theories of Drinking and Alcoholism*. 2. New York: Guilford Press; 1999. p. 14–53.
42. Alderete E, Kaplan CP, Gregorich SE, Mejia R, Perez-Stable EJ. Smoking behavior and ethnicity in Jujuy, Argentina: evidence from a low-income youth sample. *Substance use & misuse*. 2009; 44(5):632–46. Epub 2009/04/11. eng. [PubMed: 19360537]
43. Fryar CD, Merino MC, Hirsch R, Porter KS. Smoking, alcohol use, and illicit drug use reported by adolescents aged 12–17 years: United States, 1999–2004. *National health statistics reports*. 2009 May; 20(15):1–23. Epub 2009/07/29. eng.
44. MMWR. Tobacco, alcohol, and other drug use among high school students--United States, 1991. *Morbidity and mortality weekly report*. 1992 Sep 18; 41(37):698–703. Epub 1992/09/18. eng. [PubMed: 1518502]
45. Alderete E, Kaplan CP, Gregorich SE, Perez-Stable EJ. Use of alternative tobacco products in multiethnic youth from Jujuy, Argentina. *Journal of environmental and public health*. 2010; 2010:795265. Epub 2010/03/20. eng. [PubMed: 20300454]
46. Benjet C, Borges G, Medina-Mora ME, Fleiz C, Blanco J, Zambrano J, et al. Prevalence and socio-demographic correlates of drug use among adolescents: results from the Mexican Adolescent Mental Health Survey. *Addiction*. 2007 Aug; 102(8):1261–8. Epub 2007/07/13. eng. [PubMed: 17624976]
47. Vega WA, Zimmerman RS, Warheit GJ, Apospori E, Gil AG. Risk factors for early adolescent drug use in four ethnic and racial groups. *American journal of public health*. 1993 Feb; 83(2):185–9. Epub 1993/02/01. eng. [PubMed: 8427320]
48. Mejia R, Kaplan CP, Alderete E, Gregorich SE, Perez-Stable EJ. Influence of gender role attitudes on smoking and drinking among girls from Jujuy, Argentina. *Preventive medicine*. 2013 Sep; 57(3):194–7. Epub 2013/06/05. eng. [PubMed: 23732243]
49. Gibbons FX, Yeh HC, Gerrard M, Cleveland MJ, Cutrona C, Simons RL, et al. Early experience with racial discrimination and conduct disorder as predictors of subsequent drug use: a critical period hypothesis. *Drug and alcohol dependence*. 2007 Apr; 88(Suppl 1):S27–37. Epub 2007/02/06. eng. [PubMed: 17275213]
50. Shmulewitz D, Wall MM, Keyes KM, Aharonovich E, Aivadyan C, Greenstein E, et al. Alcohol use disorders and perceived drinking norms: ethnic differences in Israeli adults. *Journal of studies on alcohol and drugs*. 2012 Nov; 73(6):981–90. Epub 2012/10/06. eng. [PubMed: 23036217]
51. Lewis MA, Neighbors C, Geisner IM, Lee CM, Kilmer JR, Atkins DC. Examining the associations among severity of injunctive drinking norms, alcohol consumption, and alcohol-related negative consequences: the moderating roles of alcohol consumption and identity. *Psychology of addictive behaviors : journal of the Society of Psychologists in Addictive Behaviors*. 2010 Jun; 24(2):177–89. Epub 2010/06/23. eng. [PubMed: 20565144]
52. Holmila M, Raitasalo K, Knibbe R, Selin K. Country variations in family members' informal pressure to drink less. *Contemporary drug problems*. 2009 Apr 1;36(1/2) nihpa126808. Epub 2010/01/20. Eng.

53. Room R. Intoxication and bad behaviour: understanding cultural differences in the link. *Social science & medicine*. 2001 Jul; 53(2):189–98. Epub 2001/06/21. eng. [PubMed: 11414386]
54. Caetano R, Clark CL. Trends in situational norms and attitudes toward drinking among whites, blacks, and hispanics: 1984–1995. *Drug and alcohol dependence*. 1999 Mar 1; 54(1):45–56. Epub 1999/04/02. eng. [PubMed: 10101616]
55. Pilatti A, Godoy JC, Brussino S, Pautassi RM. Underage drinking: prevalence and risk factors associated with drinking experiences among Argentinean children. *Alcohol*. 2013 Jun; 47(4):323–31. Epub 2013/04/18. eng. [PubMed: 23591270]
56. Knight GP, Berkel C, Umana-Taylor AJ, Gonzales NA, Ettekal I, Jaconis M, et al. The Familial Socialization of Culturally Related Values in Mexican American Families. *Journal of marriage and the family*. 2011 Oct; 73(5):913–25. Epub 2011/10/25. Eng. [PubMed: 22021936]
57. Marsiglia FF, Kulis S, Hussaini SK, Nieri TA, Becerra D. Gender differences in the effect of linguistic acculturation on substance use among Mexican-origin youth in the southwest United States. *Journal of ethnicity in substance abuse*. 2010; 9(1):40–63. Epub 2010/04/15. eng. [PubMed: 20390972]
58. Wahl AM, Eitle TM. Gender, acculturation and alcohol use among Latina/o adolescents: a multi-ethnic comparison. *Journal of immigrant and minority health/Center for Minority Public Health*. 2010 Apr; 12(2):153–65. Epub 2008/09/23. eng.
59. Inter-American Drug Abuse Control Commission. Segundo Estudio Conjunto. Washington: Naciones Unidas. Oficina Contra la Droga y el Delito; 2010. Informe subregional sobre uso de drogas en poblacion escolarizada.
60. Gibbons FX, O'Hara RE, Stock ML, Gerrard M, Weng CY, Wills TA. The erosive effects of racism: reduced self-control mediates the relation between perceived racial discrimination and substance use in African American adolescents. *Journal of personality and social psychology*. 2012 May; 102(5):1089–104. Epub 2012/03/07. eng. [PubMed: 22390225]
61. Alderete, E. The Health of Indigenous Peoples. 1999. p. 98WHO/SDE/HSD/99.1
62. Brener ND, Kann L, Shanklin S, Kinchen S, Eaton DK, Hawkins J, et al. Methodology of the Youth Risk Behavior Surveillance System--2013. *MMWR Recommendations and reports : Morbidity and mortality weekly report Recommendations and reports/Centers for Disease Control*. 2013 Mar 1; 62(RR-1):1–20. Epub 2013/03/01. eng.

Highlights

- 63% of 3040 students had high appreciation for Indigenous cultures; 39% reported racial insults
- At one year, 24.4% now reported drinking, 14.8% binge drinking, and 4.1% any drug use.
- Exposure to racial insults increased odds of binge drinking but was not significant for any drug use
- Appreciation of Indigenous cultures reduced risk of drug use initiation but not alcohol
- These topics can be incorporated into programs to prevent youth substance use.

Table 1

Demographic, cultural and psychosocial variables by sex among youth age 14 to 18 years, from Jujuy, Argentina, 2006

TOTAL N=3,040	GIRLS N=1,645	BOYS N=1,395	Chi square p value
	%	%	
CULTURAL VARIABLES			
<i>Ethnic self-identification</i>			
<i>European</i>	6.8	11.8	0.003
<i>Indigenous</i>	71.1	65.8	
<i>Mixed Indigenous-European</i>	22.0	22.3	
<i>Indigenous Language spoken in the family</i>			
<i>No</i>	68.7	69.2	0.876
<i>Yes</i>	31.3	30.8	
<i>Appreciation for Indigenous Cultures Score</i>			
<i>Low</i>	35.1	39.1	0.077
<i>High</i>	64.9	60.9	
<i>Exposure to Racial Insults</i>			
<i>Never</i>	65.5	55.9	<0.001
<i>Ever</i>	34.5	44.1	
DEMOGRAPHIC VARIABLES			
<i>Age in years</i>			
<i>14–15</i>	67.2	66.4	0.699
<i>16–18</i>	32.8	33.6	
<i>Currently attending school</i>			
<i>No</i>	17.7	20.9	0.128
<i>Yes</i>	82.3	79.1	
<i>Religion</i>			
<i>Catholic</i>	85.3	85.5	0.652
<i>Evangelical/Christian</i>	10.6	9.9	
<i>Other</i>	4.1	4.6	
<i>Low SES</i>			
<i>No</i>	74.2	79.4	0.014
<i>Yes</i>	25.8	20.6	
PSYCHOSOCIAL VARIABLES			
<i>Symptoms of depression</i>			
<i>No</i>	55.3	70.8	<0.001
<i>Yes</i>	44.7	29.2	
<i>Positive expectations for the future</i>			
<i>Never to almost always</i>	30.3	29.5	0.692

TOTAL N=3,040	GIRLS N=1,645	BOYS N=1,395	Chi square p value
	%	%	
<i>Always</i>	69.7	70.5	
<i>Parental support</i>			
<i>Never/almost never</i>	13.7	6.6	<0.001
<i>Sometimes</i>	23.9	20.6	
<i>Almost always/always</i>	62.4	72.8	
<i>Thrill seeking orientation</i>			
<i>Low</i>	89.6	83.1	0.004
<i>High</i>	10.4	16.9	
<i>Body image</i>			
<i>No action</i>	31.4	44.5	<0.001
<i>Trying to maintain weight</i>	29.4	31.0	
<i>Trying to lose/gain weight</i>	39.2	24.5	
<i>Identification with defiant role model</i>			
<i>None</i>	58.9	58.0	0.522
<i>Almost none to a lot</i>	41.1	42.0	
<i>Identification with conforming role model</i>			
<i>Low</i>	40.7	20.2	<0.001
<i>High</i>	59.3	79.8	

Table 2a
Prevalence rates of alcohol and drug use among 3,040 youth in Jujuy, Argentina, 2006

		Prevalence Rates (T1)							
		Alcohol Use		Lifetime Drug Use					
	Current N=2,981 (1,606 Girls/ 1,375 Boys)	Current Binge N=3,005 (1,634 Girls/ 1,371 Boys)	Marijuana N=3,032 (1,643 Girls/ 1,389 Boys)	Inhalants N=3,029 (1,640 Girls/ 1,389 Boys)	Cocaine N=3,025 (1,638 Girls/ 1,387 Boys)	Other Drugs N=3,031 (1,641 Girls/ 1,390 Boys)	Any Drugs N=3,016 (1,634 Girls/ 1,382 Boys)		
	N	N	N	N	N	N	N	N	N
	%*	%*	%*	%*	%*	%*	%*	%*	%*
	(SE)	(SE)	(SE)	(SE)	(SE)	(SE)	(SE)	(SE)	(SE)
Girls	506 32.9% (1.5)	203 13.2% (1.5)	31 2.2% (0.4)	40 2.6% (0.4)	12 0.7% (0.2)	9 0.5% (0.3)	66 4.4% (0.5)	66	66
Boys	536 40.1% (2.1)	344 26.7% (1.8)	64 5.0% (0.8)	37 2.8% (0.3)	18 1.3% (0.3)	13 1.0% (0.3)	88 6.9% (0.8)	88	88
Total	1,042 36.2% (1.2)	547 19.3% (0.6)	95 3.5% (0.5)	77 2.7% (0.3)	30 0.9% (0.2)	22 0.7% (0.2)	154 5.5% (0.6)	154	154
p-value	0.011	<0.001	<0.001	0.631	0.077	0.281	<0.001		

Current Drinking is defined by consuming one alcohol drink in the previous 30 days

Binge drinking is defined by consuming 5 drinks at one sitting

* Percentages are weighted so the Ns will not add up to the weighted percentages of the total..

Rates of past 30-day alcohol use and incident drug use among 2,660 youth in Jujuy, Argentina, 2007

Table 2b

		Incidence Rates (T2)							
Incident Alcohol Use				Drug Use Initiation					
	Current N=1,620 (929 Girls/691 Boys)	Current Binge N=2,074 (1,225 Girls/849 Boys)	Marijuana N=2,464 (1,381 Girls/ 1,083 Boys)	Inhalants N=2,464 (1,368 Girls/ 1,096 Boys)	Cocaine N=2,506 (1,391 Girls/ 1,115 Boys)	Other Drugs N=2,511 (1,394 Girls/ 1,117 Boys)	Any Drugs N=2,391 (1,338 Girls/ 1,053 Boys)		
	N	N	N	N	N	N	N	N	N
	%*	%*	%*	%*	%*	%*	%*	%*	%*
	(SE)	(SE)	(SE)	(SE)	(SE)	(SE)	(SE)	(SE)	(SE)
Girls	203	137	29	18	15	9	38	38	38
	22.6%	12.0%	2.6%	1.4%	1.3%	0.8%	3.3%	3.3%	3.3%
	(1.5)	(0.6)	(0.5)	(0.4)	(0.4)	(0.4)	(0.5)	(0.5)	(0.5)
Boys	188	168	39	17	4	3	46	46	46
	26.9%	18.9%	4.3%	1.7%	0.4%	0.2%	5.1%	5.1%	5.1%
	(3.1)	(2.0)	(0.7)	(0.4)	(0.2)	(0.1)	(0.7)	(0.7)	(0.7)
Total	391	305	68	35	19	12	84	84	84
	24.4%	14.8%	3.3%	1.6%	0.9% (0.2)	0.5%	4.1%	4.1%	4.1%
	(1.9)	(1.1)	(0.5)	(0.3)	(0.2)	(0.2)	(0.5)	(0.5)	(0.5)
p-value	0.155	<0.001	0.060	0.609	0.034	0.167	0.024	0.024	0.024

Current Drinking is defined by consuming one alcohol drink in the previous 30 days

Current Binge drinking is defined by consuming 5 drinks at one sitting in the previous 30 days

Sample N for each behavior varies to reflect the number who did not report the behavior in 2006

* Percentages are weighted so the Ns will not add up to the weighted percentages of the total.

Cultural, demographic and psychosocial predictors of current drinking behavior and incident substance use among youth in Jujuy, Argentina, 2006–2007

Table 3

	Alcohol Use				Incident Drug Use in One Year							
	Current Drinking (past 30 days) N=1,428		Current Binge Drinking N=1,825		Marijuana N=2,176		Inhalants N=2,177		Cocaine N=2,214		Any Drug N=2,114	
	n	OR (95 % CI)	n	OR (95 % CI)	n	OR (95 % CI)	n	OR (95% CI)	n	OR (95% CI)	n	OR (95 % CI)
CUTLURAL VARIABLES												
Ethnic self-identification												
European (ref)	116	---	147	---	182	---	185	---	188	---	176	---
Indigenous	1,003	0.7 (0.5–1.0)	1,288	0.8 (0.5–1.2)	1,524	0.7 (0.3–1.8)	1,520	0.3 (0.1–0.7)*	1,542	0.7 (0.2–2.3)	1,481	0.6 (0.3–1.1)
Mixed	309	0.9 (0.6–1.2)	390	0.8 (0.6–1.1)	470	1.1 (0.5–2.5)	472	0.4 (0.1–1.0)	484	0.9 (0.2–3.3)	457	0.9 (0.5–1.5)
Indigenous Language spoken in the family												
No (ref)	951	---	1,210	---	1,443	---	1,441	---	1,472	---	1,400	---
Yes	477	1.1 (0.9–1.5)	615	1.3 (0.9–1.7)	733	0.9 (0.4–1.8)	736	1.5 (0.6–3.6)	742	1.3 (0.5–3.4)	714	0.9 (0.4–1.9)
Appreciation for Indigenous cultures:												
Low (ref)	464	---	608	---	732	---	734	---	749	---	709	---
High	964	1.1 (0.9–1.4)	1,217	1.0 (0.8–1.3)	1,444	0.5 (0.3–0.7)*	1,443	0.4 (0.2–0.7)*	1,465	0.4 (0.2–0.9)*	1,405	0.5 (0.3–0.7)*
Exposure to racial insults												
Never (ref)	961	---	1,172	---	1,363	---	1,358	---	1,375	---	1,328	---
Ever	467	1.3 (1.0–1.6)	653	1.6 (1.2–2.1)*	813	0.8 (0.5–1.4)	819	1.1 (0.5–2.6)	839	1.5 (0.5–4.3)	786	1.2 (0.7–1.8)
DEMOGRAPHIC VARIABLES												
Sex												
Girls (ref)	823	---	1,078	---	1,217	---	1,208	---	1,229	---	1,181	---
Boys	605	1.3 (1.0–1.7)	747	1.7 (1.3–2.3)*	959	2.1 (1.2–3.8)*	969	1.1 (0.5–2.1)	985	0.2 (0.1–0.7)*	933	1.7 (1.1–2.7)*
Age in years												
14–15 years (ref)	1,042	---	1,323	---	1,547	---	1,549	---	1,574	---	1,509	---
16–18 years	386	1.3 (1.0–1.7)	502	1.3 (1.0–1.7)	629	1.4 (0.8–2.2)	628	0.9 (0.4–2.3)	640	0.7 (0.2–2.3)	605	1.0 (0.6–1.6)
Religion												
Catholic (ref)	1,206	---	1,552	---	1,863	---	1,866	---	1,896	---	1,812	---
Evangelical	167	0.7 (0.4–1.1)	199	0.5 (0.3–0.9)*	227	0.8 (0.3–1.8)	225	0.7 (0.2–2.1)	231	1.5 (0.4–5.8)	221	0.8 (0.4–1.6)

	Alcohol Use						Incident Drug Use in One Year					
	Current Drinking (past 30 days) N=1,428		Current Binge Drinking N=1,825		Marijuana N=2,176		Inhalants N=2,177		Cocaine N=2,214		Any Drug N=2,114	
	n	OR (95% CI)	n	OR (95% CI)	n	OR (95% CI)	n	OR (95% CI)	n	OR (95% CI)	n	OR (95% CI)
Other	55	0.5 (0.3–0.9)*	74	0.8 (0.4–1.6)	86	0.6 (0.1–2.8)	86	2.3 (0.7–7.8)	87	11.9 (4.5–31.4)*	81	1.9 (0.7–5.0)
<i>Poverty</i> No (ref)	1,093	---	1,395	---	1,658	---	1,663	---	1,691	---	1,610	---
Yes	335	0.8 (0.6–0.9)*	430	0.8 (0.6–1.1)	518	0.7 (0.3–1.4)	514	1.0 (0.4–2.5)	523	0.9 (0.3–2.5)	504	0.9 (0.5–1.6)
PSYCHOSOCIAL VARIABLES												
<i>Positive expectations for the future</i> Always (ref)	1,024	---	1,306	---	1,541	---	1,542	---	1,566	---	1,500	---
Almost always/sometimes	85	0.9 (0.5–1.5)	121	1.6 (0.9–2.8)	168	0.7 (0.3–2.0)	168	1.2 (0.5–2.9)	173	3.3 (0.9–11.4)	161	0.8 (0.3–1.7)
Never/almost never	319	0.8 (0.7–1.0)	398	1.0 (0.7–1.5)	467	0.7 (0.4–1.2)	467	1.5 (0.5–4.4)	475	3.6 (1.3–9.9)*	453	1.3 (0.7–2.3)
<i>Parental support</i> Never/almost never (ref)	116	---	159	---	204	---	196	---	204	---	188	---
Sometimes	304	1.0 (0.6–1.6)	402	1.2 (0.7–2.0)	488	0.6 (0.3–1.3)	493	1.0 (0.3–3.6)	506	1.1 (0.3–3.9)	472	0.7 (0.3–1.5)
Always/almost always	1,008	0.7 (0.5–1.1)	1,264	0.9 (0.6–1.5)	1,484	0.3 (0.1–0.5)*	1,488	0.9 (0.3–2.3)	1,504	1.3 (0.4–4.1)	1,454	0.4 (0.2–0.7)*
<i>Thrill seeking orientation</i> Low (ref)	1,305	---	1,639	---	1,918	---	1,921	---	1,943	---	1,873	---
High	123	2.1 (1.4–3.1)*	186	1.9 (1.3–2.8)*	258	1.5 (0.9–2.7)	256	5.0 (2.4–10.4)*	271	5.7 (2.1–15.0)*	241	1.5 (0.8–2.8)
<i>Identification with defiant role model</i> None (ref)	959	---	1,146	---	1,310	---	1,304	---	1,315	---	1,279	---
Any	469	1.4 (1.1–1.9)*	679	1.6 (1.2–1.9)*	866	1.6 (0.9–3.0)	873	4.9 (2.4–10.1)*	899	1.8 (0.8–4.2)	835	1.9 (1.1–3.5)*
<i>Identification with conforming role model</i> Low (ref)	421	---	564	---	673	---	669	---	690	---	645	---
High	1,007	0.8 (0.6–1.1)	1,261	0.8 (0.6–1.1)	1,503	1.4 (0.8–2.6)	1,508	0.5 (0.2–1.1)	1,524	0.3 (0.2–0.7)*	1,469	1.2 (0.7–2.0)

* p<0.05;

All models also adjusted for school attendance, symptoms of depression and body image

OR = odds ratios