Published in final edited form as:

J Biol Inorg Chem. 2017 April ; 22(2-3): 395–405. doi:10.1007/s00775-017-1436-5.

Oxygen Activation by Mononuclear Nonheme Iron Dioxygenases Involved in the Degradation of Aromatics

Yifan Wang¹, Jiasong Li¹, and Aimin Liu¹

¹Department of Chemistry, University of Texas at San Antonio, San Antonio, TX 78249

Abstract

Molecular oxygen is utilized in numerous metabolic pathways fundamental for life. Mononuclear nonheme iron-dependent oxygenase enzymes are well-known for their involvement in some of these pathways, activating O₂ so that oxygen atoms can be incorporated into their primary substrates. These reactions often initiate pathways that allow organisms to use stable organic molecules as sources of carbon and energy for growth. From the myriad of reactions in which these enzymes are involved, this perspective recounts the general mechanisms of aromatic dihydroxylation and oxidative ring cleavage, both of which are ubiquitous chemical reactions found in life-sustaining processes. The organic substrate provides all four electrons required for oxygen activation and insertion in the reactions mediated by extradiol and intradiol ring-cleaving catechol dioxygenases. In contrast, two of the electrons are provided by NADH in the *cis*-dihydroxylation mechanism of Rieske dioxygenases. The catalytic nonheme Fe center, with the aid of active site residues, facilitates these electron transfers to O₂ as key elements of the activation processes. This review discusses some general questions for the catalytic strategies of oxygen activation and insertion into aromatic compounds employed by mononuclear nonheme iron-dependent dioxygenases. These include: (1) how oxygen is activated, (2) whether there are common intermediates before oxygen transfer to the aromatic substrate, and (3) are these key intermediates unique to mononuclear nonheme iron dioxygenases?

Graphical abstract

Keywords

Catalytic strategies; Crystal structure; High-valent iron species; Metabolism; Nonheme iron enzymes; Oxidative degradation; Reactive Oxygen Species; Ring-cleaving dioxygenase; Spectroscopy

INTRODUCTION

Discoveries of O₂ incorporation pioneered the new field of oxygenases

Oxygen often serves as a direct oxidant in biodegradation pathways of functionally relevant organic metabolites. Molecular oxygen is a stable diradical molecule with two unpaired electrons, giving it a triplet ground state. In contrast, most organic molecules have singlet ground states with no unpaired electrons, so the reaction with triplet O₂ is a spin forbidden process. Oxygenase enzymes are utilized in nature to activate O₂ and sometimes also the organic molecule to allow them to react. The most common strategies employed by oxygenases are to utilize either a transition metal center or a flavin cofactor to activate O₂ in a controlled manner, so that it can react rapidly with the organic substrate bound nearby in the enzyme active site.

It is not unusual for the first discovered phenomenon to be thought of as prevalent or predominate, thereby creating an intangible barrier to acceptance of other alternative scenarios. Often times, the truth is not what was originally assumed, or only partially illuminated by the initial finding. For instance, before the 1950's it was thought that triplet dioxygen was too stable to be easily incorporated into organic substrates, making water the logical choice for oxygen found in products. For example in carbohydrate metabolism, only oxygen from water is found in the CO₂ that results from the enzymatic oxidation of glucose *via* glycolysis, pyruvate dehydrogenase complex, and the citric acid cycle. While it is true that O₂ ultimately plays a key role in the production of energy rich molecules downstream of these processes, it only serves as a terminal electron acceptor at the terminus of the oxidative phosphorylation pathway. This understanding was well established in the literature, thus it was thought that in most natural transformations, the source of oxygen is water. However, this concept was shown to be incorrect by the elegant and independent isotopic labeling studies performed by Osamu Hayaishi and Howard S. Mason [1, 2], which demonstrated that oxygen from ¹⁸O₂ can be the direct source of oxygen incorporated during biological oxidations of organic substrates.

In the classic dioxygen labeling experiment, Hayaishi showed in 1955 that both oxygen atoms from ¹⁸O₂ were incorporated into the ring-opened dicarboxylic acid product of catechol 1,2-dioxygenase (pyrocatechase) [1]. During the same period, Mason showed that ¹⁸O₂ is the source of the oxygen incorporated by tyrosinase during the formation of 3,4-dihydroxy-L-phenylalanine (DOPA) [2]. Two years later, Hayaishi showed that the first step of tryptophan degradation involves the incorporation of two oxygen atoms to form *N*-formylkynurenine. By mass spectrometric analysis of the isolated product, Hayaishi and coworkers determined that both atoms of oxygen in *N*-formylkynurenine were derived from ¹⁸O₂ rather than H₂¹⁸O [3, 4]. These exciting discoveries, along with subsequent

findings by peers, established oxygenases as a new enzyme class. Hence, a new chapter of biological oxygen activation was started some sixty years ago.

Nonheme iron-dependent oxygenases are prevalent in nature

The most common transition metal-containing oxygenases employ an iron cofactor for oxygen activation. After activation, one or both oxygen atoms are transferred to the organic substrate, consequently generating biological functionality. Oxygenases are broadly classified into two families, dioxygenases and monooxygenases. Dioxygenases incorporate both oxygen atoms into the substrate or a primary organic substrate plus a co-substrate. They either take all four electrons needed for the reaction from the substrate (or cosubstrate) or two from the substrate and two from an external donor. In contrast, monooxygenases only incorporate one oxygen atom from O₂ into the product while reducing the second atom to water using electrons from an external donor. Both oxygenase classes differ from oxidase enzymes, which transfer two or four electrons from a donor (usually the substrate) to reduce O₂ to H₂O₂ or H₂O without oxygen incorporation. Iron-dependent dioxygenases can employ either heme and nonheme cofactors in the active site. The heme-dependent dioxygenases are discussed in our recent review [5] and by Raven in this volume [6] and thus will not be included here.

In what follows, the mechanisms of three types of mononuclear nonheme iron-dependent dioxygenases will be discussed, namely extradiol dioxygenases, intradiol dioxygenases, and Rieske dioxygenases. Several representative enzymes from each class are shown in Table I. We consider these three classes of mononuclear nonheme Fe dioxygenases to be able to promote “pure” dioxygenation reactions. The α -ketoglutarate-dependent dioxygenases and pterin-dependent aromatic amino acid hydroxylases are also prevalent and versatile in nature. However, in the latter two cases, only one atom of O₂ is incorporated into the primary organic substrate, while the other O-atom ends up on the organic cosubstrate, which provides two of the four electrons needed for oxygen reduction. These latter types of mononuclear nonheme Fe dioxygenases are discussed elsewhere in this special issue [7].

RIESKE DIOXYGENASES REQUIRE BOTH O₂ AND A REDUCING AGENT

A unique group of dioxygenases

Under aerobic conditions, Rieske-type iron-dependent dioxygenases transfer two oxygen atoms to arenes in a *cis*-dihydroxylation reaction to yield *cis*-dihydrodiol products with NADH and molecular oxygen as co-substrates (Table 1). Common examples are benzene dioxygenase, phthalate dioxygenase, toluene dioxygenase and naphthalene 1,2-dioxygenase [8]. These enzymes carry out the first step in the degradation of aromatic compounds by soil bacteria, followed by formation of catechol and then ring cleavage by intra- or extradiol dioxygenases [9, 10]. Rieske dioxygenases exhibit stereo- and enantiospecificity in their reactions, which are difficult to be achieved in synthetic organic chemistry [11].

Two or three protein components comprise the majority of Rieske dioxygenases: a reductase containing flavin and ferredoxin cofactors; sometimes a ferredoxin-containing electron transfer protein, and a terminal oxygenase. The latter includes a Rieske [2Fe-2S] cluster,

coordinated by two cysteines and two histidines, and a catalytic mononuclear iron center responsible for dioxygen activation and the *cis*-dihydroxylation of the aromatic ring.

The ~35 kD reductase component acts to transfer electrons one at a time from the two-electron donor NADH, via the small ~15 kD ferredoxin (when presents) to the Rieske [2Fe-2S] cluster found in the terminal oxygenase (MW = 150 – 220 kDa). The relatively large terminal oxygenase generally has a quaternary structure consisting of either an α_3 or $(\alpha\beta)_3$ trimer [11]. Interestingly, crystal structures show that the catalytic iron center and the Rieske cluster within the same subunit are separated by 44 Å [12], beyond the limit for an anticipated efficient electron transfer. The longest separation distance in other systems is 30 – 35 Å found in ribonucleotide reductase [13, 14]. On the other hand, the head-to-tail trimeric structure of the oxygenase component places the catalytic Fe center only 12 Å from the Rieske cluster on the adjacent subunit, which would allow fast electron transfer. Indeed, fast electron transfer between the metal centers has been reported [12]. The pathway for intersubunit electron transfer is postulated to proceed through a conserved Asp residue at the subunit-subunit interface that serves as a bridge between a histidine residue on the Rieske cluster and a histidine ligand on the iron center [15].

Proposed mechanism of Rieske dioxygenase action

In the as-isolated enzyme, the nonheme iron center is typically found in the ferrous state coordinated by two water molecules, two histidines and a bidentate aspartate/glutamate residue [15, 16]. As the first step of the catalytic cycle, the arene substrate binds to the active site adjacent to the catalytic ferrous center (Figure 1) and causes the dispatch of one of the water ligands to make the ferrous center 5-coordinate [17]. Reduction of the Rieske center causes allosteric changes in the geometry of the mononuclear iron center geometry in preparation for O₂ binding [18–20]. Oxygen is then proposed to coordinate to form a transient but yet unobserved Fe(III)-superoxo intermediate [21]. There are several possibilities for the next intermediate in the cycle. It has recently been proposed that the initially formed Fe(III)-superoxo intermediate attacks the aromatic substrate to form an Fe(III)-peroxo-aryl radical intermediate, which is then reduced by electron transfer [21] from the Rieske cluster to form the Fe(II) analog. This proposal is based on the observation that the substrate type and the electron donating capacity of aromatic ring substituents affect the rate constant for electron transfer from the Rieske center, so that electron transfer cannot precede attack on the substrate. Older mechanistic studies suggest that electron transfer results in a side-on bound bidentate hydroperoxide species [11, 22]. In support of the latter hypothesis, side-on bound Fe(III)-hydroperoxo intermediates have been observed in crystals of Rieske dioxygenases and in samples taken during “peroxide shunt” turnover in which hydrogen peroxide serves as the source of both oxygen and electrons [11, 21, 23].

Some possible mechanistic routes to *cis*-diol formation in Rieske dioxygenases are presented in Figure 1. If the intermediate in Rieske dioxygenases is the proposed Fe(II)-peroxo-aryl radical, it might evolve through homolytic cleavage of the O-O bond, leading to formation of a substrate epoxide and Fe(III)-OH. Opening of the epoxide would yield a cation on the adjacent carbon which could abstract the hydroxide from the iron to complete the formation of the non-aromatic *cis*-diol product [21]. If the Fe(III)-bound hydroperoxo species is the

reactive intermediate, it could function as the direct oxidant, transferring both oxygen atoms simultaneously to the aromatic substrate, perhaps *via* a dioxetane intermediate [11]. Two other plausible reaction routes have been proposed based on how the O-O bond is cleaved [24]. Homolytic cleavage followed by substrate oxidation would yield a substrate phenol radical and enzyme-based Fe(IV)=O species, which could transfer the second oxygen to the substrate ring in a subsequent step. The Fe(IV)=O species is chemically equivalent to compound II in most P450 catalysis [25, 26] and heme-dependent tryptophan 2,3-dioxygenase [27, 28]. Alternatively, heterolytic O-O bond cleavage would generate an HO-Fe(V)=O complex, analogous to compound I in heme enzymes, to react with the substrate to give rise to an Fe(IV)-substrate radical complex [23]. The final intermediate in all proposed mechanisms is a ferric product *cis*-diol complex. The product is released only after an external electron is transferred to the mononuclear iron *via* the Rieske cluster to regenerate the resting state.

Studies of biomimetic catalysts have provided insight into how a water ligand might facilitate the reaction of an Fe(III)-OOH intermediate. Based on isotopic labeling experiments, the Que group has proposed a water-assisted mechanism in which that H₂¹⁸O binds to the ferric center *cis* to the hydroperoxo ligand to incorporate an ¹⁸O into an H¹⁸O-Fe(V)=O oxidant that is generated *via* heterolytic cleavage of the O-O bond [29–31]. This mechanism was more recently supported by kinetic studies showing the decay of an observed Fe(III)-OOH intermediate to be accelerated as a function of water concentration [31]. An observed H₂O/D₂O kinetic isotope effect of 2.5 further demonstrated the key role of a water proton in facilitating the proposed heterolytic O-O bond cleavage step. In strong support, Costas and co-workers have reported convincing spectroscopic evidence for the existence of the [Fe^V(O)(OH)(L)]²⁺ ion by cryospray mass spectrometry [32, 33]. More mechanistic details are discussed in a review on bio-inspired nonheme iron catalysis in this issue [34]. Thus, the notion of an Fe^V(O) oxidant presented in Figure 1 for the Rieske oxygenase mechanism has a plausible basis.

OXYGEN INSERTION INTO CATECHOL: A TALE OF TWO DIOXYGENASES

Extradiol and intradiol ring-cleaving dioxygenases

It is estimated that roughly 30% of the mass of woody plant material on earth is comprised of cross-linked aromatic compounds in the form of lignin. Nature has devised a variety of strategies to break down the lignin into individual aromatic molecules, which are much more stable to degradation than aliphatic hydrocarbons [35, 36]. One strategy devised by nature to overcome the stability of aromatic compounds is to utilize oxygenases to couple ring cleavage with oxygen insertion and form ring-opened products that can re-enter the Krebs cycle.

Hashimoto and Hayaishi isolated the first intradiol ring-cleaving dioxygenase, pyrocatechase, from *Pseudomonas* (Figure 2). This enzyme breaks the catechol aromatic ring between the hydroxyl-bearing carbons with incorporation of both atoms of oxygen from O₂ to form of *cis*, *cis*-muconate [1, 37, 38]. Protocatechuate-3,4-dioxygenase, first studied by Stanier and Ingraham [39], and later by Hayaishi [40], converts 3,4-dihydroxybenzoate to 3-carboxy-*cis*,*cis*-muconate in a reaction analogous to that of catechol 1,2-dioxygenase.

Protocatechuate 4,5-dioxygenase and catechol 2,3-dioxygenase (metapyrocatechase) were the first enzymes shown to insert both atoms of oxygen from O₂ into enzyme-bound catecholic substrates at positions adjacent to the two phenolic hydroxyl groups, leading to the production of 2-hydroxymuconate semialdehydes [38, 41, 42]. Senoh and his coworkers found another enzyme, 3,4-dihydroxyphenylacetate-2,3-dioxygenase (also known as homoprotocatechuate 2,3-dioxygenase, 2,3-HPCD) [43], that performs a similar reaction oxidizing 3,4-dihydroxyphenylacetate to 2-hydroxy-5-carboxymethylmuconate semialdehyde (Table 1). Due to the position of ring cleavage, these enzymes were termed extradiol ring-cleaving dioxygenases (Figure 2).

Both intradiol and extradiol dioxygenases operate by utilizing a nonheme Fe atom to allow triplet O₂ to react rapidly with singlet substrates, but they use different mechanistic strategies. Both mechanisms progress from an initial chelate complex of the catecholic substrates with the iron. Both classes have a third iron ligand site adjacent to the substrate ligand sites that can be occupied by oxygen or an activated oxygen intermediate at some stage of the reaction cycle. The mechanism for Fe(II)-dependent extradiol dioxygenases involves nucleophilic attack by an iron-bound oxygen on the substrate to yield an alkyl peroxy intermediate. In the case of the Fe(III)-dependent intradiol dioxygenases, the iron serves to activate substrate to allow formation of a similar alkylperoxy intermediate *via* a different strategy. No additional electron source is required in either of these reactions. The catalytic Fe center, with aid from active site residues, is sufficient to promote electron transfer from the organic substrate to O₂.

Extradiol ring-cleaving dioxygenase genes are more commonly found in the NCBI protein database (<https://www.ncbi.nlm.nih.gov/protein>) than their intradiol counterparts, with 126,278 entries versus 9,994 (both with redundancies) as of November 1st, 2016, respectively. Thus, the extradiol ring-cleaving activities predominate, and they are essential biological catalysts for carbon and nitrogen cycles in the aerobic world.

Proposed mechanism of extradiol ring-cleaving dioxygenases

Extradiol dioxygenases cleave a wide variety of substrates [44–51]. There are three subfamilies of extradiol dioxygenases that differ in their protein structures and evolutionary relationships [52, 53]. Type I and II enzymes share partial sequence similarity. Type II enzymes are believed to be evolved from a type I enzyme through gene duplication [52]. As a result, the type II enzymes are composed of two domains with approximately the same folding pattern as type I [54, 55]. The well-studied catechol 2,3-dioxygenase and 2,3-HPCD belong to type I enzymes, while 2,3-dihydroxybiphenyl dioxygenase typifies type II enzymes. Type III enzymes belong to the functionally diverse Cupin superfamily [56–59]. These enzymes are far removed evolutionarily from the other groups of extradiol dioxygenases, and they are distinct in sequences as well as structures. 3-Hydroxyanthranilate-3,4-dioxygenase (HAO, also known as HAD) and protocatechuate 4,5-dioxygenase (4,5-PCD) are representative members of this group [60, 61]. However, type III enzymes do share most of the active-site characteristics and mechanistic properties of the other two types of enzymes [59, 61, 62].

Each type of extradiol dioxygenase has been well studied, especially the Type III 4,5-PCD and there is considerable evidence with which to build a consensus chemical mechanism as illustrated in Figure 3 [24, 44–46, 48–51, 63]. In particular, Fe-bound superoxo, alkylperoxo, gem-diol intermediates and product-bound enzyme portrayed in this reaction cycle have all been captured from 2,3-HPCD and characterized by X-ray crystallography in the Lipscomb laboratory [64, 65]. Notably, this work is fully supported by spectroscopic data [66–70]. The catalytic iron center of extradiol dioxygenases employs the 2-His-1-carboxylate facial triad structural platform [71]. The ferrous ion is coordinated by two histidines and one glutamate with labile water-derived ligands occupying other empty positions [54, 72]. The glutamate ligand in some enzymes binds in a bidentate mode to the catalytic iron center. In an ordered binding sequence (Figure 3), the organic substrate first binds to the ferrous center and replaces water ligand(s). A proton from only one of the two hydroxyl groups on the substrate is subtracted by an identified active site base, causing asymmetric binding of the catecholic hydroxyl groups to the iron center. Thus, the binding of the organic substrate reorganizes the iron center and promotes dioxygen binding and subsequent reactions [24, 47, 49, 59]. The dioxygen binds to the ferrous ion and may form a transient Fe(III)-bound superoxo radical. The Fe(III) ion then activates the catechol substrate by accepting an electron to generate the substrate semiquinone radical-Fe(II)-superoxo species. The two activated substrates undergo a radical-radical recombination reaction, forming an Fe(II)-bound alkylperoxo intermediate (Figure 3) [73]. In 2,3-HPCD, His200 acts as an active site base to promote formation of the alkylperoxo intermediate [67].

In support of this mechanism, an Fe(III)-superoxo radical has been trapped and spectroscopically characterized in a variant of 2,3-HPCD in which a key active site acid/base catalyst has been removed to slow the reaction chemistry [66]. The actual Fe(II)-superoxo species has been structurally characterized through *in crystallo* chemical reactions in both 2,3-HPCD and homogentisate 1,2-dioxygenase (HGDO), along with the alkylperoxo intermediate [64, 74]. The structures are similar, although, the O₂ is bound *trans* to the oxygen of glutamate in 2,3-HPCD in the alkylperoxo intermediate, whereas O₂ is observed *trans* to a His ligand in HGDO [74]. This variance appears to be due to the difference in the active site geometry of the ternary complex in the two enzymes.

Computational studies suggest that O-O bond cleavage of the alkylperoxo intermediate results in formation of a short-lived gem-diol radical at a ring carbon and then an epoxide at the position of ring cleavage [75–78]. The epoxide undergoes a rearrangement to form a seven-membered ϵ -lactone intermediate, which is hydrolyzed by the Fe(II)-hydroxide to produce the extradiol product. The product-bound intermediate has been structurally defined in both 2,3-HPCD and HGDO. The gem-diol intermediate is structurally characterized in a crystal the the E323L variant of 2,3 HPCD containing the alternative substrate 4-sulfonylcatechol [65]. The lactone intermediate has also been proposed based on the ¹⁸O-labeling experiment of 2,3-dihydroxyphenylpropionate 1,2-dioxygenase (MhpB)[79]. The hydrolysis of a lactone by the iron(II)-bound hydroxide is supported experimentally by the MhpB-catalyzed hydrolysis of a saturated lactone analog [79]. Another possibility for the O-O bond cleavage is *via* 1,2-alkyl migration, which was supported by using mechanistic probes [80] and also hybrid DFT studies [81]. This mechanism suggested that Criegee rearrangement involves concerted acid catalysis by a base residue, which deprotonates the

C-3 hydroxyl group to promote a ring expansion, forming the seven-membered ϵ -lactone intermediate by 1,2-alkyl migration [82].

Proposed mechanism of intradiol ring-cleaving dioxygenases

The mononuclear Fe(III) cofactor of the intradiol catechol dioxygenases is coordinated by two histidine and two tyrosinate residues with a fifth hydroxide ligand in a trigonal bipyramidal structure [83] (Figure 4). When the organic substrate binds to the enzyme, the axial tyrosinate ligand and equatorial hydroxide ligand of the Fe(III) center are displaced [84, 85]. The mechanism of the intradiol catechol dioxygenases has been proposed to be initiated *via* a substrate activation mechanism. The Fe(III) ion polarizes the substrate, building up the charge on one of the hydroxyl bearing carbons. MCD and IR spectroscopies have revealed numerous charge transfer interactions between the substrate and the Fe(III) [86]. One of these interactions provides a low energy path for electron transfer from the substrate to one of the d-orbitals of the iron, but actual electron transfer only occurs during a concerted O₂ binding process. Oxygen accepts one electron directly from the substrate and the second from an occupied d-orbital of the iron different from that which is accepting an electron from the substrate. As a result, two electrons of different spin are transferred to the incoming oxygen to form the alkylperoxo intermediate in one spin-allowed process [87, 88]. The collapse of the alkylperoxo intermediate and O-O bond cleavage occurs *via* a Criegee rearrangement to yield a cyclic anhydride intermediate and Fe(III)-bound hydroxide [89]. Then the anhydride intermediate undergoes hydrolysis by the Fe(III)-OH to produce muconic acid.

The reaction mechanisms of extradiol and intradiol catechol dioxygenases are both believed to proceed through alkylperoxo intermediates. However, the cleavage of the aromatic ring performed by these two enzymes is likely to diverge at this stage of the cycle due to the difference in the detailed structures of the intermediates. The crystal structures of both types of intermediates have been solved [64, 87]. Extradiol dioxygenases form the alkylperoxo intermediate with both of the catecholic oxygens bound to the iron. In contrast, one of the hydroxyl groups is released when the intermediate forms during the intradiol catalytic cycle. This change alters the alignment of the O-O bond of the peroxo with the bonds of the substrate ring and directs the insertions into the aligned bonds. Extradiol cleavage occurs *via* 1,2-alkenyl migration to give a lactone, whereas intradiol cleavage occurs *via* 1,2-acyl migration to give an anhydride [80]. It should be pointed out that the active site residues play critical roles to tune either the early oxygen activation steps or the subsequent O-atom transfer after formation of the common alkylperoxo intermediate. Acid-base catalysis is known to be an important part of the catalytic cycle in extradiol ring cleavage mechanism.

Intradiol oxidative cleavage activity was achieved from a directed evolution of the extradiol cleaving enzyme MhpB [90]. A single mutation of an active site histidine to phenylalanine (*i.e.* H200F) led to the conversion of an extradiol aromatic ring-cleaving 2,3-HPCD protein into an intradiol cleaving enzyme [91], suggesting that from chemical perspective amino acid replacements could swap these two reactions at the enzyme active site.

KEY CATALYTIC INTERMEDIATES of THE DIOXYGENASES

Catechol ring-cleaving dioxygenases are the best characterized non-heme Fe enzymes among the dioxygenases reviewed above. The common catalytic intermediate is an iron-bound alkylperoxo species (Figure 5). Extradiol dioxygenases generate an Fe(II)-alkylperoxo intermediate (Figure 5, 1), while intradiol dioxygenases make an Fe(III)-alkylperoxo intermediate (Figure 5, 2). On the other hand, the Rieske dioxygenases produce an Fe(III)-hydroperoxo oxidant (Figure 5, 3). The three intermediates, although differing from each other, share similarities. The distal oxygen is the first oxygen atom to be transferred to the substrate in the subsequent steps in the catechol ring-cleaving dioxygenases (Figures 3 & 4). This is also a possible avenue for the Rieske dioxygenases through the homolytic O-O bond cleavage pathway (Figure 1), although research on the later reaction steps of the mechanism is needed. However, a closer look indicates that these mononuclear nonheme Fe-bound peroxo intermediates differ significantly in the oxidation state of the metal ion and the interaction between the metal and the substrates.

Recently, an Fe(II)-bound persulfenate intermediate (Figure 5, 4) has been identified in a crystal structure of cysteine dioxygenase (CDO), the best characterized member of the mononuclear nonheme Fe thiol dioxygenase family [92], which includes cysteamine dioxygenase (ADO) [93, 94] and 3-mercaptopropionate dioxygenase (MDO) (Table 1) [95–97]. In this intermediate, the persulfenate is bound to the iron center, forming a three-membered Fe–S–O ring, suggesting that both oxygen atoms of the dioxygen ligand may be transferred to the thiol group simultaneously, followed by an isomerization reaction to yield cysteine sulfinic acid product [92, 98]. The persulfenate intermediate is a unique peroxo intermediate structurally distinct from the carbon-based peroxo intermediates shown in Figure 5. Although the formation of such a persulfenate species is thought to be energetically more costly [99], it was reproducibly generated by at different laboratories in the crystalline state [92, 98, 100]. However, the catalytic competence of the intermediate remains controversial [100]. Evidence is lacking to prove the persulfenate to be an on-pathway intermediate of the CDO catalytic cycle. The same intermediate in solution remains to be trapped and characterized. Nonetheless, this new persulfenate intermediate, at first sight, shares similarities with the peroxo intermediates shown in Figure 5. In the hemolysis of O-O bond catalytic cycle of Rieske dioxygenase an alkylperoxo radical intermediate is proposed (Figure 5, 5). It should be noted that an analogous alkylperoxo radical intermediate is also proposed to be part of the catalytic cycle in heme-dependent tryptophan 2,3-dioxygenase in the form of a 2-indolenylperoxo radical intermediate, (Figure 5, 6) [8], analogous to that recently proposed by Rivard et al. from studies of benzoate 1,2-dioxygenase [21].

CONCLUDING REMARKS

In the past six decades, great strides have been made towards understanding the mechanisms of oxygen activation mediated by mononuclear nonheme Fe dioxygenases. Recent successful trapping and structural characterization of intermediates have helped to establish very detailed chemical understandings of each of the aromatic ring-cleaving dioxygenase enzymes described here as well as of the α -ketoglutarate-dependent dioxygenases [7]. While

there are still uncertainties in the catalytic route for the Rieske dioxygenases, advances in the synthesis and characterization of iron-oxygen model compounds [101–103] have provided challenging new proposals to consider.

From the above survey of the non-heme iron enzyme dioxygenation strategies, a particular emphasis on understanding the general principles of how the iron center interacts with O₂ and the substrate may be reached. The active site residues, including the metal ligands and critical second sphere side chains, have been shown in numerous systems to contribute significantly to the fine tuning of the enzyme activity and substrate specificity. A considerable amount of evidence has also shown that enzyme-substrate spectral signatures are sensitive measures of the electronic structure of the iron environment, and this data can be used to calibrate computational studies and/or validate mechanistic proposals. Several general principles have emerged from these studies: (a) the maintenance of charge at the Fe center is a dominant factor in determining how the substrate binds as well as the relative stability of intermediates, (b) changes in the ligand set, *i.e.* geometry and/or coordination numbers, occur upon substrate binding to prepare the system for subsequent O₂ reactions, (c) the second ligand sphere is often an important factor to stabilize the oxidizing intermediates and direct the oxidizing power, and (d) homolysis or heterolysis of the O-O bond is the major divergence among oxygen transfer strategies and should be a key point of future mechanistic studies.

Looking forward, novel mononuclear nonheme Fe-dependent oxygenation reactions are likely to be discovered. Many are anticipated to share part of the oxygen activation schemes illustrated in this perspective while some catalytic steps may deviate from the common pathway to achieve novel chemical transformations due to the unique enzyme active site architecture and the chemical structure of the substrate. Time-resolved advanced spectroscopy and X-ray crystallography, site-directed alteration (*i.e.* substitution of a natural or unnatural amino acid) or the novel chemical mutagenesis approaches [104–106], and synthetic techniques for mechanism-based probes are expected to yield fundamental structural and chemical insights into O₂ activation and insertion for mechanistic enzymology of the non-heme Fe dioxygenases.

Acknowledgments

The research in our laboratory is supported by the National Science Foundation grants CHE-1623856 and MCB-0843537, the National Institutes of Health grants GM107529, GM108988, and MH107985, and the Litcher Brown Distinguished Chair Endowment fund. We thank Professors Lawrence Que, Jr. and John Lipscomb for helpful discussions of the catalytic mechanisms and editing of the text.

REFERENCE CITED

1. Hayaishi O, Katagiri M, Rothberg S. Mechanism of the pyrocatechase reaction. *J Am Chem Soc.* 1955; 77:5450–5451.
2. Mason HS, Fowlks WL, Peterson E. Oxygen transfer and electron transport by the phenolase complex. *J Am Chem Soc.* 1955; 77:2914–2915.
3. Hayaishi O. From oxygenase to sleep. *J Biol Chem.* 2008; 283:19165–19175. [PubMed: 18390546]
4. Hayaishi O, Rothberg S, Mehler AH, Saito Y. Studies on oxygenases; enzymatic formation of kynurenine from tryptophan. *J Biol Chem.* 1957; 229:889–896. [PubMed: 13502350]

5. Geng J, Liu A. Heme-dependent dioxygenases in tryptophan oxidation. *Arch Biochem Biophys*. 2014; 544:18–26. [PubMed: 24295960]
6. Raven EL. A short history of heme dioxygenases – rise, fall and rise again. *J Biol Inorg Chem*. 2017; 22:XX–XX. O₂ issue.
7. Kal S, Que LJ, Hausinger RP. *J Biol Inorg Chem*. 2017; 22:XX–XX. O₂ issue.
8. Gibson DT. Microbial degradation of aromatic compounds. *Science*. 1967; 161:1093–1097. [PubMed: 5597305]
9. Gibson, DT., Subramanian, V. Microbial degradation of aromatic hydrocarbons. In: Gibson, DT., editor. *Microbial degradation of organic compounds*. Marcel Dekker, Inc; New York: 1984.
10. Seo JS, Keum YS, Li QX. Bacterial degradation of aromatic compounds. *Int J Environ Res Public Health*. 2009; 6:278–309. [PubMed: 19440284]
11. Karlsson A, Parales JV, Parales RE, Gibson DT, Eklund H, Ramaswamy S. Crystal structure of naphthalene dioxygenase: Side-on binding of dioxygen to iron. *Science*. 2003; 299:1039–1042. [PubMed: 12586937]
12. Wolfe MD, Parales JV, Gibson DT, Lipscomb JD. Single turnover chemistry and regulation of O₂ activation by the oxygenase component of naphthalene 1,2-dioxygenase. *J Biol Chem*. 2001; 276:1945–1953. [PubMed: 11056161]
13. Uhlin U, Eklund H. Structure of ribonucleotide reductase protein R1. *Nature*. 1994; 370:533–539. [PubMed: 8052308]
14. Nordlund P, Eklund H. Structure and function of the *Escherichia coli* ribonucleotide reductase protein R2. *J Mol Biol*. 1993; 232:123–164. [PubMed: 8331655]
15. Kauppi B, Lee K, Carredano E, Parales RE, Gibson DT, Eklund H, Ramaswamy S. Structure of an aromatic-ring-hydroxylating dioxygenase-naphthalene 1,2-dioxygenase. *Structure*. 1998; 6:571–586. [PubMed: 9634695]
16. Carredano E, Karlsson A, Kauppi B, Choudhury D, Parales RE, Parales JV, Lee K, Gibson DT, Eklund H, Ramaswamy S. Substrate binding site of naphthalene 1,2-dioxygenase: Functional implications of indole binding. *J Mol Biol*. 2000; 296:701–712. [PubMed: 10669618]
17. Pavel EG, Martins LJ, Ellis WR, Solomon EI. Magnetic circular dichroism studies of exogenous ligand and substrate binding to the non-heme ferrous active site in phthalate dioxygenase. *Chem Biol*. 1994; 1:173–183. [PubMed: 9383387]
18. Yang TC, Wolfe MD, Neibergall MB, Mekmouche Y, Lipscomb JD, Hoffman BM. Substrate binding to NO-ferro-naphthalene 1,2-dioxygenase studied by high-resolution Q-band pulsed ²H-ENDOR spectroscopy. *J Am Chem Soc*. 2003; 125:7056–7066. [PubMed: 12783560]
19. Martins BM, Svetlitchnaia T, Dobbek H. 2-oxoquinoline 8-monooxygenase oxygenase component: Active site modulation by rieske-[2Fe-2S] center oxidation/reduction. *Structure*. 2005; 13:817–824. [PubMed: 15893671]
20. Ohta T, Chakrabarty S, Lipscomb JD, Solomon EI. Near-ir mcd of the nonheme ferrous active site in naphthalene 1,2-dioxygenase: Correlation to crystallography and structural insight into the mechanism of rieske dioxygenases. *J Am Chem Soc*. 2008; 130:1601–1610. [PubMed: 18189388]
21. Rivard BS, Rogers MS, Marell DJ, Neibergall MB, Chakrabarty S, Cramer CJ, Lipscomb JD. Rate-determining attack on substrate precedes rieske cluster oxidation during *cis*-dihydroxylation by benzoate dioxygenase. *Biochemistry*. 2015; 54:4652–4664. [PubMed: 26154836]
22. Ashikawa Y, Fujimoto Z, Usami Y, Inoue K, Noguchi H, Yamane H, Nojiri H. Structural insight into the substrate- and dioxygen-binding manner in the catalytic cycle of rieske nonheme iron oxygenase system, carbazole 1,9a-dioxygenase. *BMC Struct Biol*. 2012; 12:15. [PubMed: 22727022]
23. Neibergall MB, Stubna A, Mekmouche Y, Münck E, Lipscomb JD. Hydrogen peroxide dependent *cis*-dihydroxylation of benzoate by fully oxidized benzoate 1,2-dioxygenase. *Biochemistry*. 2007; 46:8004–8016. [PubMed: 17567152]
24. Bugg TDH, Ramaswamy S. Non-heme iron-dependent dioxygenases: Unravelling catalytic mechanisms for complex enzymatic oxidations. *Curr Opin Chem Biol*. 2008; 12:134–140. [PubMed: 18249197]

25. Tarasev M, Rhames F, Ballou DP. Rates of the phthalate dioxygenase reaction with oxygen are dramatically increased by interactions with phthalate and phthalate oxygenase reductase. *Biochemistry*. 2004; 43:12799–12808. [PubMed: 15461452]
26. Tarasev M, Ballou DP. Chemistry of the catalytic conversion of phthalate into its *cis*-dihydrodiol during the reaction of oxygen with the reduced form of phthalate dioxygenase. *Biochemistry*. 2005; 44:6197–6207. [PubMed: 15835907]
27. Fu R, Gupta R, Geng J, Dornevil K, Wang S, Zhang Y, Hendrich MP, Liu A. Enzyme reactivation by hydrogen peroxide in heme-based tryptophan dioxygenase. *J Biol Chem*. 2011; 286:26541–26554. [PubMed: 21632548]
28. Geng J, Dornevil K, Liu A. Chemical rescue of the distal histidine mutants of tryptophan 2,3-dioxygenase. *J Am Chem Soc*. 2012; 134:12209–12218. [PubMed: 22742206]
29. Chen K, Que L Jr. Stereospecific alkane hydroxylation by non-heme iron catalysts: Mechanistic evidence for an Fe(V)=O active species. *J Am Chem Soc*. 2001; 123:6327–6337. [PubMed: 11427057]
30. Chen K, Costas M, Kim J, Tipton AK, Que L Jr. Olefin *cis*-dihydroxylation versus epoxidation by non-heme iron catalysts: Two faces of an Fe(III)-OOH coin. *J Am Chem Soc*. 2002; 124:3026–3035. [PubMed: 11902894]
31. Oloo WN, Fielding AJ, Que L. Rate-determining water-assisted O-O bond cleavage of an Fe(III)-OOH intermediate in a bio-inspired nonheme iron-catalyzed oxidation. *J Am Chem Soc*. 2013; 135:6438–6441. [PubMed: 23594282]
32. Prat I, Mathieson JS, Guell M, Ribas X, Luis JM, Cronin L, Costas M. Observation of Fe(V)=O using variable-temperature mass spectrometry and its enzyme-like c-h and c=c oxidation reactions. *Nat Chem*. 2011; 3:788–793. [PubMed: 21941251]
33. Lyakin OY, Prat I, Bryliakov KP, Costas M, Talsi EP. Epr detection of Fe(V)=O active species in nonheme iron-catalyzed oxidations. *Catal Commun*. 2012; 29:105–108.
34. Olivo G, Cusso O, Borrell M, Costas M. *J Biol Inorg Chem*. 2017; 22:XX–XX. O₂ issue.
35. Leonowicz A, Matuszewska A, Luterek J, Ziegenhagen D, Wojtas-Wasilewska M, Cho NS, Hofrichter M, Rogalski J. Biodegradation of lignin by white rot fungi. *Fungal Genet Biol*. 1999; 27:175–185. [PubMed: 10441443]
36. Ruiz-Duenas FJ, Martinez AT. Microbial degradation of lignin: How a bulky recalcitrant polymer is efficiently recycled in nature and how we can take advantage of this. *Microb Biotechnol*. 2009; 2:164–177. [PubMed: 21261911]
37. Hayaishi O, Hashimoto K. Pyrocatechase: A new enzyme catalyzing oxidative breakdown of pyrocatechin. *J Biochem (Tokyo)*. 1950; 37:71.
38. Hayaishi O, Katagiri M, Rothberg S. Studies on oxygenases: Pyrocatechase. *J Biol Chem*. 1957; 229:905–920. [PubMed: 13502352]
39. Stanier RY, Ingraham JL. Protocatechuic acid oxidase. *J Biol Chem*. 1954; 210:799–808. [PubMed: 13211618]
40. Hayaishi O. Crystalline oxygenases of *pseudomonads*. *Bacteriol Rev*. 1966; 30:720–731. [PubMed: 5342517]
41. Dagley S, Evans WC, Ribbons DW. New pathways in the oxidative metabolism of aromatic compounds by microorganisms. *Nature*. 1960; 188:560–566. [PubMed: 13719300]
42. Kojima Y, Itada N, Hayaishi O. Metapyrocatechase: A new catechol-cleaving enzyme. *J Biol Chem*. 1961; 236:2223–2228. [PubMed: 13757654]
43. Adachi K, Takeda Y, Senoh S, Kita H. Metabolism of *p*-hydroxyphenylacetic acid in *pseudomonas ovalis*. *Biochim Biophys Acta*. 1964; 93:483–493. [PubMed: 14263147]
44. Lipscomb JD. Mechanism of extradiol aromatic ring-cleaving dioxygenases. *Curr Opin Struct Biol*. 2008; 18:644–649. [PubMed: 19007887]
45. Kovaleva EG, Neibergall MB, Chakrabarty S, Lipscomb JD. Finding intermediates in the O₂ activation pathways of non-heme iron oxygenases. *Acc Chem Res*. 2007; 40:475–483. [PubMed: 17567087]
46. Vaillancourt FH, Bolin JT, Eltis LD. The ins and outs of ring-cleaving dioxygenases. *Crit Rev Biochem Mol Biol*. 2006; 41:241–267. [PubMed: 16849108]

47. Abu-Omar MM, Loaiza A, Hontzeas N. Reaction mechanisms of mononuclear non-heme iron oxygenases. *Chem Rev.* 2005; 105:2227–2252. [PubMed: 15941213]
48. Bugg TDH. Dioxygenase enzymes: Catalytic mechanisms and chemical models. *Tetrahedron.* 2003; 59:7075–7101.
49. Solomon EI, Brunold TC, Davis MI, Kemsley JN, Lee S-K, Lehnert N, Neese F, Skulan AJ, Yang Y-S, Zhou J. Geometric and electronic structure/function correlations in non-heme iron enzymes. *Chem Rev.* 2000; 100:235–349. [PubMed: 11749238]
50. Broderick JB. Catechol dioxygenases. *Essays Biochem.* 1999; 34:173–189. [PubMed: 10730195]
51. Que LJ, Ho RYN. Dioxygen activation by enzymes with mononuclear non-heme iron active sites. *Chem Rev.* 1996; 96:2607–2624. [PubMed: 11848838]
52. Spence EL, Kawamukai M, Sanvoisin J, Braven H, Bugg TD. Catechol dioxygenases from *Escherichia coli* (MhpB) and *Alcaligenes eutrophus* (MpcI): Sequence analysis and biochemical properties of a third family of extradiol dioxygenases. *J Bacteriol.* 1996; 178:5249–5256. [PubMed: 8752345]
53. Eltis LD, Bolin JT. Evolutionary relationships among extradiol dioxygenases. *J Bacteriol.* 1996; 178:5930–5937. [PubMed: 8830689]
54. Han S, Eltis LD, Timmis KN, Muchmore SW, Bolin JT. Crystal structure of the biphenyl-cleaving extradiol dioxygenase from a PCB-degrading *pseudomonad*. *Science.* 1995; 270:976–980. [PubMed: 7481800]
55. Senda T, Sugiyama K, Narita H, Yamamoto T, Kimbara K, Fukuda M, Sato M, Yano K, Mitsui Y. Three-dimensional structures of free form and two substrate complexes of an extradiol ring-cleavage type dioxygenase, the bphc enzyme from *pseudomonas sp.* Strain KKS102. *J Mol Biol.* 1996; 255:735–752. [PubMed: 8636975]
56. Dunwell JM. Cupins: A new superfamily of functionally diverse proteins that include germins and plant storage proteins. *Biotechnol Genet Eng Rev.* 1998; 15:1–32. [PubMed: 9573603]
57. Khuri S, Bakker FT, Dunwell JM. Phylogeny, function, and evolution of the cupins, a structurally conserved, functionally diverse superfamily of proteins. *Mol Biol Evol.* 2001; 18:593–605. [PubMed: 11264412]
58. Dunwell JM, Culham A, Carter CE, Sosa-Aguirre CR, Goodenough PW. Evolution of functional diversity in the cupin superfamily. *Trends Biochem Sci.* 2001; 26:740–746. [PubMed: 11738598]
59. Fetzner S. Ring-cleaving dioxygenases with a cupin fold. *Appl Environ Microbiol.* 2012; 78:2505–2514. [PubMed: 22287012]
60. Colabroy KL, Zhai H, Li T, Ge Y, Zhang Y, Liu A, Ealick SE, McLafferty FW, Begley TP. The mechanism of inactivation of 3-hydroxyanthranilate-3,4-dioxygenase by 4-chloro-3-hydroxyanthranilate. *Biochemistry.* 2005; 44:7623–7631. [PubMed: 15909977]
61. Liu F, Geng J, Gumpfer RH, Barman A, Davis I, Ozarowski A, Hamelberg D, Liu A. An iron reservoir to the catalytic metal: The rubredoxin iron in an extradiol dioxygenase. *J Biol Chem.* 2015; 290:15621–15634. [PubMed: 25918158]
62. Zhang Y, Colabroy KL, Begley TP, Ealick SE. Structural studies on 3-hydroxyanthranilate-3,4-dioxygenase: The catalytic mechanism of a complex oxidation involved in nad biosynthesis. *Biochemistry.* 2005; 44:7632–7643. [PubMed: 15909978]
63. Bugg TDH, Lin G. Solving the riddle of the intradiol and extradiol catechol dioxygenases: How do enzymes control hydroperoxide rearrangements? *Chem Commun.* 2001; 11:941–952.
64. Kovaleva EG, Lipscomb JD. Crystal structures of Fe²⁺ dioxygenase superoxo, alkylperoxo, and bound product intermediates. *Science.* 2007; 316:453–457. [PubMed: 17446402]
65. Kovaleva EG, Lipscomb JD. Intermediate in the O-O bond cleavage reaction of an extradiol dioxygenase. *Biochemistry.* 2008; 47:11168–11170. [PubMed: 18826259]
66. Mbughuni MM, Chakrabarti M, Hayden JA, Bominaar EL, Hendrich MP, Münck E, Lipscomb JD. Trapping and spectroscopic characterization of an Fe(III)-superoxo intermediate from a nonheme mononuclear iron-containing enzyme. *Proc Natl Acad Sci U S A.* 2010; 107:16788–16793. [PubMed: 20837547]
67. Groce SL, Lipscomb JD. Aromatic ring cleavage by homoprotocatechuate 2,3-dioxygenase: Role of His200 in the kinetics of interconversion of reaction cycle intermediates. *Biochemistry.* 2005; 44:7175–7188. [PubMed: 15882056]

68. Mbughuni MM, Chakrabarti M, Hayden JA, Meier KK, Dalluge JJ, Hendrich MP, Münck E, Lipscomb JD. Oxy intermediates of homoprotocatechuate 2,3-dioxygenase: Facile electron transfer between substrates. *Biochemistry*. 2011; 50:10262–10274. [PubMed: 22011290]
69. Kovaleva EG, Lipscomb JD. Structural basis for the role of tyrosine 257 of homoprotocatechuate 2,3-dioxygenase in substrate and oxygen activation. *Biochemistry*. 2012; 51:8755–8763. [PubMed: 23066739]
70. Mbughuni MM, Meier KK, Münck E, Lipscomb JD. Substrate-mediated oxygen activation by homoprotocatechuate 2,3-dioxygenase: Intermediates formed by a tyrosine 257 variant. *Biochemistry*. 2012; 51:8743–8754. [PubMed: 23066705]
71. Hegg EL, Que L Jr. The 2-His-1-carboxylate facial triad. An emerging structural motif in mononuclear non-heme iron(II) enzymes. *Eur J Biochem*. 1997; 250:625–629. [PubMed: 9461283]
72. Vetting MW, Wackett LP, Que L Jr, Lipscomb JD, Ohlendorf DH. Crystallographic comparison of manganese- and iron-dependent homoprotocatechuate 2,3-dioxygenases. *J Bacteriol*. 2004; 186:1945–1958. [PubMed: 15028678]
73. Arciero DM, Lipscomb JD. Binding of ^{17}O -labeled substrate and inhibitors to protocatechuate 4,5-dioxygenase-nitrosyl complex. Evidence for direct substrate binding to the active site Fe^{2+} of extradiol dioxygenases. *J Biol Chem*. 1986; 261:2170–2178. [PubMed: 3003098]
74. Jeoung JH, Bommer M, Lin TY, Dobbek H. Visualizing the substrate-, superoxo-, alkylperoxo-, and product-bound states at the nonheme $\text{Fe}(\text{II})$ site of homogentisate dioxygenase. *Proc Natl Acad Sci U S A*. 2013; 110:12625–12630. [PubMed: 23858455]
75. Siegbahn PEM, Haefner F. Mechanism for catechol ring-cleavage by non-heme iron extradiol dioxygenases. *J Am Chem Soc*. 2004; 126:8919–8932. [PubMed: 15264822]
76. Deeth RJ, Bugg TDH. A density functional investigation of the extradiol cleavage mechanism in non-heme iron catechol dioxygenases. *J Biol Inorg Chem*. 2003; 8:409–418. [PubMed: 12761662]
77. Christian GJ, Ye S, Neese F. Oxygen activation in extradiol catecholate dioxygenases - a density functional study. *Chem Sci*. 2012; 3:1600–1611.
78. Dong G, Shaik S, Lai W. Oxygen activation by homoprotocatechuate 2,3-dioxygenase: A QM/MM study reveals the key intermediates in the activation cycle. *Chem Sci*. 2013; 4:3624–3635.
79. Sanvoisin J, Langley GJ, Bugg TDH. Mechanism of extradiol catechol dioxygenases - evidence for a lactone intermediate in the 2,3-dihydroxyphenylpropionate 1,2-dioxygenase reaction. *J Am Chem Soc*. 1995; 117:7836–7837.
80. Xin M, Bugg TD. Evidence from mechanistic probes for distinct hydroperoxide rearrangement mechanisms in the intradiol and extradiol catechol dioxygenases. *J Am Chem Soc*. 2008; 130:10422–10430. [PubMed: 18627158]
81. Borowski T, Wojcik A, Milaczewska A, Georgiev V, Blomberg MRA, Siegbahn PEM. The alkenyl migration mechanism catalyzed by extradiol dioxygenases: A hybrid dft study. *J Biol Inorg Chem*. 2012; 17:881–890. [PubMed: 22622485]
82. Mendel S, Arndt A, Bugg TDH. Acid-base catalysis in the extradiol catechol dioxygenase reaction mechanism: Site-directed mutagenesis of His-115 and His-179 in *Escherichia coli* 2,3-dihydroxyphenylpropionate 1,2-dioxygenase (MhpB). *Biochemistry*. 2004; 43:13390–13396. [PubMed: 15491145]
83. Ohlendorf DH, Lipscomb JD, Weber PC. Structure and assembly of protocatechuate 3,4-dioxygenase. *Nature*. 1988; 336:403–405. [PubMed: 3194022]
84. Orville AM, Elango N, Lipscomb JD, Ohlendorf DH. Structures of competitive inhibitor complexes of protocatechuate 3,4-dioxygenase: Multiple exogenous ligand binding orientations within the active site. *Biochemistry*. 1997; 36:10039–10051. [PubMed: 9254599]
85. Orville AM, Lipscomb JD, Ohlendorf DH. Crystal structures of substrate and substrate analog complexes of protocatechuate 3,4-dioxygenase: Endogenous $\text{Fe}(\text{III})$ ligand displacement in response to substrate binding. *Biochemistry*. 1997; 36:10052–10066. [PubMed: 9254600]
86. Davis MI, Wasinger EC, Decker A, Pau MYM, Vaillancourt FH, Bolin JT, Eltis LD, Hedman B, Hodgson KO, Solomon EI. Spectroscopic and electronic structure studies of 2,3-dihydroxybiphenyl 1,2-dioxygenase: O_2 reactivity of the non-heme ferrous site in extradiol dioxygenases. *J Am Chem Soc*. 2003; 125:11214–11227. [PubMed: 16220940]

87. Knoot CJ, Purpero VM, Lipscomb JD. Crystal structures of alkylperoxo and anhydride intermediates in an intradiol ring-cleaving dioxygenase. *Proc Natl Acad Sci U S A*. 2015; 112:388–393. [PubMed: 25548185]
88. Pau MY, Lipscomb JD, Solomon EI. Substrate activation for O₂ reactions by oxidized metal centers in biology. *Proc Natl Acad Sci U S A*. 2007; 104:18355–18362. [PubMed: 18003930]
89. Que L Jr, Lipscomb JD, Münck E, Wood JM. Protocatechuate 3,4-dioxygenase. Inhibitor studies and mechanistic implications. *Biochim Biophys Acta*. 1977; 485:60–74. [PubMed: 199266]
90. Schlosrich J, Eley KL, Crowley PJ, Bugg TDH. Directed evolution of a non-heme-iron-dependent extradiol catechol dioxygenase: Identification of mutants with intradiol oxidative cleavage activity. *ChemBioChem*. 2006; 7:1899–1908. [PubMed: 17051653]
91. Groce SL, Lipscomb JD. Conversion of extradiol aromatic ring-cleaving homoprotocatechuate 2,3-dioxygenase into an intradiol cleaving enzyme. *J Am Chem Soc*. 2003; 125:11780–11781. [PubMed: 14505375]
92. Simmons CR, Krishnamoorthy K, Granett SL, Schuller DJ, Dominy JE Jr, Begley TP, Stipanuk MH, Karplus PA. A putative Fe²⁺-bound persulfenate intermediate in cysteine dioxygenase. *Biochemistry*. 2008; 47:11390–11392. [PubMed: 18847220]
93. Dominy JE Jr, Simmons CR, Hirschberger LL, Hwang J, Coloso RM, Stipanuk MH. Discovery and characterization of a second mammalian thiol dioxygenase, cysteamine dioxygenase. *J Biol Chem*. 2007; 282:25189–25198. [PubMed: 17581819]
94. Richerson RB, Ziegler DM. Cysteamine dioxygenase. *Methods Enzymol*. 1987; 143:410–415. [PubMed: 3657558]
95. Bruland N, Wubbeler JH, Steinbuechel A. 3-mercaptopropionate dioxygenase, a cysteine dioxygenase homologue, catalyzes the initial step of 3-mercaptopropionate catabolism in the 3,3-thiodipropionic acid-degrading bacterium *variovorax paradoxus*. *J Biol Chem*. 2009; 284:660–672. [PubMed: 19001372]
96. Tchesnokov EP, Fellner M, Siakkou E, Kleffmann T, Martin LW, Aloï S, Lamont IL, Wilbanks SM, Jameson GN. The cysteine dioxygenase homologue from *pseudomonas aeruginosa* is a 3-mercaptopropionate dioxygenase. *J Biol Chem*. 2015; 290:24424–24437. [PubMed: 26272617]
97. Pierce BS, Subedi BP, Sardar S, Crowell JK. The “Gln-type” thiol dioxygenase from *Azotobacter vinelandii* is a 3-mercaptopropionic acid dioxygenase. *Biochemistry*. 2015; 54:7477–7490. [PubMed: 26624219]
98. Driggers CM, Cooley RB, Sankaran B, Hirschberger LL, Stipanuk MH, Karplus PA. Cysteine dioxygenase structures from pH 4 to 9: Consistent Cys-persulfenate formation at intermediate pH and a Cys-bound enzyme at higher pH. *J Mol Biol*. 2013; 425:3121–3136. [PubMed: 23747973]
99. Kumar D, Thiel W, de Visser SP. Theoretical study on the mechanism of the oxygen activation process in cysteine dioxygenase enzymes. *J Am Chem Soc*. 2011; 133:3869–3882. [PubMed: 21344861]
100. Souness RJ, Kleffmann T, Tchesnokov EP, Wilbanks SM, Jameson GB, Jameson GN. Mechanistic implications of persulfenate and persulfide binding in the active site of cysteine dioxygenase. *Biochemistry*. 2013; 52:7606–7617. [PubMed: 24084026]
101. Hohenberger J, Ray K, Meyer K. The biology and chemistry of high-valent iron-oxo and iron-nitrido complexes. *Nat Commun*. 2012; 3:720. [PubMed: 22395611]
102. Bassan A, Blomberg MR, Siegbahn PE, Que L Jr. Two faces of a biomimetic non-heme HO-Fe(V)=O oxidant: Olefin epoxidation versus *cis*-dihydroxylation. *Angew Chem Int Ed Engl*. 2005; 44:2939–2941. [PubMed: 15812868]
103. Costas M, Tipton AK, Chen K, Jo DH, Que L Jr. Modeling rieske dioxygenases: The first example of iron-catalyzed asymmetric *cis*-dihydroxylation of olefins. *J Am Chem Soc*. 2001; 123:6722–6723. [PubMed: 11439071]
104. Harder D, Hirschi S, Ucurum Z, Goers R, Meier W, Muller DJ, Fotiadis D. Engineering a chemical switch into the light-driven proton pump proteorhodopsin by cysteine mutagenesis and thiol modification. *Angew Chem Int Ed Engl*. 2016; 55:8846–8849. [PubMed: 27294681]
105. Yang A, Ha S, Ahn J, Kim R, Kim S, Lee Y, Kim J, Soll D, Lee HY, Park HS. A chemical biology route to site-specific authentic protein modifications. *Science*. 2016; 354:623–626. [PubMed: 27708052]

106. Bose JL. Chemical and uv mutagenesis. *Methods Mol Biol.* 2016; 1373:111–115. [PubMed: 25646611]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Homolysis of the O–O bond:

Heterolysis of the O–O bond:

Figure 1.

Proposed catalytic routes of Rieske dioxxygenases through hemolysis (upper) and heterolysis (bottom) of O–O bond.

Figure 2. Distinct natural catechol dioxygenase activities. (A) Extradiol and intradiol cleavages, (B) the alkylperoxo intermediate structurally characterized in the extradiol dioxygenase 2,3-HPCD (from 2IGA.pdb), and (C) the structure of the anhydride intermediate in the intradiol dioxygenase 3,4-PCD (from 4WHR.pdb).

Figure 3.
The catalytic mechanism of extradiol ring-cleaving dioxygenases

Figure 4.
The catalytic cycle of intradiol ring-cleaving dioxygenases.

Figure 5.

Comparison of the Fe-bound peroxo intermediates of mononuclear nonheme Fe dioxygenases with the analogous intermediates of a nonheme Fe-dependent thiol dioxygenase and heme-dependent tryptophan 2,3-dioxygenase. Key catalytic intermediates of alkylperoxo intermediate found in extradiol dioxygenase (1) and intradiol dioxygenase (2), a hydroperoxo intermediate in the catalytic route of Rieske dioxygenases (3), a putative persulfenate intermediate in thiol dioxygenase (4), an alkylperoxo radical intermediate proposed for the Rieske dioxygenases (5) and an analogous 2-indolenylperoxo radical intermediate in the catalytic cycle of heme-dependent tryptophan 2,3-dioxygenase (6).

Table 1

Nonheme Fe dioxygenases discussed in this review

Enzyme	Substrate	Product
Rieske Dioxygenase		
Benzene 1,2-dioxygenase
Phthalate 4,5-dioxygenase
Toluene dioxygenase
Naphthalene 1,2-dioxygenase
Thiol Dioxygenase		
Cysteine dioxygenase
3-Mercaptopropionic acid dioxygenase
Cysteamine dioxygenase
Intradiol Dioxygenase		
Catechol 1,2-dioxygenase
Protocatechuate 3,4-dioxygenase
Extradiol Dioxygenase		
Catechol 2,3-dioxygenase
Homoprotocatechuate 2,3-dioxygenase

Enzyme	Substrate	Product
2,3-dihydroxybiphenyl dioxygenase
3-Hydroxyanthranilate 3,4-dioxygenase
Protocatechuate 4,5-dioxygenase
Homogentisate 1,2-dioxygenase
2,3-dihydroxyphenyl propionate 1,2-dioxygenase

The ring cleavage site is indicated by a jagged red line in the intradiol and extradiol dioxygenases.