Published in final edited form as:

Gastroenterology. 2017 April ; 152(5): 983–986.e6. doi:10.1053/j.gastro.2016.12.010.

Germline Mutations in *PALB2*, *BRCA1*, and *RAD51C*, Which Regulate DNA Recombination Repair, in Patients with Gastric Cancer

Ruta Sahasrabudhe^{1,*}, Paul Lott^{1,*}, Mabel Bohorquez², Ted Toal¹, Ana P. Estrada², John J. Suarez², Alejandro Brea-Fernández³, José Cameselle-Teijeiro⁴, Carla Pinto⁵, Irma Ramos⁶, Alejandra Mantilla⁷, Rodrigo Prieto^{2,a}, Alejandro Corvalan⁸, Enrique Norero⁸, Carolina Alvarez⁹, Teresa Tapia⁹, Pilar Carvallo⁹, Luz M. Gonzalez¹⁰, Alicia Cock-Rada¹⁰, Angela Solano¹¹, Florencia Neffa¹², Adriana Della Valle¹², Chris Yau¹³, Gabriela Soares¹⁴, Alexander Borowsky¹⁵, Nan Hu¹⁶, Li-Ji He¹⁷, Xiao-You Han¹⁸, Latin American Gastric Cancer Genetics Collaborative Group¹⁹, Philip R. Taylor¹⁶, Alisa M. Goldstein¹⁶, Javier Torres⁵, Magdalena Echeverry², Clara Ruiz-Ponte³, Manuel R. Teixeira^{5,20}, and Luis G. Carvajal Carmona^{1,21}

¹Genome Center and Department of Biochemistry and Molecular Medicine, School of Medicine, University of California, Davis, USA

²Grupo de Citogenética, Filogenia y Evolución de Poblaciones, Facultades de Ciencias y Facultad de Ciencias de la Salud, Universidad del Tolima, Ibagué, Colombia

³Fundación Pública Galega de Medicina Xenómica (FPGMX), Centro de Investigación Biomédica en Red de Enfermedades Raras (CIBERER), Genomics Medicine Group, Hospital Clínico, 15706 Santiago de Compostela, University of Santiago de Compostela, Galicia, Spain

⁴Servicio de Anatomía Patológica, Hospital Clínico, Santiago de Compostela, Spain

⁵Department of Genetics, Portuguese Oncology Institute of Porto (IPO-Porto), Porto, Portugal

⁶Unidad de Investigación en Enfermedades Infecciosas, UMAE Pediatría, IMSS, México City, México

⁷Departamento de Patología, UMAE Oncología, IMSS, Mexico City, Mexico

Corresponding Author: Luis G. Carvajal-Carmona, PhD Genome Center and Department of Biochemistry and Molecular Medicine School of Medicine, University of California, Davis UC Davis Genome Center, 451 Health Sciences Drive Davis, California 95616, USA Phone: +1 530-752-9654, Fax: +1 530-754-9658 lgcavajal@ucdavis.edu.

*These authors contributed equally to the manuscript

^aIn memoriam

Disclosure: The authors have no conflict of interest to disclose.

Author contribution: Study concept and design = RS, MT, LGCC; Acquisition of data = RS, PL, MB, APE, JJS, ABF, CP, IR, AM, RP, JS, AC, EN, CA, TT, PC, LMG, ACR, AS, FN, ADV, CY, GS, AB, NH, LJH, XYH, PRT, AMG, JT, ME, CRP, MT, LGCC; Analysis and interpretation of data = PL, RS, TT, LGCC; Drafting of the manuscript = RS, LGCC; Critical revision of the manuscript for important intellectual content = RS, PL, TT, AG, JT, ME, JCT, AB, CRP, MT, LGCC; Statistical analysis = RS, PL, TT, LGCC; Obtained funding = LGCC; Administrative, technical, or material support = RS, PL, MT, JT, CRP, ME, LGCC; Study supervision = RS, PL, MT, JT, CRP, ME, LGCC

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

⁸Facultad de Medicina, Pontificia Universidad Católica de Chile, Santiago, Chile

⁹Departamento de Biología Celular y Molecular, Facultad de Ciencias Biológicas, Pontificia Universidad Católica de Chile, Santiago, Chile

¹⁰Universidad de Antioquia, Medellín, Colombia

¹¹INBIOMED, UBA/CONICET y CEMIC, CABA, Argentina

¹²Laboratorio GENIA, Montevideo, Uruguay

¹³Wellcome Trust Centre for Human Genetics, University of Oxford, UK

¹⁴Centro de Genética Médica Dr. Jacinto Magalhães (CGMJM), Centro Hospitalar do Porto, Porto, Portugal

¹⁵Department of Pathology, School of Medicine, University of California, Davis, USA

¹⁶Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, Bethesda, Maryland, USA

¹⁷Yangcheng Cancer Hospital, Yangcheng, Shanxi, PR China

¹⁸Shanxi Cancer Hospital, Taiyuan, Shanxi, PR China

¹⁹Full details listed in the acknowledgements

²⁰Institute of Biomedical Sciences (ICBAS), University of Porto, Porto, Portugal

²¹Fundación de Genómica y Genética Molecular, Colombia

Abstract

Up to 10% of cases of gastric cancer are familial, but so far, only mutations in *CDHI* have been associated with gastric cancer risk. To identify genetic variants that affect risk for gastric cancer, we collected blood samples from 28 patients with hereditary diffuse gastric cancer (HDGC) not associated with mutations in *CDHI* and performed whole-exome sequence analysis. We then analyzed sequences of candidate genes in 333 independent HDGC and non-HDGC cases. We identified 11 cases with mutations in *PALB2*, *BRCA1*, or *RAD51C* genes, which regulate homologous DNA recombination. We found these mutations in 2 of 31 patients with HDGC (6.5%) and 9 of 331 patients with sporadic gastric cancer (2.8%). Most of these mutations had been previously associated with other types of tumors and partially co-segregated with gastric cancer in our study. Tumors that developed in patients with these mutations had a mutation signature associated with somatic homologous recombination deficiency. Our findings indicate that defects in homologous recombination increase risk for gastric cancer.

Keywords

stomach; tumor; WES; interaction

Worldwide, gastric cancer (GC) is the fifth most commonly diagnosed malignancy and the third cause of cancer-related deaths ¹. Up to 10% of cases show familial clustering, suggesting a genetic basis ². *CDHI* mutations are a known cause of hereditary diffuse gastric

cancer (HDGC), explaining ~ 40% of cases^{3,4}, but the genetics of non-HDGC remain largely unknown. To identify novel GC genes, we analyzed *CDHI* mutation-negative HDGC cases using whole exome sequencing (WES) followed by candidate gene targeted analyses in independent HDGC and non-HDGC cases.

WES of 28 *CDHI*-negative European HDGC cases identified three with candidate causal variants (Table 1): nonsense (p.Arg414Ter) and splice site (c.3201+1G>T) *PALB2* mutations, and a nonsense *RAD51C* (p.Arg237Ter) mutation. No deleterious mutations were seen in other known cancer genes (Supplementary methods). *PALB2* and *RAD51C* are both critical in homologous recombination (HR), a major DNA repair pathway⁵. Both of the above *PALB2* mutations have been previously reported as pathogenic in breast cancer (BC) families⁶ and *RAD51C* p.Arg237Ter is reported as pathogenic in ClinVar⁷.

We then performed targeted sequencing of *PALB2* and *RAD51C*, their interaction partners *BRCA1/2* and *CDHI* in 173 additional Latin American GC cases. Based upon enrichment of HR mutations in our discovery cohort and a recent report showing multiple intestinal, diffuse and mixed histology gastric tumors with a somatic HR deficiency signature⁸, our validation cohort included both HDGC and non-HDGC cases of diffuse and non-diffuse histology (Supplementary methods). Targeted sequencing identified four additional mutation carriers: two sharing a known Hispanic *BRCA1* founder mutation (p.Gln1111Asnfs)⁹ and two with novel *PALB2* mutations (p.Pro918Gln and p.Lys628_Cys630del) with predicted deleterious effects. Residue Pro918 falls in the PALB2 WD40 domain, which mediates interactions with BRCA2, RAD51 and RAD51C, whereas Lys628-Cys630 resides in the binding domain of MRG15, a transcription regulator and whose PALB2 interaction is required for homology directed DNA double-strand break repair indicating potential pathogenicity of these two novel mutations^{10, 11}.

In a third phase of the study, we genotyped all six *PALB2*, *RAD51C* and *BRCA1* mutations described above plus four known Hispanic *BRCA1/2* founder mutations (Supplementary methods) in 160 independent Latin American non-HDGC cases and found three additional mutation carriers, one with a *BRCA1* mutation (p.Gly559Valfs) and two with *PALB2* mutations (p.Lys628_Cys630del and p.Arg414Ter, Table 1). Interestingly, during the preparation of this manuscript, our clinic-based Portuguese collaborator (MT), identified one additional GC case (GM037589) with *PALB2* p.Arg414Ter. None of the seven *PALB2*, *RAD51C* and *BRCA1* mutations, detected in 11 unrelated Caucasian and Latin American cases, was detected in 1,170 population-matched controls (see mutation details in Supplementary Table 1).

Clinical details of our mutation carriers are given in Table 1. Most of them had diffuse histology, two had HDGC syndrome (CG-05 and GM022584) and one reported history of hereditary breast and ovarian cancer (HBOC, case CG-36, not shown). These mutation carriers were predominantly non-smokers and/or negative for *Helicobacter pylori* infection (Table 1), which suggest that GC risk in most of these cases was not driven by these two known environmental risk factors¹².

To obtain additional evidence of the causality of our HR gene mutations, we carried out loss of heterozygosity (LOH), mutational signature and co-segregation analyses in available samples from tumors and relatives. For LOH and mutational signatures, we performed WES in four available tumor samples from three *PALB2* (CG-12/p.Arg414Ter, CG-028/p.Lys628_Cys630del and 3CG-103/p.Pro918Gln) and *RAD51C* mutation carriers (Table 1). We found no LOH or compound heterozygosity in these tumor samples (not shown). Interestingly, when we analyzed the somatic WES data for mutational signatures, we found that all four tumors were enriched for a signature indicative of HR defects^{13, 14}, providing evidence for the causality of these mutations (Supplementary methods, Supplementary Figures 1 and 2).

Figure 1 shows available pedigrees from mutation carriers. Case 3CG-103 and her daughter were both diagnosed with GC and carried the *PALB2* p.Pro918Gln mutation (Figure 1A). GM037589, a *PALB2* p.Arg414Ter carrier, developed GC and BC and had a sister diagnosed with ovarian and endometrial cancer who also carried *PALB2* p.Arg414Ter (Figure 1B). The *RAD51C* p.Arg237Ter carrier's son died of colon cancer but did not carry the mutation (Figure 1C). We found that GC was the predominantly diagnosed malignancy among unavailable relatives of these carriers (Figures 1A–1D). Although we did not have access to samples from relatives of the *PALB2* p.Lys628_Cys630del carriers, our local collaborators found this mutation co-segregating in an unrelated breast cancer family (unpublished). Albeit limited, our co-segregation data partially support GC causality of *PALB2* mutations. The *RAD51C* co-segregation data is however inconclusive but the presence of a strong HR signature in the gastric tumor (see above) of this mutation carrier warrants further studies on *RAD51C* as a candidate GC gene.

In summary, our study identified eleven cases with mutations in *PALB2*, *BRCA1* and *RAD51C*, three closely-related HR genes. Some of these mutations are known to be pathogenic in other cancer types. Out of 362 cases analyzed, 6.45% of the HDGC cases (2 out of 31) and 2.7% (9 out of 331) of non-HDGC cases had *PALB2*, *BRCA1* or *RAD51C* mutations, suggesting that HR genes play a role in GC risk. Our data also provide evidence of a germline basis for the recently reported HR mutational signature in gastric tumors and strengthens the evidence for a causal role of these genes, specifically *PALB2*, in GC, as previously observed^{4, 15}. Future larger studies are needed to definitively assign causality and understand the penetrance and prevalence of HR gene mutations in GC and to further understand if and why some individuals from HBOC families with HR gene mutations develop GC. Further characterizations of the GC histology in HR gene mutation carriers are also needed as we found instances where the same mutation was found in cases with different histologies (CG-12 and CG-008 with *PALB2* p.Arg414ter and CG-039 and CG-028 with *PALB2* p.Lys628_Cys630del, Table 1). *CDHI* mutation negative families might benefit from HR gene testing and increased endoscopic surveillance and targeted therapies, such as PARP inhibitors⁸.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Grant support – No Stomach For Cancer Inc and the University of California, Davis provided the principal funding for this study. AC received funding from Conicyt-Fondap (15130011), CR-P received funding from Investigacion Sanitaria/FEDER (14/00230, 14/00173) and from the European Commission H2020-PHC (GA n°635290). ME and LGC-C received funding from Fondo de Investigaciones, Universidad del Tolima (Projects 30113, 160120516, 450110, 160114) and from COLCIENCIAS (Project 470115). JT was supported by CONACYT-Fronteras de la Ciencia (clave 773), and Instituto Mexicano del Seguro Social (FIS/IMSS/PROT/PRI0/13/027). This work was also supported in part by the Intramural Research Program of the National Cancer Institute, the National Institutes of Health, the Division of Cancer Epidemiology and Genetics. LGC-C received funding from the V Foundation from Cancer Research and from the National Cancer Institute (K12CA138464 and R21CA199631) of the National Institutes of Health. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

We thank all patients and their families for participating in the study and to multiple clinicians and research nurses who helped with patient recruitment. We are thankful to No Stomach For Cancer Inc. and the University of California for providing principal financial support to the study. We are grateful to former Carvajal-Carmona laboratory members John Williamson, Julian Halmi, Anna Marie Tuazon and Cathy Wang for their support during the study. We are also grateful to Drs. Yuichi Shiraishi and Matthew Stephens for their support and stimulating discussions about mutational signature analyses.

Members of the Latin American Gastric Cancer Genetics Collaborative Group: Magdalena Echeverry¹, Mabel Bohorquez¹, Rodrigo Prieto¹, John Suarez¹, Gilbert Mateus², Maria Mercedes Bravo³, Fernando Bolaños⁴, Alejandro Vélez⁵, Alejandro Corvalán⁶, Pilar Carvallo⁷, Javier Torres⁸, Luis Carvajal-Carmona⁹.

¹ Universidad del Tolima, Ibagué, Colombia; ² Hospital Federico Lleras Acosta, Ibagué, Colombia; ³ Instituto Nacional de Cancerología, Bogotá, D.C, Colombia; ⁴ Hospital Hernando Moncaleano Perdomo, Neiva, Colombia; ⁵ Hospital Pablo Tobón Uribe, Medellín, Colombia; ⁶ Facultad de Medicina, Pontificia Universidad Católica de Chile, Santiago, Chile; ⁷ Departamento de Biología Celular y Molecular, Facultad de Ciencias Biológicas, Pontificia Universidad Católica de Chile, Santiago, Chile; ⁸ Unidad de Investigación en Enfermedades Infecciosas, UMAE Pediatría, IMSS, México City, México; ⁹ University of California, Davis.

Abbreviations

GC	gastric cancer
HDGC	hereditary diffuse gastric cancer
WES	whole exome sequencing
LOH	loss of heterozygosity
BC	breast cancer
R	homologous recombination

References

Author names in bold designate shared co-first authorship

1. Ferlay J. *Int J Cancer*. 2015; 136:E359–86. [PubMed: 25220842]
2. La Vecchia C. *Cancer*. 1992; 70:50–5. [PubMed: 1606546]
3. Guilford P. *Nature*. 1998; 392:402–5. [PubMed: 9537325]
4. Hansford S. *JAMA Oncol*. 2015; 1:23–32. [PubMed: 26182300]
5. Prakash R. *Cold Spring Harb Perspect Biol*. 2015; 7:a016600. [PubMed: 25833843]
6. Antoniou AC. *N Engl J Med*. 2014; 371:1651–2. [PubMed: 25337756]
7. Landrum MJ. *Nucleic Acids Res*. 2016; 44:D862–8. [PubMed: 26582918]
8. Alexandrov LB. *Nat Commun*. 2015; 6:8683. [PubMed: 26511885]
9. Torres D. *Breast Cancer Res Treat*. 2007; 103:225–32. [PubMed: 17080309]

10. Park JY. *Oncogene*. 2014; 33:4803–12. [PubMed: 24141787]
11. Sy SM. *J Biol Chem*. 2009; 284:21127–31. [PubMed: 19553677]
12. Cover TL. *Gut Microbes*. 2013; 4:482–93. [PubMed: 23989802]
13. Alexandrov LB. *Nature*. 2013; 500:415–21. [PubMed: 23945592]
14. Shiraishi Y. *PLoS Genet*. 2015; 11:e1005657. [PubMed: 26630308]
15. Lu C, Xie M, Wendl MC, et al. *Nat Commun*. 2015; 6:10086. [PubMed: 26689913]

Table 1

Details of clinical information of the mutation carriers

Mutation details	ID	Age of onset	Sex	Histology	Satisfied HDGC criteria ?	<i>Helicobacter Pylori</i> infection	History of smoking
<i>PALB2</i> c.1240C>T, p.Arg414Ter	CG-12 ^{a*}	69	M	Intestinal	No	NA	NA
	CG-008 ^c	48	F	Diffuse	NA	NA	Yes
	GM037589	46	F	NA	No	Negative	No
<i>PALB2</i> c.3201+1G>T	CG-05 ^a	50	M	Diffuse	Yes	Negative	No
	CG-039 ^b	47	F	Diffuse	NA	Negative	No
	CG-028 ^{c*}	81	M	Intestinal	NA	Negative	Yes
<i>PALB2</i> c.2753C>A, p.Pro918Gln	3CG-103 ^{b*}	79	F	Mixed	No	Negative	Yes
	CG-036 ^b	67	F	Diffuse	No	NA	No
	CG-059 ^b	54	M	Diffuse	No	NA	No
<i>BRCA1</i> c.3331_3334delCAAG, p.Gln1111Asnfs	CG-001 ^c	65	M	NA	No	Positive	Yes
	GM022584 ^{a*}	73	M	Diffuse	Yes	Negative	No

Identified by:

^aWES,^btargeted sequencing or^cgenotyping.^{*} LOH and mutational signature analyzed.

NA: Not available