RESEARCH

Development and validation of the Bowel Cleansing Impact Review (BOCLIR)

Lynda Doward,¹ Jeanette Wilburn,¹ Stephen P McKenna,¹
Roger Leicester,² Owen Epstein,³ Vicki Hedley,² Sanjeeva Korala,³
James Twiss,¹ Deborah Jones,⁴ Mike Geraint⁴

¹Galen Research Ltd,
Manchester, UK

²Department of Endoscopy,
St George's Hospital, London,
UK

³Department of
Gastroenterology, The Royal Free
Hospital, London, UK

⁴Norgine Pharmaceuticals Ltd,
Uxbridge, UK

Correspondence to

Jeanette Wilburn, Galen
Research Ltd, Manchester
Science Park, Lloyd Street North,
Manchester M15 6SE, UK;
jwilburn@galen-research.com,
http://www.galen-research.com

Received 30 October 2012

Revised 10 December 2012

Accepted 12 December 2012

Published Online First

15 January 2013

ABSTRACT

Objective Acceptability and tolerability of bowel cleansers influence whether patients are able to complete the prescribed dose and, consequently, the quality of the cleansing achieved. No standardised means of assessing patients' experience of using bowel cleansing is currently available. The aim of the study was to develop the Bowel Cleansing Impact Review (BOCLIR) to assess patient response to bowel cleansing products.

Design Content was derived from qualitative interviews. Face and content validity were assessed via cognitive-debriefing interviews. Finally, patients completed the BOCLIR and a demographic questionnaire. Item response theory (Rasch analysis) was employed for item reduction and assessment of unidimensionality. Internal consistency and construct validity were also assessed.

Results Analysis of 40 interviews resulted in the production of three scales; patient satisfaction, symptomatic impact and activity limitations. Scales were designed to be used alone or together. 19 debriefing interviews demonstrated BOCLIR acceptability, relevance and ease of completion. The validation survey involved 166 patients (52% male, mean (SD) age 54.3 (15.2) years). After misfitting and redundant items were removed all scales fit the Rasch model confirming their unidimensionality. Cronbach's α -coefficients were high (0.77–0.94) indicating good internal consistency. Scores on the BOCLIR were related to patients' willingness to use the product in future and ease of drinking the full preparation ($p < 0.01$ for each scale).

Conclusions The BOCLIR is a new measure consisting of three unidimensional scales (satisfaction, symptoms and activity limitations) with good psychometric and scaling properties. The BOCLIR will allow accurate assessment of patients' response to bowel cleansing preparations.

INTRODUCTION

Procedures such as colonoscopy may be performed for the diagnosis, evaluation or therapeutic treatment of certain diseases of the gastrointestinal system. Adequate cleansing of the colon is a critical factor in achieving diagnostic accuracy and maximising the safety and efficacy of endoscopic or surgical resection.¹ A number of options are available for colon cleansing. The most widely employed are the orthograde gut lavage preparations using sodium or balanced electrolyte solutions. While patients' experiences of using bowel cleansing preparations have not been adequately researched, evidence suggests that this is dependent on the type of preparation used.^{2–3} The literature also suggests that the acceptability and tolerability of the specific product determines whether the patient is able to complete the prescribed dose and, hence, whether the bowel will be adequately cleansed. Thus, in addition to other factors such as dedicated time and information provided by the specialist screening practitioner, compliance has a direct impact on the accuracy and safety of the procedure. Serious consequences may follow from inadequate cleansing. These include missed underlying pathology, delayed treatment, complications and the cost of the failed procedures.

Studies assessing the relative effectiveness and tolerability of gut cleansing regimes have been hampered by the lack of a standardised measure. This is despite it becoming commonplace in many diagnostic areas to assess patient's views. Patient-Reported Outcome (PRO) is an

To cite: Doward L, Wilburn J, McKenna SP, et al. *Frontline Gastroenterology* 2013;**4**:112–119.

umbrella term that covers a wide range of potential measurement endpoints but is used specifically to refer to outcomes reported directly by the patient, without interpretation by clinicians or others.^{4,5} PRO data are collected with standardised instruments or questions designed to measure an explicit concept (construct). Constructs of interest may include impairments (symptoms), activity limitations (measures of physical, social or psychological functioning), quality of life, patient satisfaction, compliance or treatment preferences.⁶

A PRO measure is only capable of providing useful information if it meets certain developmental, psychometric and scaling standards.⁷ Specifically PROs should be:

- ▶ Based on an explicit description of the construct assessed.
- ▶ Relevant for, and well-targeted to, the patient group with which they are to be used. For a measure to be truly patient-based its content should be generated from relevant patients.⁶
- ▶ Reliable, valid and responsive.
- ▶ Unidimensional, allowing scale items to be summed to provide a valid total score.

The aim of the current study was to develop and validate the Bowel Cleansing Impact Review (BOCLIR). The acceptability, comprehensiveness and psychometric properties of the new instrument are reported below.

METHODOLOGY

The development of the BOCLIR involved three key stages:

1. Item generation and production of draft scales.
2. Assessment of face and content validity via cognitive debriefing interviews.
3. Validation by means of a survey designed to assess the psychometric and scaling properties of the scales.

The methodology adopted recent advances in the conceptual and practical bases of measurement; combining strong theoretical underpinnings with the statistical and diagnostic power of the Rasch model.⁸ The application of Rasch analysis ensures that the fundamental scaling properties of an instrument (eg, unidimensionality and level of measurement) are assessed in addition to the traditional psychometric assessments of reliability and external construct validity.

Patient samples

Approval for the study was granted by London-Surrey Borders Research Ethics Committee (LREC 06/Q0806/110). Patients for all stages of the study were recruited through two colonoscopy centres in London; St George's Hospital Tooting and the Royal Free Hospital in Hampstead. All patients attending for colonoscopy were invited to participate in the study. Although different patients were involved at each stage of the study the reasons for the

colonoscopies were comparable. Informed written consent was obtained from all participants.

Patients were eligible for the study if they were aged 18 or above, were able to provide written informed consent and could understand and complete questionnaires independently (as judged by the clinical team). Patients with major morbidity were excluded from the study.

Conceptual basis of the BOCLIR scales

The Symptom and Activity Limitations scales took as their conceptual basis the WHO classification of impairments (physiological and anatomical) and activity limitations (capacity and performance).^{9,10} WHO defines impairments as loss or abnormality of psychological, physiological or anatomical structure or function which represents disturbances at the level of the organ. Activity limitation (functioning) is defined as any restriction or lack of ability to perform an activity in the manner or within the range considered normal for a human being.

Patient satisfaction is a more difficult concept to define as there is no current consensus on what 'satisfaction' means or how best to measure it. However, this construct has become increasingly popular as an endpoint as patient satisfaction may influence a variety of other outcomes, including compliance, treatment success and continued use of health services.^{11–13}

Stages in the development of the BOCLIR scales

Stage 1: Item generation and production of draft scales

Qualitative interviews were conducted with patients in a private room following bowel cleansing and prior to their colonoscopy. The aim of the interviews was twofold. First, to identify the key impacts of the bowel cleansing process. Second, to provide statements that could form the basis of questionnaire items. Interviews took the form of an informal, focused conversation and participants were encouraged to talk freely about all aspects of their experience. Interviews were audio-recorded and verbatim transcripts produced. Thematic and interpretive phenomenological analyses¹⁴ were applied to the interview transcripts to identify key constructs for measurement and potential questionnaire items. Each interview transcript was analysed independently by two of five experienced researchers. Item pools were scrutinised to remove items that were idiosyncratic, ambiguous, badly phrased or duplicated.

Stage 2: Assessment of face and content validity

Semi-structured cognitive debriefing interviews were conducted with patients to assess the applicability, clarity, relevance, comprehensiveness and practicality of the draft measure. Respondents completed the BOCLIR at clinic, again after bowel cleansing and prior to their colonoscopy, in the presence of an

interviewer. Items found to be problematic were removed and a second draft version of the BOCLIR was prepared.

Stage 3: Validation survey

The purpose of the survey was to reduce the number of items in the BOCLIR scales and to assess the scaling and psychometric properties of the final scales. Participants completed the BOCLIR and a demographic questionnaire.

Scaling and item reduction analysis

Rasch analysis (one-parameter logistic item response theory) was conducted to determine the unidimensionality of the BOCLIR scales using the RUMM 2020 programme.¹⁵ Rasch analysis¹⁶ is a key tool in the development and improvement of questionnaires with advantages over traditional test theory and factor analysis.^{17–18} The Rasch model tests the assumption that a questionnaire scale has the basic property of unidimensionality and establishes whether all items capture a single underlying construct and can be combined to derive a valid total score.

Fit to the model is represented by a non-significant χ^2 fit statistic. A significance level of 0.01 was used to account for multiple testing. Items were assessed to see if they fit the Rasch model. Those that showed overall misfit to the model or had excessive fit residuals (> 2.5 or < -2.5) were considered for removal. Highly negative residuals indicate overfit to the model (suggesting that the item is duplicating other items in the scale) while highly positive residuals indicate that the item violates unidimensionality. The person separation index (PSI) or the extent to which items distinguish between distinct levels of the underlying construct was also assessed. For PSI, 0.7 is considered the minimal required value.¹⁹ Finally, differential item functioning (DIF), which represents the stability of the instrument irrespective of the group being evaluated, was assessed.^{20–21} Assessment of DIF ensures that responses to individual items are not affected by factors that are external to the measurement tool, such as age and gender. Such DIF was assessed using analysis of variance models. Items failing to fit the Rasch model and/or that showed DIF by gender or age (above or below the median) were removed. Throughout the item reduction process the content and meaning of items was taken into account before their removal, to maintain content validity.

Traditional psychometric analysis

Internal consistency was assessed using Cronbach's α . A correlation of 0.70 is accepted as indicating adequate consistency.²² Construct validity was determined by examining BOCLIR scores of respondents who differed according to certain known factors. For all three scales, respondent scores were related to: the amount of cleansing product the patient was able to consume ('full' or 'part'), ease of drinking the whole

volume ('very easy' to 'very difficult') and how willing they would be to use the same preparation in the future ('very happy' to 'very unhappy'). In addition, scores on the symptoms and activity limitations scales were related to how well the patient felt after taking the product, ('very well' to 'very unwell'), satisfaction with the overall experience ('very pleasant' to 'very unpleasant') and the convenience of the preparation ('very convenient' to 'very inconvenient'). Individuals reporting a worse experience and those who had greatest difficulty consuming the product were expected to have significantly higher BOCLIR scores. Non-parametric tests for independent samples (Mann-Whitney U test for two groups and Kruskal-Wallis one-way analysis of variance for three or more groups) were employed. Psychometric testing was performed using the SPSS V.16.0 statistical package.

RESULTS

Participant details for all study stages of the study are shown in table 1.

Item generation and production of draft scales

Interviews were conducted with 20 patients from each centre and lasted between 20 and 45 min. Three key areas of impact were identified; satisfaction with preparation, symptomatic impact and impact on the patient's ability to carry out their usual daily activities.

Common themes reported within these key areas are presented in table 2. Using these areas as a guide, interview transcripts were searched to identify statements that could be used as the basis for items. This allowed a draft questionnaire to be produced consisting of three scales; patient satisfaction (12 items), symptoms (29 items) and activity limitations (13 items). In order to ensure that individual symptoms were expressed in the most appropriate way for patients, some alternative expressions for the same concept were included in the draft questionnaire. It was intended that the less appropriate items would be removed from the scale during the later testing stages. Polytomous response formats were selected for the scales; five options for the patient satisfaction items and four for the symptoms and activity limitations items.

Assessment of face and content validity

Cognitive debriefing interviews were conducted with 19 patients. Interviews lasted approximately 20 min. Patients reported that they found the BOCLIR clear, acceptable and relevant. Minor changes were made to item ordering to improve the flow of the scale. A new draft BOCLIR was produced for use in the validation survey.

Validation survey

The BOCLIR and a demographic questionnaire were completed by 166 patients following bowel cleansing and prior to colonoscopy.

Table 1 Participant details for each stage of study

	Qualitative interviews	Cognitive debriefing interviews	Psychometric and scaling survey
n	40	19	166
Demographic details			
Male; n (%)	23 (57.5)	12 (63.2)	86 (51.8)
Age; mean (SD)	52.8 (15.8)	60.2 (14.9)	54.3 (15.2)
Married; n (%)	25 (62.5)	12 (63.2)	96 (57.8)
Employed (full or part time); n (%)	24 (60.0)	8 (42.1)	92 (55.4)
Colonoscopy details			
First; n (%)	14 (35.0)	12 (63.2)	103 (62.0)
Repeat; n (%)	26 (65.0)	7 (36.8)	62 (37.3)
Missing; n (%)	–	–	1 (0.6)
Amount of preparation taken			
All; n (%)	–	–	142 (85.5)
Most; n (%)	–	–	14 (8.4)
About 50%; n (%)	–	–	4 (2.4)
Missing; n (%)	–	–	6 (3.7)
Additional clear fluids			
Yes; n (%)	–	–	155 (93.4)
No; n (%)	–	–	5 (3.0)
Missing; n (%)	–	–	6 (3.6)
General health			
Very good; n (%)	13 (32.5)	2 (10.5)	–
Good; n (%)	14 (35.0)	7 (36.8)	–
Fair; n (%)	9 (22.5)	7 (36.8)	–
Poor; n (%)	4 (10.0)	3 (15.8)	–
Other health problems			
Yes; n (%)	21 (52.5)	9 (47.4)	–
No; n (%)	19 (47.5)	10 (52.6)	–

Scaling properties and item reduction

The initial 12-item satisfaction and 29-item symptom scales showed misfit to the Rasch model. Four items were removed from the satisfaction scale due to item misfit, extreme fit residuals or item redundancy. After

item reduction the scale fit the Rasch model (table 3). Low levels of uniform DIF by age were observed for three items and DIF by gender for one item. However, these factors were judged unlikely to compromise the scale's overall fit. Fifteen items were

Table 2 Key areas of impact

Patient satisfaction	Symptoms	Activity limitations
General satisfaction	Bloating	General ability to do usual activities
Volume	Cramps/gurgling/stomach pains	Household chores for example, cooking
Taste	Nausea/dizziness	Family impact
Drinking full volume/keeping it down	Energy/fatigue/concentration problems	Social impact
Preparation/instructions	Sleep disturbance	Work impact
Adhering to diet/fast	Headache	
Certainty of thoroughness of cleansing	Need to urinate	
Speed of effect	Bowel evacuation	
Comparison to other preparations	► Urgency/control	
	► Frequency	
	► Leakage	
	► Pain/soreness	
	Hunger	
	Dry mouth/thirst	
	Irritability	

Table 3 Rasch fit statistics for final Bowel Cleansing Impact Review scales

Scale	Item-trait interaction χ^2	PSI	Item-person interaction			
			Items		Persons	
			Mean	SD	Mean	SD
Satisfaction	p<0.13	0.86	0.22	1.12	-10.29	1.09
Symptoms	p<0.38	0.71	-0.07	0.96	-0.17	0.87
Activity limitations	p<0.43	0.94	0.05	1.4	-0.28	1.30

PSI, person separation index.

removed from the symptom scale due to item redundancy, misfit, extreme fit residuals, DIF, poor item total correlations or low endorsement rate. The final scale fit the Rasch model. After item reduction only one item exhibited low levels of DIF by age. The draft 13-item activity limitations scale fit the Rasch model.

Table 4 Bowel Cleansing Impact Review scores

	Satisfaction	Symptoms	Activity limitations
Mean (SD)	10.7 (5.9)	11.9 (5.7)	17.0 (10.0)
Median (IQR)	10 (6–15)	11.5 (8–15)	17 (9–25)
Range	0–24	2–28	0–36
% Scoring minimum	2.6	0.0	4.6
% Scoring maximum	0.0	0.0	0.0

However, one item displayed a high fit residual and was removed from the scale. Low levels of DIF by age were observed for three items and by gender for two items. Again, this was judged not to compromise overall scale fit. The PSI was good for all scales indicating appropriate inter-relatedness of items. The Rasch analyses demonstrated that the final versions of all scales were unidimensional ($p<0.05$) indicating that they each measure a single construct (table 3). Consequently, scale responses can be summed to provide valid scale scores.

Traditional psychometric analysis

The final satisfaction scale contains eight items scored 0 (highly satisfied) to 32 (highly dissatisfied). The symptoms scale includes 14 items scored 0 (no symptoms) to 42 (severe symptoms). The activity limitations scale is made up of 12 items with possible scores ranging from 0 (no effect on activities) to 36 (activities greatly affected). Table 4 shows descriptive statistics for the final scales. Limited ceiling or floor effects were found for all scales. Internal consistency was good for all three scales. Cronbach's α values were Satisfaction 0.84, Symptoms 0.77 and Activities 0.94.

Table 5 shows symptoms and activity limitation scale scores by satisfaction factors. Both scales were able to distinguish between groups of patients categorised by how well or unwell they felt after taking the bowel cleansing preparation, how pleasant or unpleasant they rated the experience of using the product and self-rated convenience of product use.

Figures 1–3 show BOCLIR scale scores by product consumption factors; specifically, volume of cleanser

Table 5 Known group validity for Bowel Cleansing Impact Review symptoms and activity limitations scales

	n	Symptoms Mean (%)	n	Activity limitations Mean (%)
How well or unwell did you feel as a result of taking the bowel preparation?				
Very well	41	7.9 (3.3)	41	12.0 (9.0)
Quite well	48	11.1 (4.8)	48	15.8 (9.6)
Neither well nor unwell	43	13.1 (5.5)	42	19.6 (9.5)
Quite / very unwell	21	18.3 (4.4)	19	24.8 (8.5)
p		< 0.001		< 0.001
Overall, how would you rate your experience of using this bowel preparation?				
Very pleasant	14	6.7 (4.1)	13	7.4 (9.1)
Quite pleasant	35	9.3 (4.1)	34	14.3 (9.2)
Neither pleasant nor unpleasant	62	11.2 (4.4)	62	15.5 (8.5)
Quite / very unpleasant	42	16.5 (5.6)	41	24.5 (8.6)
p		< 0.001		< 0.001
How convenient was it to fit taking the bowel preparation into your daily routine?				
Very convenient	33	8.8 (3.6)	34	12.8 (9.3)
Quite convenient	54	10.5 (4.8)	53	15.3 (10.6)
Neither convenient nor inconvenient	27	13.0 (6.0)	25	17.9 (9.4)
Quite inconvenient	28	15.4 (6.7)	27	20.9 (8.3)
Very inconvenient	10	13.9 (3.7)	11	26.2 (5.8)
p		< 0.001		< 0.001

Figure 1 Bowel Cleansing Impact Review (BOCLIR) scores by volume of bowel cleanser consumed.

consumed, ease of use and willingness to use in the future. Patients who reported a worse experience in terms of these factors, scored higher on the BOCLIR scales.

DISCUSSION

Adequate colon cleansing is a prerequisite to effective endoscopic examination and surgical procedures associated with the large colon and distal portion of the small bowel.²³ The accuracy with which colorectal polyps or lesions can be detected is largely associated with the cleanliness of the luminal environment.^{24–25} Poor cleansing is also associated with an increased risk of peritoneal contamination should perforation occur.^{1–2} Similarly, poor cleansing may increase the risk of colonic gas explosion during colonoscopy with electrocautery.²⁶ Indeed, up to a third of failed procedures have been attributed to inadequate bowel cleansing.¹

Bowel cleansing may be achieved by several different methods that vary by volume of liquid required for ingestion, the length of time over which the preparation is consumed and taste. Preparations are taken at prescribed intervals either all on the day before the colonic procedure or on a combination of the day before and the morning of the procedure. In addition, all preparations require the patient to maintain a liquid diet for at least 12 h prior to the procedure. It

has been reported that the requirement to use a bowel preparation may be a major factor in deterring patients from attending for screening colonoscopies.²³ Reports on adverse events and on ability to complete prescribed doses suggest that many patients find them unpleasant to use or even distressing. Symptoms such as nausea, abdominal cramps and vomiting are commonly reported.²⁷ Problems with sleep disturbance have also been reported.²⁸

For clinicians and commissioners to judge the best cleansing agent it is necessary to employ Patient-Reported Outcome Measures with good psychometric properties. The BOCLIR was developed to address this need. It consists of three stand-alone scales (satisfaction, symptoms and activity limitations) which together form a suite of instruments. The content of the BOCLIR was generated from qualitative interviews with relevant patients and the scales represent their key areas of concern. Respondents find the instrument acceptable, comprehensive and relevant. The BOCLIR is completed by the patient independently and takes approximately 5 min. It can be completed at any stage following bowel cleansing and prior to the procedure. As it is quick and easy to complete it is suitable for use in clinical settings.

Application of the Rasch model showed each BOCLIR scale to be unidimensional and to have minimal DIF. Confirmation of unidimensionality

Figure 2 Bowel Cleansing Impact Review (BOCLIR) scores by ease of drinking full volume of bowel cleanser.

Figure 3 Bowel Cleansing Impact Review (BOCLIR) scores by willingness to use cleanser in the future.

means that each scale provides an index or valid single total score. Construct validity was demonstrated by evaluating the scales' ability to distinguish between groups of patients categorised by known factors. Participants who reported a worse experience (as indicated by several variables) had worse scores on the BOCLIR scales.

It is concluded that the BOCLIR will provide a valuable tool for measuring patients' response to bowel cleansing preparations prior to colonoscopy.

The study has limitations. As it was designed to develop and validate the new measure it did not allow the comparison of BOCLIR scores with clinical judgement or markers of the degree of bowel cleansing achieved. It would be useful for future studies to compare BOCLIR scores with clinical assessments of bowel cleansing. It is anticipated that further use of the BOCLIR will provide such data. The work was conducted entirely in London. However, there is no reason to suggest that individuals from other locations would respond differently to bowel cleansing.

What is already known on this topic

- ▶ Satisfactory colonic cleansing is considered a crucial factor for the efficacy and safety of colonic procedures.
- ▶ Acceptability and tolerability of bowel cleansers influence the ability of the patient to complete the prescribed dose and, consequently, the quality of the cleansing achieved.
- ▶ The available literature suggests that there are clear differences in patient tolerability of different bowel cleansing preparations.
- ▶ No standardised means of assessing patients' experience of bowel cleansers is available.

What this study adds

- ▶ Three key factors affecting patient response to, and compliance with, bowel cleansing preparations were identified; satisfaction with the preparation, symptomatic impact and effect on patients' ability to carry out their usual daily activities.
- ▶ A new suite of Patient-reported outcome scales, the Bowel Cleansing Impact Review (BOCLIR), has been developed covering these key areas.
- ▶ The BOCLIR scales have been shown to be unidimensional and to have excellent content and construct validity.

How might it impact on clinical practice in the foreseeable future

- ▶ BOCLIR will allow accurate assessment of patients' response to bowel cleansing preparations.
- ▶ The measure will help clinicians, commissioners and the pharmaceutical industry to judge the benefits of one cleansing agent over another from the patients' perspective.

Acknowledgements The authors would like to acknowledge all those patients who helped with the study.

Contributors JW was involved with the design of the study, acquisition, analysis and interpretation of data and drafting of the manuscript. LD and SPM were involved in the conception and design of the study, acquisition and interpretation of data and contributed to the manuscript. RL and OE were involved with the design of the study, acquisition of data and reviewed and contributed to the manuscript. VH and SK were involved in the acquisition of data and reviewed and contributed to the manuscript. DJ and MG were involved in the design of the study and contributed to the manuscript. All authors read and approved the final manuscript.

Funding Norgine Pharmaceuticals Ltd.

Competing interests None.

Ethics approval London-Surrey Borders Research Ethics Committee.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Toledo TK, DiPalma JA. Review article: colon cleansing preparation for gastrointestinal procedures. *Aliment Pharmacol Ther* 2001;15:605–11.
- 2 Berry MA, DiPalma JA. Review article: orthograde gut lavage for colonoscopy. *Aliment Pharmacol Ther* 1994;8:391–5.
- 3 Belsey J, Epstein O, Heresbac D. Systematic review: oral bowel preparation for colonoscopy. *Aliment Pharmacol Ther* 2007;25:373–84.

- 4 Acquadro C, Berzon R, Dubois D, *et al.* Incorporating the patient's perspective into drug development and communication: an Ad Hoc Task Force report on the Patient-Reported Outcomes (PRO) Harmonization Group Meeting at the Food and Drug Administration, February 16, 2001. *Value Health* 2003;5:522–31.
- 5 Burke LB, Kennedy DL, Miskala PH, *et al.* The use of patient-reported outcome measures in the evaluation of medical products for regulatory approval. *Clin Pharmacol Ther* 2008;84:281–3.
- 6 McKenna SP. Measuring patient-reported outcomes: moving beyond misplaced common sense to hard science. *BMC Med* 2011;9:86. Review.
- 7 Berzon R, Hays RD, Shumaker SA. International use, application and performance of health-related quality of life instruments. *Qual Life Res* 1993;2:367–8.
- 8 Rasch G. *Probabilistic models for some intelligence and attainment tests*. Chicago: University of Chicago Press, 1960 (Reprinted 1980).
- 9 World Health Organisation. *The international classification of impairments, disabilities and handicaps*. Geneva: WHO, 1980.
- 10 World Health Organisation. *International classification of functioning and disability: ICDH-2*. Geneva: WHO, 2001.
- 11 Préau M, Protopopescu C, Raffi F, *et al.* Satisfaction with care in HIV-infected patients treated with long-term follow-up antiretroviral therapy: the role of social vulnerability. *AIDS Care* 2012;24:434–43.
- 12 Schoenthaler AM, Schwartz BS, Wood C, *et al.* Patient and physician factors associated with adherence to diabetes medications. *Diabetes Educ* 2012;38:397–408.
- 13 Kestemont P, Cartier H, Trevidic P, *et al.* Sustained efficacy and high patient satisfaction after cheek enhancement with a new hyaluronic acid dermal filler. *J Drugs Dermatol* 2012;11 (1 Suppl):s9–16.
- 14 Smith JA, Osborne M. Interpretive phenomenological analysis. In: Smith JA, ed. *Qualitative research: a practical guide to research methods*. London: Sage, 2008:53–80.
- 15 Andrich D. *Rasch models for measurement*. London: Sage Publications, 1988.
- 16 Tennant A, McKenna SP, Hagell P. Application of Rasch analysis in the development and application of quality of life instruments. *Value Health* 2004;7(Suppl 1):S22–6.
- 17 Wright BD. Comparing Rasch measurement and factor analysis. *Struct Equation Model* 1996;3:3–24.
- 18 Svensson E. Guidelines to statistical evaluation of data from rating scales and questionnaires. *J Rehabil Med* 2001; 33:47–8.
- 19 Streiner D, Norman G. *Health measurement scales*. Oxford: Oxford University Press, 1989.
- 20 Hagquist C, Andrich D. Is the sense of coherence-instrument applicable on adolescents? A latent trait analysis using Rasch-modelling. *Pers Indiv Dif* 2004;36:955–8.
- 21 Angoff WH. Perspectives on differential item functioning methodology. In: Holland PW, Wainer H, eds. *Differential item functioning*. Hillsdale, New Jersey: Lawrence Erlbaum, 1993.
- 22 Nunally JC Jr. *Psychometric theory*. 2nd edn. New York: McGraw-Hill, 1978:3–23.
- 23 Brown AR, DiPalma JA. Bowel preparation for gastrointestinal procedures. *Curr Gastroenterol Rep* 2004;6:395–401.
- 24 Kaminski MF, Regula J. Colorectal cancer screening by colonoscopy—current issues. *Digestion* 2007;76:20–5.
- 25 Dykes C, Cash BD. Key safety issues of bowel preparations for colonoscopy and importance of adequate hydration. *Gastroenterol Nurs* 2008;31:30–5;quiz 36–7.
- 26 Ladas SD, Karamanolis G, Ben-Soussan E. Colonic gas explosion during therapeutic colonoscopy with electrocautery. *World J Gastroenterol* 2007;13:5295–8.
- 27 Cohen SM, Wexner SD, Binderow SR, *et al.* Prospective, randomised, endoscopic-blinded trial comparing precolonoscopy bowel cleansing methods. *Dis Colon Rectum* 1994;37:689–96.
- 28 Fasih T, Tabaqchali MA, Varma JS. Patients' perception of colonoscopy. The Royal College of Surgeons of Edinburgh. Surgical Knowledge and Skills Website. Lecture Notes Series. <http://www.edu.rcsed.ac.uk/lectures/lt35.htm> (accessed 2006).