Published in final edited form as:

Am J Reprod Immunol. 2017 March ; 77(3): . doi:10.1111/aji.12617.

Neutrophil extracellular traps in acute chorioamnionitis: a mechanism of host defense

Nardhy Gomez-Lopez, MSc, PhD^{1,2,3}, Roberto Romero, MD DMedSci^{1,4,5,6}, Yaozhu Leng, MSc^{1,2}, Valeria Garcia-Flores, MSc^{1,2}, Yi Xu, PhD^{1,2}, Derek Miller, BSc^{1,2,3}, and Sonia S. Hassan, MD^{1,2}

¹Perinatology Research Branch, Program for Perinatal Research and Obstetrics, Division of Intramural Research, Eunice Kennedy Shriver National Institute of Child Health and Human Development, NICHD/NIH/DHHS, Bethesda, Maryland, and Detroit, Michigan, USA

²Department of Obstetrics and Gynecology, Wayne State University School of Medicine, Detroit, Michigan, USA

³Department of Immunology and Microbiology, Wayne State University School of Medicine, Detroit, Michigan, USA

⁴Department of Obstetrics and Gynecology, University of Michigan, Ann Arbor, Michigan, USA

⁵Department of Epidemiology and Biostatistics, Michigan State University, East Lansing, Michigan, USA

⁶Center for Molecular Medicine and Genetics, Wayne State University, Detroit, Michigan, USA

Abstract

Problem—Neutrophil extracellular traps (NETs) were recently described as a mechanism for microbial killing in the amniotic cavity of women with intra-amniotic infection. Such a clinical condition can result in acute chorioamnionitis, a placental lesion characterized by the infiltration of maternal neutrophils in the chorioamniotic membranes. Herein, we investigated whether these infiltrating neutrophils form NETs in the chorioamniotic membranes from women who underwent spontaneous term or preterm labor with acute chorioamnionitis.

Method of Study—Chorioamniotic membrane samples were collected from women who underwent spontaneous term or preterm labor with acute chorioamnionitis (n=10 each). Controls included chorioamniotic membrane samples from women who delivered at term or preterm with or without labor, in the absence of acute chorioamnionitis (n=10 each). NETs were visualized and semi-quantified in the chorioamniotic membranes by using antibodies against neutrophil elastase and histone H3, in combination with DAPI staining.

Address correspondence to: Nardhy Gomez-Lopez, PhD, Department of Obstetrics and Gynecology, Wayne State University School of Medicine, Perinatology Research Branch, NICHD/NIH/DHHS, Detroit, Michigan 48201, USA, Tel (313) 577-8904, ngomezlo@med.wayne.edu, Roberto Romero, MD, D. Med. Sci, Perinatology Research Branch, NICHD/NIH/DHHS, Wayne State University/Hutzel Women's Hospital, 3990 John R, Box 4, Detroit, MI 48201, USA, Telephone: (313) 993-2700, Fax: (313) 993-2694, prbchiefstaff@med.wayne.edu.

Disclosure/Conflict of Interest

The authors disclose no conflicts of interest.

Results—NETs were abundant in the chorioamniotic membranes from women who underwent spontaneous term or preterm labor with acute chorioamnionitis. NETs were rarely found, or not visualized at all, in the chorioamniotic membranes from women who delivered at term or preterm with or without labor, in the absence of acute chorioamnionitis.

Conclusion—NETs are abundant in the chorioamniotic membranes from women who underwent spontaneous term or preterm labor with acute chorioamnionitis. These findings suggest that chorioamniotic neutrophils can form NETs as a mechanism of host defense against infection or danger signals.

Keywords

Amniotic fluid; alarmins; cytokines; DAMPs; DNA; elastase; fetal inflammatory response; fever; funisitis; histone; intra-amniotic inflammation; microbial invasion of the amniotic cavity; parturition; pregnancy; prematurity; preterm birth; sterile intra-amniotic inflammation

INTRODUCTION

Acute chorioamnionitis is strongly associated with spontaneous preterm labor^{1–4}; yet, it is also frequently observed in the placentas of women who delivered after spontaneous labor at term^{5, 6}. In both spontaneous preterm and term labor, this placental lesion is associated with elevated concentrations of pro-inflammatory cytokines such as IL-1 α , IL-1 β , TNF- α , IL-8, and IL-6 in the amniotic fluid^{7–26} and umbilical cord blood^{27–33}. Elevated concentrations of these cytokines are linked to adverse neonatal outcomes^{20, 21, 27, 34–45}. Therefore, the study herein focused on the mechanisms implicated in acute chorioamnionitis.

The defining morphologic feature of acute chorioamnionitis is diffuse infiltration of neutrophils into the chorioamniotic membranes^{46, 47}. Neutrophils are rarely seen in the chorioamniotic membranes of patients without acute chorioamnionitis⁴⁷; therefore, we refer to these innate immune cells as chorioamniotic neutrophils. Their maternal origin was observed when two X chromosomes were detected by fluorescence *in situ* hybridization in the chorioamniotic leukocytes of women who delivered male preterm neonates and whose placenta was diagnosed with acute chorioamnionitis^{48, 49}. The current hypothesis⁴⁷ states that such maternal neutrophils migrate from the decidual vessels towards the chorion and amnion following a chemotactic gradient established by amniotic fluid chemokines such as IL-8^{12, 13, 19, 21, 22, 50–58}, CXCL6⁵⁹ and GRO α ^{52, 60}. Since acute chorioamnionitis generally represents the presence of intra-amniotic infection^{47, 61, 62}, we propose that chorioamniotic neutrophils play a role in the maternal host response against microbes invading the amniotic cavity.

In line with our hypothesis, we recently demonstrated that amniotic fluid neutrophils form neutrophil extracellular traps (NETs) as a mechanism for microbial killing in cases with intra-amniotic infection⁶³. NETs were initially described as web-like structures that contain DNA, histones, and anti-microbial products such as neutrophil elastase⁶⁴. NET formation is a specialized cell death process, which represents the final containment effort of a neutrophil to lyse pathogens⁶⁵. Although NET formation (or NETosis⁶⁶) was initially described as an *in vitro* phenomenon⁶⁴, *in vivo* NETosis can occur in tissues⁶⁴ and intravascular^{67/}

extravascular fluids (e.g. amniotic fluid⁶³). *In vitro*-induced NETs release their components freely as those traps formed in intravascular/extravascular fluids^{68, 69}. However, tissue NETs display a unique appearance in each tissue since the release of their components is restricted by the surrounding cellular structures^{68, 69}. Tissue NETs are generated in response to a local infection whereas intravascular NETs are formed in response to a systemic infection (i.e. sepsis)⁷⁰. Since acute chorioamnionitis represents the presence of a local inflammatory response in the amniotic cavity, we investigated whether infiltrating neutrophils form NETs in the chorioamniotic membranes.

MATERIALS AND METHODS

Human subjects, clinical specimens, and definitions

Chorioamniotic membrane samples were obtained from the Bank of Biological Specimens of the Detroit Medical Center, Wayne State University, and the Perinatology Research Branch (Detroit, MI, USA), an intramural program of the *Eunice Kennedy Shriver* National Institute of Child Health and Human Development, National Institutes of Health, U. S. Department of Health and Human Services (NICHD/NIH/DHHS). The collection and utilization of biological materials for research purposes were approved by the Institutional Review Boards of Wayne State University and NICHD. All participating women provided written informed consent. The following six study groups were included: 1) women who delivered at term without labor (TNL, n=10); 2) women who underwent spontaneous labor at term without acute chorioamnionitis (TIL, n=10); 3) women who underwent spontaneous labor at term with acute chorioamnionitis (TIL-ACA, n=10); 4) women who delivered preterm without labor (PTNL, n=10); 5) women who underwent spontaneous preterm labor without acute chorioamnionitis (PTL, n=10); and 6) women who underwent spontaneous preterm labor with acute chorioamnionitis (PTL-ACA, n=10). Table I includes the demographic and clinical characteristics of the study population. Multiparous women and women with neonates having congenital or chromosomal abnormalities were excluded. Labor at term was defined by the presence of regular uterine contractions at a frequency of at least two contractions every 10 minutes with cervical changes resulting in delivery. Preterm labor was diagnosed by the presence of regular uterine contractions (at least 3 in 30 min) and documented cervical changes in patients with a gestational age between 20 and 36 6/7 weeks. Preterm delivery was defined as birth prior to the 37th week of gestation.

Placental histopathological examinations

Five- μ m-thick sections of formalin-fixed, paraffin-embedded tissue specimens were cut and mounted on SuperFrostTM Plus microscope slides (Erie Scientific LLC, Portsmouth, NH, USA). In each case, several tissue sections of the chorioamniotic membranes, umbilical cord, and placental disc were examined. After deparaffinization, slides were rehydrated, stained with hematoxylin-eosin, and evaluated by pathologists who had been blinded to the clinical outcome, according to published criteria^{46, 47, 71}. Acute chorioamnionitis was diagnosed when the infiltration of neutrophils was observed in the chorionic trophoblast layer or chorioamniotic connective tissue^{46, 47, 71}.

Identification of neutrophil extracellular traps in the chorioamniotic membranes

Chorioamniotic membrane samples were frozen in Tissue-Plus O.C.T. compound (Fisher HealthCare, Houston, TX, USA) immediately after collection. Cryogenic sections were cut to 8 μ m and placed on glass microscope slides (Fisherbrand Superfrost Plus slides; Thermo Scientific, Waltham, MA, USA). The sections were fixed using 4% paraformaldehyde (Electron Microscopy Sciences, Hatfield, PA, USA) for 20 min at room temperature and rinsed with 1 $\times$ phosphate-buffered saline (PBS, Life Technologies, Grand Island, NY, USA). Prior to staining, non-specific antibody interactions were blocked using serum-free Protein Blocker (Cat#X09090; DAKO North America, Carpinteria, CA, USA) for 30 min at room temperature. The slides were then incubated at 4°C overnight with a mouse anti-human neutrophil elastase (Cat# M0752, clone NP57, DAKO, Glostrup, Denmark) and a rabbit anti-histone H3 antibody (Cat# ab5103, Abcam, Cambridge, MA, USA). Mouse IgG and rabbit IgG were used as negative controls, respectively. Following staining, slides were washed with 1 $\times$ PBS with 0.1% Tween 20. Next, a second blocking step was performed by adding 10% goat serum (KPL, Gaithersburg, MD, USA) for 10 mins at room temperature. The slides were then incubated with a secondary goat anti-mouse IgG–Alexa Fluor 488 antibody (Cat# A11029, Life Technologies) and a goat anti-rabbit IgG–Alexa Fluor 594 antibody (Cat#A11072, Life Technologies) for 30mins at room temperature in the dark. Finally, slides were washed with 1 $\times$ PBS and mounted with ProLong Diamond Antifade Mountant with DAPI (Thermo Fisher Scientific, Eugene, OR, USA). Slides were visualized on a Zeiss LSM 780 laser scanning confocal microscope (Carl Zeiss Microscopy GmbH, Jena, Germany) at the Microscopy, Imaging, and Cytometry Resources Core at the Wayne State University School of Medicine (<http://micr.med.wayne.edu/>). Tile-scans were performed from the chorioamniotic membranes and the complete imaging field was divided into eight-by-eight quadrants. Z-stack scans (8 μ m deep) were performed in order to create 3D reconstructions.

Semi-quantification of neutrophil extracellular traps in the chorioamniotic membranes

Following immunostaining, tissue slides were scanned using a Panoramic MIDI Digital Slide Scanner (PerkinElmer, Inc., Waltham, MA, USA). The chorioamniotic membrane section was divided into quadrants using the scanner software (3DHISTECH Ltd., Budapest, Hungary), and NET semi-quantification was performed in two opposing quadrants. Within each quadrant, five 1mm-wide sections of the choriodecidua and five 1-mm-wide sections of the amnion were chosen in pairs. The width of each section remained constant while the height spanned the full thickness of the amnion or choriodecidua. A NET was defined as a structure in which blue (DAPI), green (neutrophil elastase), and red (histone H3) fluorescence signals were co-localized. The total number of NETs was semi-quantified in the amnion and choriodecidua.

Statistical analyses

The SPSS v.19.0 software (SPSS Inc., Chicago, IL, USA) was used to analyze demographic, clinical and NET semi-quantification data. Normality of the data was tested using the Wilk-Shapiro test. Comparisons among groups were performed using the Kruskal–Wallis test followed by two group comparisons using the Mann-Whitney U-test. Comparison of

proportions was made using the Fisher's exact test. A p-value of <0.05 was used to determine statistical significance.

RESULTS

NETs were abundant in the chorioamniotic membranes from women who underwent spontaneous labor at term with acute chorioamnionitis (TIL-ACA; Figure 1A). However, NETs were rarely seen, or not found at all, in the chorioamniotic membranes from women who underwent spontaneous labor at term without acute chorioamnionitis or those who delivered at term without labor (TIL and TNL; Figure 1A). Magnifications of the NETs found in the chorioamniotic membranes from women who underwent spontaneous term labor with acute chorioamnionitis demonstrated that these traps contain neutrophil elastase and histone H3 (white arrows; Figure 1B), as well as DNA (white arrows; Figure 1C). Semi-quantification revealed that NETs were more abundant in the amnion (Figure 1D) and choriodecidua (Figure 1E) from women who underwent spontaneous labor at term with acute chorioamnionitis than in those without this placental lesion who delivered at term with or without labor. A 3D reconstruction shows that NETs are located in the amnion and choriodecidua from women who underwent spontaneous labor at term with acute chorioamnionitis (Video 1). A snapshot of this 3D reconstruction shows that chorioamniotic membrane NETs contain neutrophil elastase and histone H3 (white arrows; Figure 2A) as well as DNA (white arrows; Figure 2B).

NETs were also abundant in the chorioamniotic membranes from women who underwent spontaneous preterm labor with acute chorioamnionitis (PTL-ACA; Figure 3A). However, NETs were rarely seen, or not found at all, in the chorioamniotic membranes from women who underwent spontaneous preterm labor without acute chorioamnionitis or those who delivered preterm without labor (PTL and PTNL; Figure 3A). Magnifications of the NETs found in the chorioamniotic membranes from women who underwent spontaneous preterm labor with acute chorioamnionitis demonstrated that these traps contain neutrophil elastase and histone H3 (white arrows; Figure 3B), as well as DNA (white arrows; Figure 3C). Semi-quantification revealed that NETs were more abundant in the amnion (Figure 3D) and choriodecidua (Figure 3E) from women who underwent spontaneous preterm labor with acute chorioamnionitis than in those without this placental lesion who delivered preterm with or without labor. A 3D reconstruction shows that NETs are located in the amnion and choriodecidua from women who underwent spontaneous preterm labor with acute chorioamnionitis (Video 2). A snapshot of this 3D reconstruction shows that chorioamniotic membrane NETs contain neutrophil elastase and histone H3 (white arrows; Figure 4A) as well as DNA (white arrows; Figure 4B).

DISCUSSION

Acute chorioamnionitis generally represents the presence of intra-amniotic infection^{47, 61, 62}, a clinical condition that is characterized by a local inflammatory response containing abundant leukocytes^{72–76} and elevated concentrations of pro-inflammatory mediators such as cytokines^{25, 58}. Recently, we characterized the cellular composition of this local inflammatory response using immunophenotyping⁷⁷. We found that neutrophils are the most

abundant leukocyte subset in the amniotic cavity of women with intra-amniotic infection⁷⁷, which is consistent with previous observations⁷². Such neutrophils mainly express pro-inflammatory cytokines such as TNF- α , MIP-1 β , and IL-8⁷⁷. These cytokines are implicated in the processes of term and preterm parturition^{10–13, 25, 78–82}. In addition, amniotic fluid neutrophils form NETs in cases with intra-amniotic infection, which represents a new mechanism for trapping and/or killing microbes invading the amniotic cavity⁶³. Amniotic fluid neutrophils are considered to be of fetal origin^{83, 84}; however, these innate immune cells have also been observed in cases with a severe maternal inflammatory response (i.e., acute chorioamnionitis) but without a fetal inflammatory response (i.e. funisitis and chorionic vasculitis), suggesting that, in some cases, amniotic fluid neutrophils are of maternal origin or a mixture of both fetal and maternal neutrophils. In such cases, maternal neutrophils could be migrating from the decidual vessels into the chorion and amnion, causing acute inflammation of the chorioamniotic membranes (i.e. acute chorioamnionitis)⁴⁷ and ultimately reaching the amniotic cavity. Therefore, the function of chorioamniotic neutrophils in acute chorioamnionitis may be comparable to their role in intra-amniotic infection since, in both pathological processes, these innate immune cells form NETs and may participate in the maternal host response against microbes invading the amniotic cavity.

Yet, acute chorioamnionitis can also occur in the setting of sterile intra-amniotic inflammation^{61, 85–88}, an inflammatory process in which microorganisms cannot be detected using a combination of cultivation and molecular microbiology techniques^{61, 85–87}. Sterile inflammation is induced by danger signals termed damage-associated molecular patterns (DAMPs)⁸⁹ or alarmins⁹⁰, derived from necrotic cells or cellular stress⁹¹. NETs can also be formed in sterile inflammation since both alarmins and pathogen-associated molecular patterns (PAMPs) use the same sensor molecules or pattern recognition receptors⁹². Particularly, the high-mobility group box-1 (HMGB1, a prototypical alarmin^{93, 94}) protein can induce NET formation via TLR4⁹⁵, which is the sensor molecule for lipopolysaccharide from Gram negative bacteria⁹⁶. The fact that HMGB1 induces NETs is relevant since: (1) amniotic fluid HMGB1 concentrations are higher in women with intra-amniotic infection⁹⁷ or clinical chorioamnionitis⁹⁸ than in those without these clinical conditions; (2) patients with sterile intra-amniotic inflammation and high amniotic fluid HMGB1 concentrations delivered earlier than those with low concentrations of this alarmin⁸⁵; (3) the intra-amniotic administration of HMGB1 induces preterm labor and birth in mice⁹⁹; and (4) the chorioamniotic membranes from women who underwent spontaneous preterm labor releases high concentrations of HMGB1¹⁰⁰. Alarmin-induced NETs can exacerbate immune responses by directly causing tissue damage⁹². Together, these data suggest that in the setting of sterile intra-amniotic inflammation, chorioamniotic neutrophils form NETs in response to danger signals derived from the amniotic fluid or the chorioamniotic membranes which, in turn, could aggravate the local immune response observed in acute chorioamnionitis.

In summary, the study herein provides evidence that neutrophils infiltrating the chorioamniotic membranes in preterm and term cases of acute chorioamnionitis form NETs. These data suggest that chorioamniotic neutrophils form NETs in response to microbes invading the amniotic cavity (i.e. intra-amniotic infection) or danger signals derived from the amniotic fluid or chorioamniotic membranes (i.e. sterile intra-amniotic inflammation).

Collectively, these findings provide insight into the functions of infiltrating neutrophils in the chorioamniotic membranes from women who underwent spontaneous term or preterm labor with acute chorioamnionitis.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This research was supported, in part, by the Perinatology Research Branch, Division of Intramural Research, *Eunice Kennedy Shriver* National Institute of Child Health and Human Development, National Institutes of Health, U. S. Department of Health and Human Services (NICHD/NIH/DHHS), and, in part, with federal funds from the NICHD/NIH/DHHS under Contract No. HHSN275201300006C. This research was also supported by the Wayne State University Perinatal Initiative in Maternal, Perinatal and Child Health. We thank the physicians and nurses from the Center for Advanced Obstetrical Care and Research and the Intrapartum Unit for their help in collecting samples. We also thank staff members from the PRB Clinical Laboratory and the PRB Histology Unit for processing and preparing the pathological sections. Finally, we thank Tara Mial for her critical readings of the manuscript.

REFERENCES

1. Russell P. Inflammatory lesions of the human placenta: Clinical significance of acute chorioamnionitis. *Am J Diagn Gynecol Obstet.* 1979; 2:127–137.
2. Guzik DS, Winn K. The association of chorioamnionitis with preterm delivery. *Obstet Gynecol.* 1985; 65:11–16. [PubMed: 3966012]
3. van Hoesen KH, Anyaegbunam A, Hochster H, Whitty JE, Distant J, Crawford C, Factor SM. Clinical significance of increasing histologic severity of acute inflammation in the fetal membranes and umbilical cord. *Pediatr Pathol Lab Med.* 1996; 16:731–744. [PubMed: 9025872]
4. Lee SM, Park JW, Kim BJ, Park CW, Park JS, Jun JK, Yoon BH. Acute histologic chorioamnionitis is a risk factor for adverse neonatal outcome in late preterm birth after preterm premature rupture of membranes. *PLoS One.* 2013; 8:e79941. [PubMed: 24324586]
5. Seong HS, Lee SE, Kang JH, Romero R, Yoon BH. The frequency of microbial invasion of the amniotic cavity and histologic chorioamnionitis in women at term with intact membranes in the presence or absence of labor. *Am J Obstet Gynecol.* 2008; 199:375, e371–e375. [PubMed: 18928978]
6. Park HS, Romero R, Lee SM, Park CW, Jun JK, Yoon BH. Histologic chorioamnionitis is more common after spontaneous labor than after induced labor at term. *Placenta.* 2010; 31:792–795. [PubMed: 20655108]
7. Romero R, Brody DT, Oyarzun E, Mazor M, Wu YK, Hobbins JC, Durum SK. Infection and labor. III. Interleukin-1: a signal for the onset of parturition. *Am J Obstet Gynecol.* 1989; 160:1117–1123. [PubMed: 2786341]
8. Romero R, Parvizi ST, Oyarzun E, Mazor M, Wu YK, Avila C, Athanassiadis AP, Mitchell MD. Amniotic fluid interleukin-1 in spontaneous labor at term. *J Reprod Med.* 1990; 35:235–238. [PubMed: 2325034]
9. Romero R, Mazor M, Brandt F, Sepulveda W, Avila C, Cotton DB, Dinarello CA. Interleukin-1 alpha and interleukin-1 beta in preterm and term human parturition. *Am J Reprod Immunol.* 1992; 27:117–123. [PubMed: 1418402]
10. Romero R, Manogue KR, Mitchell MD, Wu YK, Oyarzun E, Hobbins JC, Cerami A. Infection and labor. IV. Cachectin-tumor necrosis factor in the amniotic fluid of women with intraamniotic infection and preterm labor. *Am J Obstet Gynecol.* 1989; 161:336–341. [PubMed: 2764054]
11. Romero R, Mazor M, Sepulveda W, Avila C, Copeland D, Williams J. Tumor necrosis factor in preterm and term labor. *Am J Obstet Gynecol.* 1992; 166:1576–1587. [PubMed: 1595815]

12. Romero R, Ceska M, Avila C, Mazor M, Behnke E, Lindley I. Neutrophil attractant/activating peptide-1/interleukin-8 in term and preterm parturition. *Am J Obstet Gynecol.* 1991; 165:813–820. [PubMed: 1951537]
13. Cherouny PH, Pankuch GA, Romero R, Botti JJ, Kuhn DC, Demers LM, Appelbaum PC. Neutrophil attractant/activating peptide-1/interleukin-8: association with histologic chorioamnionitis, preterm delivery, and bioactive amniotic fluid leukoattractants. *Am J Obstet Gynecol.* 1993; 169:1299–1303. [PubMed: 8238198]
14. Romero R, Avila C, Santhanam U, Sehgal PB. Amniotic fluid interleukin 6 in preterm labor. Association with infection. *J Clin Invest.* 1990; 85:1392–1400. [PubMed: 2332497]
15. Romero R, Yoon BH, Kenney JS, Gomez R, Allison AC, Sehgal PB. Amniotic fluid interleukin-6 determinations are of diagnostic and prognostic value in preterm labor. *Am J Reprod Immunol.* 1993; 30:167–183. [PubMed: 8311926]
16. Hillier SL, Witkin SS, Krohn MA, Watts DH, Kiviat NB, Eschenbach DA. The relationship of amniotic fluid cytokines and preterm delivery, amniotic fluid infection, histologic chorioamnionitis, and chorioamnion infection. *Obstet Gynecol.* 1993; 81:941–948. [PubMed: 8497360]
17. Saito S, Kasahara T, Kato Y, Ishihara Y, Ichijo M. Elevation of amniotic fluid interleukin 6 (IL-6), IL-8 and granulocyte colony stimulating factor (G-CSF) in term and preterm parturition. *Cytokine.* 1993; 5:81–88. [PubMed: 7683506]
18. Halgunset J, Johnsen H, Kjollesdal AM, Qvigstad E, Espevik T, Austgulen R. Cytokine levels in amniotic fluid and inflammatory changes in the placenta from normal deliveries at term. *Eur J Obstet Gynecol Reprod Biol.* 1994; 56:153–160. [PubMed: 7821485]
19. Arntzen KJ, Kjollesdal AM, Halgunset J, Vatten L, Austgulen R. TNF, IL-1, IL-6, IL-8 and soluble TNF receptors in relation to chorioamnionitis and premature labor. *J Perinat Med.* 1998; 26:17–26. [PubMed: 9595363]
20. Yoon BH, Jun JK, Romero R, Park KH, Gomez R, Choi JH, Kim IO. Amniotic fluid inflammatory cytokines (interleukin-6, interleukin-1beta, and tumor necrosis factor-alpha), neonatal brain white matter lesions, and cerebral palsy. *Am J Obstet Gynecol.* 1997; 177:19–26. [PubMed: 9240577]
21. Yoon BH, Romero R, Jun JK, Park KH, Park JD, Ghezzi F, Kim BI. Amniotic fluid cytokines (interleukin-6, tumor necrosis factor-alpha, interleukin-1 beta, and interleukin-8) and the risk for the development of bronchopulmonary dysplasia. *Am J Obstet Gynecol.* 1997; 177:825–830. [PubMed: 9369827]
22. Figueroa R, Garry D, Elimian A, Patel K, Sehgal PB, Tejani N. Evaluation of amniotic fluid cytokines in preterm labor and intact membranes. *J Matern Fetal Neonatal Med.* 2005; 18:241–247. [PubMed: 16318974]
23. Holst RM, Laurini R, Jacobsson B, Samuelsson E, Savman K, Doverhag C, Wennerholm UB, Hagberg H. Expression of cytokines and chemokines in cervical and amniotic fluid: relationship to histological chorioamnionitis. *J Matern Fetal Neonatal Med.* 2007; 20:885–893. [PubMed: 18050018]
24. Kacerovsky M, Celec P, Vlkova B, Skogstrand K, Hougaard DM, Cobo T, Jacobsson B. Amniotic fluid protein profiles of intraamniotic inflammatory response to *Ureaplasma* spp. and other bacteria. *PLoS One.* 2013; 8:e60399. [PubMed: 23555967]
25. Romero R, Grivel JC, Tarca AL, Chaemsaitong P, Xu Z, Fitzgerald W, Hassan SS, Chaiworapongsa T, Margolis L. Evidence of perturbations of the cytokine network in preterm labor. *Am J Obstet Gynecol.* 2015; 213:836.e1–836.e18. [PubMed: 26232508]
26. Yoneda S, Shiozaki A, Ito M, Yoneda N, Inada K, Yonezawa R, Kigawa M, Saito S. Accurate Prediction of the Stage of Histological Chorioamnionitis before Delivery by Amniotic Fluid IL-8 Level. *Am J Reprod Immunol.* 2015; 73:568–576. [PubMed: 25600281]
27. Yoon BH, Romero R, Yang SH, Jun JK, Kim IO, Choi JH, Syn HC. Interleukin-6 concentrations in umbilical cord plasma are elevated in neonates with white matter lesions associated with periventricular leukomalacia. *Am J Obstet Gynecol.* 1996; 174:1433–1440. [PubMed: 9065108]
28. Dollner H, Vatten L, Halgunset J, Rahimipoor S, Austgulen R. Histologic chorioamnionitis and umbilical serum levels of pro-inflammatory cytokines and cytokine inhibitors. *BJOG.* 2002; 109:534–539. [PubMed: 12066943]

29. Tasci Y, Dilbaz B, Uzmez Onal B, Caliskan E, Dilbaz S, Doganci L, Han U. The value of cord blood interleukin-6 levels for predicting chorioamnionitis, funisitis and neonatal infection in term premature rupture of membranes. *Eur J Obstet Gynecol Reprod Biol.* 2006; 128:34–39. [PubMed: 16459014]
30. Andrys C, Drahosova M, Hornychova H, Tambor V, Musilova I, Tosner J, Flidrova E, Kacerovsky M. Umbilical cord blood concentrations of IL-6, IL-8, and MMP-8 in pregnancy complicated by preterm premature rupture of the membranes and histological chorioamnionitis. *Neuro Endocrinol Lett.* 2010; 31:857–863. [PubMed: 21196908]
31. Duncombe G, Veldhuizen RA, Gratton RJ, Han VK, Richardson BS. IL-6 and TNFalpha across the umbilical circulation in term pregnancies: relationship with labour events. *Early Hum Dev.* 2010; 86:113–117. [PubMed: 20171025]
32. Roberts DJ, Celi AC, Riley LE, Onderdonk AB, Boyd TK, Johnson LC, Lieberman E. Acute histologic chorioamnionitis at term: nearly always noninfectious. *PLoS One.* 2012; 7:e31819. [PubMed: 22412842]
33. Chan CJ, Summers KL, Chan NG, Hardy DB, Richardson BS. Cytokines in umbilical cord blood and the impact of labor events in low-risk term pregnancies. *Early Hum Dev.* 2013; 89:1005–1010. [PubMed: 24045131]
34. Yoon BH, Romero R, Park JS, Kim CJ, Kim SH, Choi JH, Han TR. Fetal exposure to an intra-amniotic inflammation and the development of cerebral palsy at the age of three years. *Am J Obstet Gynecol.* 2000; 182:675–681. [PubMed: 10739529]
35. Hitti J, Tarczy-Hornoch P, Murphy J, Hillier SL, Aura J, Eschenbach DA. Amniotic fluid infection, cytokines, and adverse outcome among infants at 34 weeks' gestation or less. *Obstet Gynecol.* 2001; 98:1080–1088. [PubMed: 11755557]
36. Moon JB, Kim JC, Yoon BH, Romero R, Kim G, Oh SY, Kim M, Shim SS. Amniotic fluid matrix metalloproteinase-8 and the development of cerebral palsy. *J Perinat Med.* 2002; 30:301–306. [PubMed: 12235718]
37. Combs CA, Gravett M, Garite TJ, Hickok DE, Lapidus J, Porreco R, Rael J, Grove T, Morgan TK, Clewell W, Miller H, Luthy D, Pereira L, Nageotte M, Robilio PA, Fortunato S, Simhan H, Baxter JK, Amon E, Franco A, Trofatter K, Heyborne K. ProteoGenix/Obstetrix Collaborative Research N: Amniotic fluid infection, inflammation, and colonization in preterm labor with intact membranes. *Am J Obstet Gynecol.* 2014; 210:836.e1–836.e18.
38. Kunze M, Klar M, Morfeld CA, Thorns B, Schild RL, Markfeld-Erol F, Rasenack R, Proempeler H, Hentschel R, Schaefer WR. Cytokines in noninvasively obtained amniotic fluid as predictors of fetal inflammatory response syndrome. *Am J Obstet Gynecol.* 2016; 215:96.e1–96.e8. [PubMed: 26829512]
39. Gomez R, Romero R, Ghezzi F, Yoon BH, Mazor M, Berry SM. The fetal inflammatory response syndrome. *Am J Obstet Gynecol.* 1998; 179:194–202. [PubMed: 9704787]
40. Berner R, Niemeyer CM, Leititis JU, Funke A, Schwab C, Rau U, Richter K, Tawfeek MS, Clad A, Brandis M. Plasma levels and gene expression of granulocyte colony-stimulating factor, tumor necrosis factor-alpha, interleukin (IL)-1beta, IL-6, IL-8, and soluble intercellular adhesion molecule-1 in neonatal early onset sepsis. *Pediatr Res.* 1998; 44:469–477. [PubMed: 9773833]
41. Dollner H, Vatten L, Linnebo I, Zanussi GF, Laerdal A, Austgulen R. Inflammatory mediators in umbilical plasma from neonates who develop early-onset sepsis. *Biol Neonate.* 2001; 80:41–47. [PubMed: 11474148]
42. Goepfert AR, Andrews WW, Carlo W, Ramsey PS, Cliver SP, Goldenberg RL, Hauth JC. Umbilical cord plasma interleukin-6 concentrations in preterm infants and risk of neonatal morbidity. *Am J Obstet Gynecol.* 2004; 191:1375–1381. [PubMed: 15507968]
43. An H, Nishimaki S, Ohyama M, Haruki A, Naruto T, Kobayashi N, Sugai T, Kobayashi Y, Mori M, Seki K, Yokota S. Interleukin-6, interleukin-8, and soluble tumor necrosis factor receptor-I in the cord blood as predictors of chronic lung disease in premature infants. *Am J Obstet Gynecol.* 2004; 191:1649–1654. [PubMed: 15547537]
44. Armstrong-Wells J, Donnelly M, Post MD, Manco-Johnson MJ, Winn VD, Sebire G. Inflammatory predictors of neurologic disability after preterm premature rupture of membranes. *Am J Obstet Gynecol.* 2015; 212:212 e211–212 e219. [PubMed: 25223243]

45. Cordeiro CN, Savva Y, Vaidya D, Argani CH, Hong X, Wang X, Burd I. Mathematical Modeling of the Biomarker Milieu to Characterize Preterm Birth and Predict Adverse Neonatal Outcomes. *Am J Reprod Immunol*. 2016; 75:594–601. [PubMed: 26892347]
46. Redline RW. Classification of placental lesions. *Am J Obstet Gynecol*. 2015; 213:S21–S28. [PubMed: 26428500]
47. Kim CJ, Romero R, Chaemsaitong P, Chaityasit N, Yoon BH, Kim YM. Acute chorioamnionitis and funisitis: definition, pathologic features, and clinical significance. *Am J Obstet Gynecol*. 2015; 213:S29–S52. [PubMed: 26428501]
48. McNamara MF, Wallis T, Qureshi F, Jacques SM, Gonik B. Determining the maternal and fetal cellular immunologic contributions in preterm deliveries with clinical or subclinical chorioamnionitis. *Infect Dis Obstet Gynecol*. 1997; 5:273–279. [PubMed: 18476151]
49. Steel JH, O'Donoghue K, Kennea NL, Sullivan MH, Edwards AD. Maternal origin of inflammatory leukocytes in preterm fetal membranes, shown by fluorescence in situ hybridisation. *Placenta*. 2005; 26:672–677. [PubMed: 16085046]
50. Gomez R, Ghezzi F, Romero R, Munoz H, Tolosa JE, Rojas I. Premature labor and intra-amniotic infection. Clinical aspects and role of the cytokines in diagnosis and pathophysiology. *Clin Perinatol*. 1995; 22:281–342. [PubMed: 7671540]
51. Ghezzi F, Gomez R, Romero R, Yoon BH, Edwin SS, David C, Janisse J, Mazor M. Elevated interleukin-8 concentrations in amniotic fluid of mothers whose neonates subsequently develop bronchopulmonary dysplasia. *Eur J Obstet Gynecol Reprod Biol*. 1998; 78:5–10. [PubMed: 9605441]
52. Hsu CD, Meaddough E, Aversa K, Copel JA. The role of amniotic fluid L-selectin, GRO-alpha, and interleukin-8 in the pathogenesis of intraamniotic infection. *Am J Obstet Gynecol*. 1998; 178:428–432. [PubMed: 9539502]
53. Hsu CD, Meaddough E, Aversa K, Hong SF, Lu LC, Jones DC, Copel JA. Elevated amniotic fluid levels of leukemia inhibitory factor, interleukin 6, and interleukin 8 in intra-amniotic infection. *Am J Obstet Gynecol*. 1998; 179:1267–1270. [PubMed: 9822513]
54. Jacobsson B, Mattsby-Baltzer I, Andersch B, Bokstrom H, Holst RM, Wennerholm UB, Hagberg H. Microbial invasion and cytokine response in amniotic fluid in a Swedish population of women in preterm labor. *Acta Obstet Gynecol Scand*. 2003; 82:120–128. [PubMed: 12648172]
55. Jacobsson B, Mattsby-Baltzer I, Andersch B, Bokstrom H, Holst RM, Nikolaitchouk N, Wennerholm UB, Hagberg H. Microbial invasion and cytokine response in amniotic fluid in a Swedish population of women with preterm prelabor rupture of membranes. *Acta Obstet Gynecol Scand*. 2003; 82:423–431. [PubMed: 12752072]
56. Witt A, Berger A, Gruber CJ, Petricevic L, Apfalter P, Husslein P. IL-8 concentrations in maternal serum, amniotic fluid and cord blood in relation to different pathogens within the amniotic cavity. *J Perinat Med*. 2005; 33:22–26. [PubMed: 15841609]
57. Cobo T, Kacerovsky M, Palacio M, Hornychova H, Hougaard DM, Skogstrand K, Jacobsson B. Intra-amniotic inflammatory response in subgroups of women with preterm prelabor rupture of the membranes. *PLoS One*. 2012; 7:e43677. [PubMed: 22916296]
58. Romero R, Chaemsaitong P, Korzeniewski SJ, Tarca AL, Bhatti G, Xu Z, Kusanovic JP, Dong Z, Docheva N, Martinez-Varea A, Yoon BH, Hassan SS, Chaiworapongsa T, Yeo L. Clinical chorioamnionitis at term II: the intra-amniotic inflammatory response. *J Perinat Med*. 2016; 44:5–22. [PubMed: 25938217]
59. Mittal P, Romero R, Kusanovic JP, Edwin SS, Gotsch F, Mazaki-Tovi S, Espinoza J, Erez O, Nhan-Chang CL, Than NG, Vaisbuch E, Hassan SS. CXCL6 (granulocyte chemotactic protein-2): a novel chemokine involved in the innate immune response of the amniotic cavity. *Am J Reprod Immunol*. 2008; 60:246–257. [PubMed: 18782286]
60. Cohen J, Ghezzi F, Romero R, Ghidini A, Mazor M, Tolosa JE, Goncalves LF, Gomez R. GRO alpha in the fetomaternal and amniotic fluid compartments during pregnancy and parturition. *Am J Reprod Immunol*. 1996; 35:23–29. [PubMed: 8789556]
61. Romero R, Miranda J, Chaiworapongsa T, Chaemsaitong P, Gotsch F, Dong Z, Ahmed AI, Yoon BH, Hassan SS, Kim CJ, Korzeniewski SJ, Yeo L. A novel molecular microbiologic technique for the rapid diagnosis of microbial invasion of the amniotic cavity and intra-amniotic infection in

- preterm labor with intact membranes. *Am J Reprod Immunol*. 2014; 71:330–358. [PubMed: 24417618]
62. Romero R, Chaemsaihong P, Docheva N, Korzeniewski SJ, Kusanovic JP, Yoon BH, Kim JS, Chaiyasit N, Ahmed AI, Qureshi F, Jacques SM, Kim CJ, Hassan SS, Chaiworapongsa T, Yeo L, Kim YM. Clinical chorioamnionitis at term VI: acute chorioamnionitis and funisitis according to the presence or absence of microorganisms and inflammation in the amniotic cavity. *J Perinat Med*. 2016; 44:33–51. [PubMed: 26352071]
 63. Gomez Lopez N, Romero R, Xu Y, Miller D, Unkel R, Shaman M, Jacques SM, Panaitescu B, Garcia-Flores V, Hassan SS. Neutrophil extracellular traps in the amniotic cavity of women with intra-amniotic infection: a new mechanism of host defense. *Reprod Sci*. Epub 2016 Nov. 23.
 64. Brinkmann V, Reichard U, Goosmann C, Fauler B, Uhlemann Y, Weiss DS, Weinrauch Y, Zychlinsky A. Neutrophil extracellular traps kill bacteria. *Science*. 2004; 303:1532–1535. [PubMed: 15001782]
 65. Fuchs TA, Abed U, Goosmann C, Hurwitz R, Schulze I, Wahn V, Weinrauch Y, Brinkmann V, Zychlinsky A. Novel cell death program leads to neutrophil extracellular traps. *J Cell Biol*. 2007; 176:231–241. [PubMed: 17210947]
 66. Steinberg BE, Grinstein S. Unconventional roles of the NADPH oxidase: signaling, ion homeostasis, and cell death. *Sci STKE*. 2007; 2007:pe11. [PubMed: 17392241]
 67. Tanaka K, Koike Y, Shimura T, Okigami M, Ide S, Toiyama Y, Okugawa Y, Inoue Y, Araki T, Uchida K, Mohri Y, Mizoguchi A, Kusunoki M. In vivo characterization of neutrophil extracellular traps in various organs of a murine sepsis model. *PLoS One*. 2014; 9:e111888. [PubMed: 25372699]
 68. Brinkmann V, Zychlinsky A. Beneficial suicide: why neutrophils die to make NETs. *Nat Rev Microbiol*. 2007; 5:577–582. [PubMed: 17632569]
 69. Yipp BG, Kubes P. NETosis: how vital is it? *Blood*. 2013; 122:2784–2794. [PubMed: 24009232]
 70. Sorensen OE, Borregaard N. Neutrophil extracellular traps - the dark side of neutrophils. *J Clin Invest*. 2016; 126:1612–1620. [PubMed: 27135878]
 71. Redline RW. Placental pathology: a systematic approach with clinical correlations. *Placenta*. 2008; 29(Suppl A):S86–S91. [PubMed: 17950457]
 72. Romero R, Quintero R, Nores J, Avila C, Mazor M, Hanaoka S, Hagay Z, Merchant L, Hobbins JC. Amniotic fluid white blood cell count: a rapid and simple test to diagnose microbial invasion of the amniotic cavity and predict preterm delivery. *Am J Obstet Gynecol*. 1991; 165:821–830. [PubMed: 1951538]
 73. Romero R, Yoon BH, Mazor M, Gomez R, Diamond MP, Kenney JS, Ramirez M, Fidel PL, Sorokin Y, Cotton D, et al. The diagnostic and prognostic value of amniotic fluid white blood cell count, glucose, interleukin-6, and gram stain in patients with preterm labor and intact membranes. *Am J Obstet Gynecol*. 1993; 169:805–816. [PubMed: 7694461]
 74. Romero R, Yoon BH, Mazor M, Gomez R, Gonzalez R, Diamond MP, Baumann P, Araneda H, Kenney JS, Cotton DB, et al. A comparative study of the diagnostic performance of amniotic fluid glucose, white blood cell count, interleukin-6, and gram stain in the detection of microbial invasion in patients with preterm premature rupture of membranes. *Am J Obstet Gynecol*. 1993; 169:839–851. [PubMed: 7694463]
 75. Gomez R, Romero R, Galasso M, Behnke E, Insunza A, Cotton DB. The value of amniotic fluid interleukin-6, white blood cell count, and gram stain in the diagnosis of microbial invasion of the amniotic cavity in patients at term. *Am J Reprod Immunol*. 1994; 32:200–210. [PubMed: 7533501]
 76. Yoon BH, Yang SH, Jun JK, Park KH, Kim CJ, Romero R. Maternal blood C-reactive protein, white blood cell count, and temperature in preterm labor: a comparison with amniotic fluid white blood cell count. *Obstet Gynecol*. 1996; 87:231–237. [PubMed: 8559530]
 77. Martinez-Varea A, Romero R, Xu Y, Miller D, Ahmed AI, Chaemsaihong P, Chaiyasit N, Yeo L, Shaman M, Lannaman K, Cher B, Hassan SS, Gomez-Lopez N. Clinical chorioamnionitis at term VII: the amniotic fluid cellular immune response. *J Perinat Med*. Epub 2016 Oct 20.
 78. Osmer RG, Blaser J, Kuhn W, Tschesche H. Interleukin-8 synthesis and the onset of labor. *Obstet Gynecol*. 1995; 86:223–229. [PubMed: 7617353]

79. Elliott CL, Loudon JA, Brown N, Slater DM, Bennett PR, Sullivan MH. IL-1 β and IL-8 in human fetal membranes: changes with gestational age, labor, and culture conditions. *Am J Reprod Immunol.* 2001; 46:260–267. [PubMed: 11642674]
80. Young A, Thomson AJ, Ledingham M, Jordan F, Greer IA, Norman JE. Immunolocalization of proinflammatory cytokines in myometrium, cervix, and fetal membranes during human parturition at term. *Biol Reprod.* 2002; 66:445–449. [PubMed: 11804961]
81. Osman I, Young A, Ledingham MA, Thomson AJ, Jordan F, Greer IA, Norman JE. Leukocyte density and pro-inflammatory cytokine expression in human fetal membranes, decidua, cervix and myometrium before and during labour at term. *Mol Hum Reprod.* 2003; 9:41–45. [PubMed: 12529419]
82. Gomez-Lopez N, Tong WC, Arenas-Hernandez M, Tanaka S, Hajar O, Olson DM, Taggart MJ, Mitchell BF. Chemotactic activity of gestational tissues through late pregnancy, term labor, and RU486-induced preterm labor in Guinea pigs. *Am J Reprod Immunol.* 2015; 73:341–352. [PubMed: 25329235]
83. Sampson JE, Theve RP, Blatman RN, Shipp TD, Bianchi DW, Ward BE, Jack RM. Fetal origin of amniotic fluid polymorphonuclear leukocytes. *Am J Obstet Gynecol.* 1997; 176:77–81. [PubMed: 9024093]
84. Lee SD, Kim MR, Hwang PG, Shim SS, Yoon BH, Kim CJ. Chorionic plate vessels as an origin of amniotic fluid neutrophils. *Pathol Int.* 2004; 54:516–522. [PubMed: 15189506]
85. Romero R, Miranda J, Chaiworapongsa T, Korzeniewski SJ, Chaemsathong P, Gotsch F, Dong Z, Ahmed AI, Yoon BH, Hassan SS, Kim CJ, Yeo L. Prevalence and clinical significance of sterile intra-amniotic inflammation in patients with preterm labor and intact membranes. *Am J Reprod Immunol.* 2014; 72:458–474. [PubMed: 25078709]
86. Romero R, Miranda J, Chaiworapongsa T, Chaemsathong P, Gotsch F, Dong Z, Ahmed AI, Yoon BH, Hassan SS, Kim CJ, Korzeniewski SJ, Yeo L, Kim YM. Sterile intra-amniotic inflammation in asymptomatic patients with a sonographic short cervix: prevalence and clinical significance. *J Matern Fetal Neonatal Med.* Epub 2014 Sep 24.
87. Romero R, Miranda J, Chaemsathong P, Chaiworapongsa T, Kusanovic JP, Dong Z, Ahmed AI, Shaman M, Lannaman K, Yoon BH, Hassan SS, Kim CJ, Korzeniewski SJ, Yeo L, Kim YM. Sterile and microbial-associated intra-amniotic inflammation in preterm prelabor rupture of membranes. *J Matern Fetal Neonatal Med.* 2015; 28:1394–1409. [PubMed: 25190175]
88. Romero R, Miranda J, Kusanovic JP, Chaiworapongsa T, Chaemsathong P, Martinez A, Gotsch F, Dong Z, Ahmed AI, Shaman M, Lannaman K, Yoon BH, Hassan SS, Kim CJ, Korzeniewski SJ, Yeo L, Kim YM. Clinical chorioamnionitis at term I: microbiology of the amniotic cavity using cultivation and molecular techniques. *J Perinat Med.* 2015; 43:19–36. [PubMed: 25720095]
89. Lotze MT, Zeh HJ, Rubartelli A, Sparvero LJ, Amoscato AA, Washburn NR, Devera ME, Liang X, Tor M, Billiar T. The grateful dead: damage-associated molecular pattern molecules and reduction/oxidation regulate immunity. *Immunol Rev.* 2007; 220:60–81. [PubMed: 17979840]
90. Oppenheim JJ, Yang D. Alarmins: chemotactic activators of immune responses. *Curr Opin Immunol.* 2005; 17:359–365. [PubMed: 15955682]
91. Chen GY, Nunez G. Sterile inflammation: sensing and reacting to damage. *Nat Rev Immunol.* 2010; 10:826–837. [PubMed: 21088683]
92. Gupta S, Kaplan MJ. The role of neutrophils and NETosis in autoimmune and renal diseases. *Nat Rev Nephrol.* 2016; 12:402–413. [PubMed: 27241241]
93. Wang H, Bloom O, Zhang M, Vishnubhakat JM, Ombrellino M, Che J, Frazier A, Yang H, Ivanova S, Borovikova L, Manogue KR, Faist E, Abraham E, Andersson J, Andersson U, Molina PE, Abumrad NN, Sama A, Tracey KJ. HMG-1 as a late mediator of endotoxin lethality in mice. *Science.* 1999; 285:248–251. [PubMed: 10398600]
94. Lotze MT, Tracey KJ. High-mobility group box 1 protein (HMGB1): nuclear weapon in the immune arsenal. *Nat Rev Immunol.* 2005; 5:331–342. [PubMed: 15803152]
95. Tadie JM, Bae HB, Jiang S, Park DW, Bell CP, Yang H, Pittet JF, Tracey K, Thannickal VJ, Abraham E, Zmijewski JW. HMGB1 promotes neutrophil extracellular trap formation through interactions with Toll-like receptor 4. *Am J Physiol Lung Cell Mol Physiol.* 2013; 304:L342–L349. [PubMed: 23316068]

96. Poltorak A, He X, Smirnova I, Liu MY, Van Huffel C, Du X, Birdwell D, Alejos E, Silva M, Galanos C, Freudenberg M, Ricciardi-Castagnoli P, Layton B, Beutler B. Defective LPS signaling in C3H/HeJ and C57BL/10ScCr mice: mutations in Tlr4 gene. *Science*. 1998; 282:2085–2088. [PubMed: 9851930]
97. Romero R, Chaiworapongsa T, Alpay Savasan Z, Xu Y, Hussein Y, Dong Z, Kusanovic JP, Kim CJ, Hassan SS. Damage-associated molecular patterns (DAMPs) in preterm labor with intact membranes and preterm PROM: a study of the alarmin HMGB1. *J Matern Fetal Neonatal Med*. 2011; 24:1444–1455. [PubMed: 21958433]
98. Romero R, Chaiworapongsa T, Savasan ZA, Hussein Y, Dong Z, Kusanovic JP, Kim CJ, Hassan SS. Clinical chorioamnionitis is characterized by changes in the expression of the alarmin HMGB1 and one of its receptors, sRAGE. *J Matern Fetal Neonatal Med*. 2012; 25:558–567. [PubMed: 22578261]
99. Gomez-Lopez N, Romero R, Plazyo O, Panaitescu B, Furcron AE, Miller D, Roumayah T, Flom E, Hassan SS. Intra-Amniotic Administration of HMGB1 Induces Spontaneous Preterm Labor and Birth. *Am J Reprod Immunol*. 2016; 75:3–7. [PubMed: 26781934]
100. Plazyo O, Romero R, Unkel R, Balancio A, Mial TN, Xu Y, Dong Z, Hassan SS, Gomez Lopez N. HMGB1 induces an inflammatory response in the chorioamniotic membranes that is partially mediated by the inflammasome. *Biol Reprod*. 2016 Epub 2016 Nov. 2.

Figure 1.

Neutrophil extracellular traps (NETs) in the chorioamniotic membranes from women who delivered at term. (A) A tile-scan image of the chorioamniotic membranes from women who delivered at term without labor or underwent spontaneous labor at term with or without acute chorioamnionitis. Merged images show DAPI (nuclei) in blue, neutrophil elastase in green, and histone H3 in red. Tile-scan images were acquired at 400X. The area outlined in (A) is enlarged in (B&C), demonstrating a higher resolution view of a NET in the choriodecuidua from women who underwent spontaneous labor at term with acute chorioamnionitis. Merged images show neutrophil elastase in green and histone H3 in red

(B; white arrow) or DAPI (nuclei) in blue, neutrophil elastase in green, and histone H3 in red (C; white arrow). NETs are structures in which DAPI, neutrophil elastase, and histone H3 fluorescence signals are co-localized. Semi-quantification of the total number of NETs in the amnion (D) and choriodecidua (E).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 2.

A snapshot of the 3D reconstruction of neutrophil extracellular traps (NETs) in the chorioamniotic membranes from women who underwent spontaneous labor at term with acute chorioamnionitis. A merged image shows neutrophil elastase in green and histone H3 in red (A; white arrows). A merged image shows DAPI (nuclei) in blue, neutrophil elastase in green and histone H3 in red (B; white arrows). 400 × magnification. NETs are structures in which DAPI, neutrophil elastase, and histone H3 fluorescence signals are co-localized.

Figure 3.

Neutrophil extracellular traps (NETs) in the chorioamniotic membranes from women who delivered preterm. (A) A tile-scan image of the chorioamniotic membranes from women who delivered preterm without labor or underwent spontaneous preterm labor with or without acute chorioamnionitis. Merged images show DAPI (nuclei) in blue, neutrophil elastase in green, and histone H3 in red. Tile-scan images were acquired at 400X. The area outlined in (A) is enlarged in (B, C), demonstrating a higher resolution view of a NET in the amnion from women who underwent spontaneous preterm labor with acute chorioamnionitis. Merged images show neutrophil elastase in green and histone H3 in red

(B; white arrow) or DAPI (nuclei) in blue, neutrophil elastase in green, and histone H3 in red (C; white arrow). NETs are structures in which DAPI, neutrophil elastase, and histone H3 fluorescence signals are co-localized. Semi-quantification of the total number of NETs in the amnion (D) and choriodecidua (E).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 4.

A snapshot of the 3D reconstruction of neutrophil extracellular traps (NETs) in the chorioamniotic membranes from women who underwent spontaneous preterm labor with acute chorioamnionitis. The merged image shows neutrophil elastase in green and histone H3 in red (A; white arrows). The merged image shows DAPI (nuclei) in blue, neutrophil elastase in green, and histone H3 in red (B; white arrows). 400 × magnification. NETs are structures in which DAPI, neutrophil elastase, and histone H3 fluorescence signals are co-localized.

Table 1

Demographic and clinical characteristics of the study population

	TNL (n=10)	TIL (n=10)	TIL-ACA (n=10)	PTNL (n=10)	PTL (n=10)	PTL-ACA (n=10)	p value
Age (y; median [IQR]) ^a	28 (25.3–29.8)	24.5 (21.5–31.3)	22.5 (20.5–26)	29 (22.5–33)	25.5 (22.5–28.3)	27 (21.5–30.8)	NS
Body mass index (kg/m ² ; median [IQR]) ^a	33.7 (29.8–36.6)	23 (21.1–27.2)	27.9 (24.6–29)	27.5 (21.9–32.8)	23.4 (22.7–27.3)	25.2 (21.4–27.1)	p=0.037
Gestational age at delivery (wk; median [IQR]) ^a	39.3 (39–39.6)	39.2 (38.9–39.9)	39.2 (38.8–40)	31 (29.7–32.6)	33.6 (32.8–34.1)	32.75 (31.6–34.5)	p<0.001
Birth weight (g; median [IQR]) ^a	3692.5 (3006.3–3853.8)	3145 (2677.5–3255.75)	3322.5 (3093.8–3476.3)	1170 (931.3–1290)	1895 (1717.5–2078.8)	1827.5 (1517.5–2848.8)	p<0.001
Race (n[%]) ^b							
African-American	6 (60%)	10 (100%)	9 (90%)	8 (80%)	10 (100%)	7 (70%)	NS
Caucasian	3 (30%)	0 (0%)	1 (10%)	1 (10%)	0 (0%)	1 (10%)	
Other	1 (10%)	0 (0%)	0 (0%)	1 (10%)	0 (0%)	2 (20%)	
Primiparity (n[%]) ^b	0 (0%)	3 (30%)	1 (10%)	1 (10%)	1 (10%)	1 (10%)	NS
C-section (n[%]) ^b	10 (100%)	2 (20%)	1 (10%)	10 (100%)	4 (40%)	1 (10%)	NS
Acute fetal inflammatory response (n[%]) ^b							
Stage 1 (acute phlebitis/chorionic vasculitis)	0 (0%)	0 (0%)	2 (20%)	0 (0%)	0 (0%)	4 (40%)	NS
Stage 2 (acute arteritis)	0 (0%)	0 (0%)	7 (70%)	0 (0%)	0 (0%)	3 (30%)	NS
Stage 3 (necrotizing funisitis)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	1 (10%)	NS

^aKruskal-Wallis test^bFisher's exact test

IQR, interquartile range; NS, non-significant