CORRESPONDENCE

Dependence on Prescription Opioids—Prevention, Diagnosis and Treatment

by Dr. Johannes Just, Dr. Martin Mücke, and Dr. Markus Bleckwenn in issue 13/2016

psychological and physical function among opioid-using chronic pain patients. *J Pain Symptom Manage* 2014; 48: 1091–9.

Dr. Dirk K. Wolter
 Psykiatrien i Region Syddanmark
 Gerontopsykiatrisk Afdeling
 Aabenraa, Danmark
 dirk.wolter@rsyd.dk

Abuse of Opioid Analgesics Is Missing

I am pleased to see the subject of opioid analgesic abuse being broached by primary care physicians (1). In the USA and Canada, abuse of prescription opioids led to an “opioid overdose epidemic”: for a while more deaths have been caused by opioid analgesics than by heroin and cocaine put together. Drug addicts in particular use oxycodone as a heroin substitute (2).

The diagnostic criteria of ICD-10 proceed from the assumption of illegal heroin use. They fail to take proper account of abuse of opioid analgesics. Pain treatment with opioid analgesics can lead to “complex persistent dependence” (3) that cannot be equated with heroin dependence. The exhaustive, 25-year-long discussion of alternative diagnostic criteria is not mentioned, and no source is given for the “typical behavior patterns” listed in *Box 3* in the article. The same is true for the “basic concepts of dependence” (*Box 2*), for some of which the consensus on definition suggested in the article does not in fact exist. The extent to which the North American discussion and the almost exclusively North American epidemiological findings can be extrapolated to the situation in Germany is not discussed.

Without exception, the screening instruments mentioned in the article were developed in North America and are aimed at dissocial behavior and (illegal) drug use; their suitability for application in the context of pain treatment or in the elderly (for whom in Germany—in contrast to North America—most opioid analgesics are prescribed) is questionable. There is an instrument, however, which is suitable as a guide for medical history and medical decision making, namely the Diagnosis, Intractability, Risk, Efficacy (DIRE) tool (German version available at www.dggpp.de/publik_mat.html).

Furthermore, “mindfulness-oriented recovery enhancement” (MORE) is a treatment approach with a positive effect both on pain and on dependence problems/opioid analgesic abuse (4).

DOI: 10.3238/arztebl.2016.0676a

REFERENCES

1. Just J, Mücke M, Bleckwenn M: Dependence on prescription opioids—prevention, diagnosis and treatment. *Dtsch Arztebl Int* 2016; 113: 213–20.
2. Wolter DK: Schmerzen und Schmerzmittelabhängigkeit im Alter – die gerontopsychiatrische Perspektive. Stuttgart: Kohlhammer 2016.
3. Ballantyne JC, Sullivan MD, Kolodny A: Opioid dependence vs addiction: a distinction without a difference? *Arch Intern Med* 2012; 172: 1342–3.
4. Garland EL, Thomas E, Howard MO: Mindfulness-oriented recovery enhancement ameliorates the impact of pain on self-reported

Early Cooperation With Psychiatrists

With regard to the contraindications to treatment with opioids, differential diagnostic investigation of the possibility that chronic pain is caused by a mental disorder requires early cooperation with psychiatrists. In addition, the diagnosis of dependence as the cause of pain could be swiftly confirmed by consulting an addiction specialist. For example, the conjunction of positive experience and loss of control mentioned in *Box 2* in the article is not so characteristic. On the one hand loss of control often occurs without pleasant sensations, and on the other hand repeated consumption because of pleasant experiences on previous consumption does not correspond to loss of control. It is more helpful simply to use the descriptive diagnostic criteria of ICD-10 for classification.

Even if the dosage is reduced in a structured fashion and the patient is switched to a long-acting opioid, the goal of opioid abstinence is often hard to achieve. If psychopharmaceuticals such as antidepressants or antipsychotics are prescribed in the attempt to reduce the opioid dose, their adverse effects and interactions have to be borne in mind in continued opioid treatment. “Structured” use supported by behavioral therapy is a good option.

In the case of substitution therapy, take-home prescriptions are an option (1). The patient can take the substitute medication at home, reducing the number of office visits to one per week.

In conclusion, we would like to stress that the authors' work (2) provides an important reminder of the benefit of closer cooperation between addiction and pain specialists—a project that we at the Bavarian Academy for Addiction and Health Issues (*Bayerische Akademie für Sucht- und Gesundheitsfragen*) are also pursuing (3).

DOI: 10.3238/arztebl.2016.0676b

REFERENCES

1. Bundesärztekammer: Richtlinie der Bundesärztekammer zur Durchführung der substitionsgestützten Behandlung Opiatabhängiger. www.bundesaerztekammer.de/fileadmin/user_upload/downloads/RL-Substitution_19-Februar-2010.pdf (last accessed on 20 June 2016).
2. Just J, Mücke M, Bleckwenn M: Dependence on prescription opioids—prevention, diagnosis and treatment. *Dtsch Arztebl Int* 2016; 113: 213–20.
3. BLÄK informiert: Schmerz(medizin) trifft Sucht(medizin): 15. Suchtforum in Bayern. www.bayerisches-aerzteblatt.de/fileadmin/aerzteblatt/

ausgaben/2016/05/einzel/pdf/BAB_5_2016_218_219.pdf (last accessed on 27 June 2016)

Dr. Felix Tretter

Bayerische Akademie für Suchtfragen, BAS e.V.

Dr. Beate Erbas, MPH

Bayerische Akademie für Sucht- und Gesundheitsfragen, BAS UG
erbas@bas-muenchen.de

Lower Saxony Is an Exception

“Psychosocial care” in Germany is tailored to the treatment requirements of heroin-dependent patients. Patients dependent on painkillers form a different group; most of them are occupationally and socially integrated, and they are not burdened by the numerous diseases associated with heroin use. Therefore, their needs are better served by a facility for the treatment of dependence on medicinal drugs than by the “psychosocial care”. Psychotherapy can be considered if the medication dependence rests on an underlying psychic illness.

As laid down in the German law regulating the prescription of narcotic drugs (*Betäubungsmittel-Verschreibungsverordnung*, BtMVV), a precondition for substitution therapy is “opiate dependence.” In Germany, this is traditionally understood to mean dependence on heroin. However, in most regions the Association of Statutory Health Insurance Physicians accepts dependence on fully synthetic opioids as a justified reason for substitution therapy. Not so in the federal state of Lower Saxony, where, for example, the statutory health insurance provider of a patient dependent on tramadol will not pay for substitution. It is to be hoped that this discrimination will be eliminated in the next reform of the BtMVV.

The literature contains no recommendations for the duration of outpatient withdrawal from opioids. The time needed depends on the duration of the foregoing consumption. In our experience at least 2 weeks should be allowed, plus 1 week for each year of opioid use, with considerable interindividual variation. In our practice we use buprenorphine to support gentle withdrawal. Switching to the opiate antagonist naltrexone after dose reduction has been demonstrated to help maintain abstinence. The naltrexone treatment should be continued for at least 6 months, preferably for a full year. Particularly in this phase of treatment an accompanying program in a facility for patients dependent on medicinal drugs is advisable.

D01: 10.3238/arztebl.2016.0677a

REFERENCES

1. Piest B: Ambulanter Entzug bei Opioidabhängigkeit und anschließende Abstinenztherapie. Suchttherapie 2015; 16: 42–8.
2. Just J, Mücke M, Bleckwenn M: Dependence on prescription opioids—prevention, diagnosis and treatment. Dtsch Arztebl Int 2016; 113: 213–20.

Dr. Bernhard Piest

Braunschweig
bernhardpiest@arcor.de

In Reply:

We are grateful for the important and informative comments on our article about dependence on prescription opioids (1). This is a complex problem that cannot be solved without active interdisciplinary cooperation.

As Dr. Wolter emphasizes—and as we stated in the concluding section of our article—the currently available epidemiological data stem overwhelmingly from North America. These have to be supplemented by German data so we can better estimate the scale of the problem in our healthcare system. The exclusively English-language screening instruments should be validated for the German-speaking countries and adapted as required.

As stressed by Dr. Tretter and Dr. Erbas, ICD-10 provides a good basis for recognition of addiction. The quality of care is, as they mention, probably highly dependent on efficient cooperation among various specialists. In our view these should include primary care physicians just as much as addiction and pain specialists or psychiatrists/psychotherapists.

There is no sound medical or scientific justification for the restricted indication for substitution in Lower Saxony described by Dr. Piest. In the interests of the patients affected, urgent change is necessary. We would like to add to Dr. Piest's remarks on “psychosocial care”. In our experience, some patients who are dependent on prescription opioids do in fact benefit from such care. We suggest one appointment. If the physician, the psychosocial service, and the patient then decide that further psychosocial care would not be beneficial, this decision should be documented. Moreover, additional therapeutic elements (psychotherapy, self-help groups, and so on) can be added on a case to case basis.

D01: 10.3238/arztebl.2016.0677b

REFERENCES

1. Just J, Mücke M, Bleckwenn M: Dependence on prescription opioids—prevention, diagnosis and treatment. Dtsch Arztebl Int 2016; 113: 213–20.

Dr. Johannes Just

Institut für Hausarztmedizin, Allgemeinmedizin, Bonn
johannesjust@gmail.com

Conflict of interest statement

The authors of all contributions state that no conflict of interests exists.