Published in final edited form as:

Eur J Clin Nutr. 2017 April ; 71(4): 486–493. doi:10.1038/ejcn.2016.179.

Sociodemographic Disparity in the Diet Quality Transition among Chinese adults, 1991 to 2011

Zhihong Wang^{1,3}, Penny Gordon-Larsen¹, Anna Maria Siega-riz¹, Jianwen Cai², Huijun Wang³, Linda S. Adair¹, and Barry M. Popkin^{1,*}

¹Department of Nutrition, Carolina Population Center, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, USA (ZW, AMSR, PG-L, LA and BP)

²Department of Biostatistics, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, USA (JC)

³National Institute for Nutrition and Health, Chinese Center for Disease Control and Prevention, Beijing, China (ZW,HW)

Abstract

Objective—This study investigates secular trends in diet quality distribution and related socioeconomic disparity from 1991 to 2011 in the Chinese adult population.

Methods—The analysis uses the 1991–2011 China Health and Nutrition Survey data on 13,853 participants (6,876 men; 6,977 women) ages 18 to 65 with 56,319 responses. Dietary assessment was carried out over a 3-day period with 24-hour recalls combined a household food inventory. We tailored Alternative Healthy Eating Index 2010 (named as tAHEI) to measure diet quality and performed quantile regression to investigate shifts in tAHEI scores at different percentiles and used mixed-effect linear regression to examine average diet quality trend and potential sociodemographic disparity.

Results—The energy-adjusted mean tAHEI scores increased from 36.9 (36.7–37.1) points in 1991 to 50.3 (50.1–50.5) in 2011 for men ($p < 0.001$) and from 35.6 (35.4–35.8) to 46.9 (46.7–47.1) for women ($p < 0.001$). The covariates-adjusted score of polyunsaturated fatty acids increased by 6.8 (6.6, 7.0) and 7.0 (6.9, 7.2) and the score of long chain (ω -3) fats increased by 5.3 (5.2, 5.4) and 5.3 (5.2, 5.5) in men and women, respectively, while the cereal fiber and red meat scores decreased slightly. Increasing tAHEI score occurred across the entire distribution and diet quality transition varied across sociodemographic groups.

*Corresponding author: Barry M. Popkin, W. R. Kenan Jr. Distinguished Professor, Carolina Population Center, department of nutrition, University of North Carolina at Chapel Hill, CB #8120 University Square, Chapel Hill, NC 27516-3997, Phone: 919-962-6139, Fax: 919-966-9159 (backup: 919-966-6638) popkin@unc.edu.

Contributors

ZW designed and conducted the analysis and drafted the manuscript. PG-L, AMSR, JC, LA, HW and BP contributed to the interpretation of the data analysis and critical revision of the manuscript for important intellectual content. ZW and BP had full access to all of the data in the study and had primary responsibility for the final content.

Disclosure Statement

The authors declare no conflict interest.

Supplementary information is available at European Journal of Clinical Nutrition's website

Conclusion—Chinese diet quality is far from optimal with moderate improvement over 21-year period. Findings suggest that nutritional intervention should give priority to low income, low urbanized communities, and southern provincial adults with low diet quality in China.

Keywords

diet quality; distribution; quantile regression; China; adults

INTRODUCTION

The World Health Organization reports that unhealthy diet, excessive alcohol consumption, smoking, and decreased physical activity (PA) are modifiable risk factors for the increasing chronic disease epidemic worldwide ¹. Over the past two decades, China has experienced marked shifts in diet ^{2, 3} along with its rapid economic growth and social changes and the concurrent shifts in disease patterns ³. Chinese food consumption has been characterized as rapid declines in intake of coarse grains, vegetables, and legumes and increases in intake of edible oils and animal-source foods²⁻⁴. One key issue is how to capture the dynamic complexity of multidimensional diet changes.

Diet quality indexes, based on current healthy diet recommendations, have gained increasing attention for assessing diet quality as a whole and allowing for standardization of the scores, reproducibility of results and thus, comparability of results across studies from different populations. The Harvard Healthy Eating Pyramid (HEP), a popular healthy diet guideline, is based on global scientific evidence on diet-disease relationships^{1, 5, 6}. The Alternative Healthy Eating Index (AHEI) ⁷ and its most recent version, AHEI-2010 ⁸ assess adherence to the HEP and have strongly predicted major chronic disease risk in many Western populations⁹⁻¹³. Two recent studies showed the negative association of the tailored AHEI-2010 (named as tAHEI) with risk of low density lipoprotein cholesterol in Chinese adults aged 18 to 65¹⁴ and hip fracture risk among Singapore Chinese¹⁵, which indicated the potential of the AHEI 2010 to assess Chinese diet quality in terms of validated health benefit.

One recent study showed poor diet quality with a steady improvement in diet quality, assessed by AHEI-2010 scores, among the US population between 1999 and 2010¹⁶. To date, no study has investigated long-term trends in index-based diet quality and related sociodemographic characteristics in China.

The present study aimed to examine the long-term Chinese diet quality transition and the role that sociodemographic characteristics play in the transition. We therefore applied the tAHEI to assess overall diet quality and used longitudinal quantile regression to estimate trends in percentiles of diet quality and socioeconomic influence among adults ages 18 to 65 in China from 1991 to 2011 of the China Health and Nutrition Survey (CHNS).

SUBJECTS AND METHODS

Study Population

All data used in this study were derived from the CHNS, an ongoing longitudinal study. Initiated in 1989, it focuses on assessing the relationships between the social and economic transformation in China and the resulting effects on the health and nutritional status of the Chinese population^{9, 17}.

The CHNS used a multistage, random cluster process to draw the sample from eight provinces, and then 24 communities in each province were selected randomly as the primary sampling units. In each type of community, 20 households were randomly selected, and all individuals in each household were surveyed for all data in each wave. The sampling procedure has been described in detail elsewhere^{5, 17}. The CHNS has completed nine rounds (1989, 1991, 1993, 1997, 2000, 2004, 2006, 2009, and 2011). We used data from 1991 to 2011, because only adults ages 20 to 45 were involved in 1989. Of all 18- to 65-year-old participants who had complete socioeconomic and dietary data, we excluded women currently pregnant or lactating during a survey year and those having implausible energy intakes (< 800 kilocalories [kcal] per day or > 6,000 kcal for men and < 600 kcal or > 4,000 kcal for women)¹⁸. We also excluded participants with only one wave of data. Our final sample included 13,853 participants (6,876 men; 6,977 women) clustered in 234 communities with an average of 4.1 responses from each subject (56,319 responses).

The protocols of the survey were approved by the institutional review committees of the University of North Carolina at Chapel Hill and the National Institute of Nutrition and Food Safety, Chinese Center for Disease Control and Prevention. All subjects provided written informed consent for their participation in the protocols.

Assessment of Diet Quality

Dietary intake was assessed by collecting three consecutive 24-hour dietary recalls (2 weekdays and 1 weekend day) for each individual combined with a household weighing inventory of all available foods over the same three periods and an interviewer-administered past year food frequency questionnaire of sugar sweetened beverages (SSB) and fruit juices since 2004 and alcohol consumption since 1993. The dietary data collection details have been described elsewhere^{19, 20}.

We used the tAHEI, tailored from the Harvard AHEI-2010⁸, to assess diet quality. The tailoring methods are described in detail elsewhere¹⁴. In general, major tailoring includes: (1) change in scale from serving to grams for vegetables, whole fruits, nuts and legumes, red/processed meat, SSBs and fruits juices; (2) estimation of alcohol and SSB and fruit juices intake from the past year FFQ in available survey years to reduce the potential underestimation of 24 h recall due to episodic consumption; (3) replacement the whole grain component with cereal fiber component due to extremely low intake and lack of variation in Chinese adults; (4) scaling only fresh red meat intake to increase the variation given that Chinese processed red meat intake is extremely low (about 3.1% of total meat) and few adults consumed processed meat higher than 64g²¹; (5) linking all Chinese foods to the US Food and Drug Administration (USDA) Food and Nutrient Database for Dietary Studies

(FNDDS)²² and National Nutrient Database for Standard Reference²³ to estimate polyunsaturated fatty acid (PUFA) and long chain (ω -3) fatty acid intake; (6) omission trans fat component in the tAHEI given a lack of information on trans fat composition of all eaten food in both China and USDA food composition tables (FCT). Supplemental Table 1 illustrates the components and scaling methods of the tAHEI. We used the three-day average intakes of total energy, nutrients, and foods and food groups to calculate total tAHEI scores ranging from 0 to 100. A higher score indicates a better diet quality and a score of 100 points indicates optimal diet quality.

Assessment of Sociodemographic Factors

Trained interviewers used standard questionnaires to collect information on household and individual incomes, individual education levels, and community environments. We calculated per capita annual family income by dividing annual family income by household size in each survey, inflated the income to 2011 values by adjusting for the consumer price index, and then categorized incomes as wave-specific tertiles. We grouped individual education levels into less than primary school, primary school, and higher than primary school.

The community urbanicity index, a complex measure of urbanization, is based on 12 multidimensional components reflecting the heterogeneity in economic, social, demographic, and infrastructural changes at the community level²⁴. We categorized the continuous urbanicity index into wave-specific tertiles. We also included the geographic regions (northern, central, and southern) due to the different dietary intakes shown in previous studies^{25, 26}.

Statistical Analysis

We first present descriptive statistics (means and standard deviations for continuous variables, and frequencies for categorical variables) on socioeconomic factors of interest stratified by sex for adults by survey years. We used chi-square tests for categorical variables, analysis of variance tests for continuous variables, and nonparametric median test for median of tertiles to test differences among survey years. For cross-sectional analysis, we first conducted a Kolmogorov-Smirnov test for normality and confirmed that the distribution for tAHEI is not normal. We then estimated the quantiles of the tAHEI scores based on simple frequencies. We also used LMS method [the L curve (Yeo-Johnson to remove skewness), M curve (median) and S curve (coefficient of variation)]²⁷ to present graphically smoothing probability density distribution of the tAHEI scores for select waves of the CHNS.

Next we used the longitudinal quantile regression method^{28, 29} to see whether secular trends and associations of diet quality vary at different percentiles of the tAHEI scores after adjustment for all potential covariates. For comparison, we also performed three-level [survey years (level 1) for individuals (level 2) nested in communities (level 3)] mixed-effect linear random intercept regression models³⁰ to estimate average secular trend of the total tAHEI scores and each component score, adjusting for all potential covariates including survey year, baseline age, income, education, urbanicity, geographic regions and total energy

intake. We included baseline age as continuous and modeled survey years as dummy variables rather than continuous, given potentially uneven changes over time.

Lastly, we introduced statistically significant time-sociodemographic product terms (time-income, time-education, time-urbanicity, and time-region) based on likelihood ratio test into linear mixed-effect model and used this fully adjusted model for predicted mean of tAHEI scores over time across socio-demographic subpopulations, respectively.

All statistical tests were two-tailed, and we regarded differences as significant at $p < 0.05$. We fitted three-level mixed-effect linear models using Stata/SE version 14.1, plotted density and quantile distribution using the VGAM package and performed quantile regression models using the LQMM package in R version 2.15.1. For all other descriptive analyses, we used SAS version 9.3.

RESULTS

As shown in Table 1, the energy-adjusted mean tAHEI scores and 95% confidence intervals (CI) increased from 36.9 (36.7–37.1) points in 1991 to 50.3 (50.1–50.5) points in 2011 for men ($p < 0.001$) and from 35.6 (35.4–35.8) to 46.9 (46.7–47.1) points for women ($p < 0.001$), both with bigger increase between 2009 and 2011 (men 8.2; women 12.5) than between other adjacent survey years. Moreover, distributions of education level and geographic region and mean baseline age were significantly different across survey years ($p < 0.05$). The median of each income and urbanicity index tertiles increased significantly over time ($p < 0.05$).

Shifts in the Distribution of the unadjusted tAHEI Scores in Chinese Adults

Supplemental Table 2 presents mean and distribution (at the 10th, 25th, 50th, 75th, 90th, and 95th percentiles of the unadjusted tAHEI scores along with 95% CI from 1991 to 2011 for men and women, respectively. We can see the degree of increase in unadjusted tAHEI score was uneven, with upper percentile score having larger rate of increase than lower percentile. The increase in the 95th percentile from 1991 to 2011 was around 18.4 points in men and 14.8 points in women, whereas the increase for the median was 13.6 and 11.6 and for the 10th percentile only 7.3 and 4.7, respectively.

We also show graphically the shifts in the distribution of the unadjusted tAHEI scores using LMS methods in men and women for selected survey years (Supplemental Figure 1). For both sexes the distribution generally shifts to the right, with the distributions becoming wider and flatter with a larger proportion of the subjects having higher tAHEI scores over time. We also observed different degrees of shifts in the whole distribution over time. For men the shift was gradually flattening over time, with slightly larger shifts between 2009 and 2011. In contrast, for women there was a remarkable rightward shift and flattening only between 2006 and 2011 with no significant changes before 2006.

Secular trends in covariate-adjusted tAHEI score at different percentiles

As shown in Tables 2 and Table 3, increasing trends in the tAHEI score occurred at various percentiles for both sexes after adjustment for all potential covariates. We also observed

gradually steeper rate of increase with increasing percentiles of the tAHEI score and remarkable upward jump between 2009 and 2011 at various percentiles. In addition, the median (50th percentile) estimates from quantile regression shows fairly steeper than the mean estimate of secular trends for both sexes.

For men, after additional adjustment for income and education (model 2), only the 25th percentile estimates in 2011 as compared to 1991 attenuated by 1.0 point. After adjustment for urbanicity and geographic region, (model 3) the 25th and the 95th percentile estimates in 2011 attenuated by 1.1 points and 0.7 points, respectively.

For women, the coefficients at various percentiles did not change significantly from adjustment for income and education in model 2. Then only the 25th percentile estimates in 2011 attenuated by 0.8 points from additional adjustment from urbanicity and geographic region.

Secular trends in covariate-adjusted mean tAHEI component score

Figure 1 illustrates how tAHEI component scores changed over time. For both sexes, PUFAs, EPA and DHA scores showed the most remarkable increasing trends between 2009 and 2011. Compared with the scores in 1991, PUFAs and EPA plus DHA scores in 2011 increased by 6.8 points and 5.3 points in men and 7.0 points and 5.3 points in women, respectively. In contrast, the cereal fiber score, SSB score and red meat score showed slight declines for both sexes. In addition, whole fruits score and nuts and legumes score showed slight increases by about 1.0 points. Alcohol scores, on the other hand, increased by 2.3 points only among men over the 21-year period.

Table 4 presents predicted mean of the tAHEI score across demographic and socioeconomic factors from a fully adjusted mixed-effect linear regression. Figure 2 graphically illustrates potential disparity in tAHEI score transition. The increasing tAHEI score within northern adults indicated larger increase than that seen among southern adults and the difference in tAHEI score between northern and southern adults increased significantly from 2.4 points in 1991 to 7.2 points in 2011 in men and from 1.8 points to 6.6 points in women. High urbanicity was associated with lower tAHEI score before 2004 but with higher diet quality after 2004 for both sexes given that slightly large increase in lower urbanicity adults over time. In addition, men with primary school had slightly higher scores than those with other corresponding groups in 2011, whereas tAHEI scores in women were not significantly different across income or education groups in 2011.

DISCUSSION

Using data from the CHNS between 1991 and 2011, we found a moderate increase across the entire distribution of the tAHEI scores with the most remarkable improvement between 2009 and 2011. Our study also showed the diet quality transition varied greatly across score percentiles and that adults with higher percentile score tended to have larger increase than those with lower percentile score over time. Improvements in diet quality were mainly attributable to increased scores of PUFAs, EPA and DHA. In addition, improved diet quality was observed in all sociodemographic groups with significant sociodemographic disparity.

Limited studies have used the AHEI-2010 to assess diet quality and/or to examine longitudinal trends. One recent study in a nationally representative sample of 29,124 US adults aged 20 to 85 years indicated increased energy-adjusted AHEI-2010 scores without the *trans* fat component from 34.2 (33.1–35.2) in 1999 to 2000 to 37.1 (36.6–37.7) in 2009 to 2010¹⁶. Increased scores in US adults mainly resulted from an increase of 0.9 points for SSB and fruit juice, 0.7 points for whole fruit, 0.5 points for whole grains, and 0.4 points for nuts and legumes and slight decrease in sodium¹⁶. Our study observed an increase of 11.3 points for Chinese men and women ages 18 to 65 between 2000 and 2011. Chinese adults had better diet quality and gained larger improvement in diet quality than US adults in the past decade. Different from that in US adults, the changed profiles in Chinese adults were characterized by an increase 7.0 points for PUFAs, 5.3 points for long chain (ω -3) fatty acids score, 1.0 points for whole fruit and nuts and legume, as well as slight decline in the scores of cereal fiber, SSB and fruit juices, and red meat over time. Significant increases in the intake of edible oils^{2, 31} and sea foods^{2, 32} in Chinese adults in the past two decades may be a possible explanation for great increase in PUFAs and EPA plus DHA score. However, such comparison should be understood with caution given that the tAHEI was not quite identical to AHEI-2010 due to tailoring.

Previous studies using the CHNS data examined the role of sociodemographic characteristics for changes in specific nutrients, foods or food groups in Chinese adults: positive association of income with intakes of animal-source foods^{31, 33, 34}, edible oils^{31, 33, 34} and percentage of energy from dietary fat^{33, 34} and negative association of income with cereal intake³¹, relationship of higher urbanicity with higher intake of total animal-source foods and lower intake of coarse grains in adults ages 18 to 59 years between 1991 and 2011². Our results add valuable information on sociodemographic disparity in Chinese diet quality transition. We found wider gaps in diet quality between southern and northern adults over time and lower diet quality in lower urbanized adults since 2004. Income and education levels were not associated with women's diet quality, while gaps in diet quality across income or education in men narrowed. In contrast, Wang et al. found that US adults with higher socioeconomic status had greater improvement in diet quality, assessed by AHEI-2010, and the gaps in diet quality between higher and lower SES adults widen over 12-year period^{16, 35}. One recent study using NHANES cycles (1999–2012) also showed that the disparities in the diet scores, constructed based on the AHA 2020 Strategic Impact Goals for diet, widen over time with smaller overall improvements in those with lower levels of family income and lower education levels in US adults aged 20 year or older³⁶.

Our study is the first to investigate 21-year trends in index-based diet quality distribution in a large longitudinal sample in a country undergoing rapid nutrition transitions. The use of tAHEI was an easier and more practical way to standardize and track diet quality transitions than research on specific nutrients, foods or food groups over time^{2, 4, 31, 32, 37}. The longitudinal quantile regression and mixed-effect linear regression we used are well suited for the repeated diet measure in the CHNS and best represent trends in diet quality distribution and transition. Both approaches have the ability to handle unbalanced panel data, insensitivity to outliers, and provide more precise and robust estimates than ordinary least square regressions. Moreover, quantile analysis can provide multiple estimates to

reflect characteristics of diet quality transition across the whole distribution. Survey years modeled as dummy variables contribute to discovering the uneven degree of diet quality transition over time instead of a continuous coding presenting average changes per year.

Several limitations should be considered. First, the tailored tAHEI was not identical to the AHEI-2010 and may not reflect the nature of the AHEI-2010. For example, we may underestimate cereal fiber intake due to limited insoluble fibre in China FCT. Fatty acid composition may be different due to country-specific food types and planting conditions. We omitted a trans fat component, but it was shown as one important dimension of the overall diet quality^{2, 38, 39}. In addition, the tAHEI does not consider cooking methods or eating behaviors. Previous studies reported a marked increase in the proportion of energy from deep-fried and stir-fried foods and from snacking over time^{2, 40}, which may be important aspect of overall diet quality in relation to disease risk.

In conclusion, Chinese adults gained moderate improvements in the overall quality across the whole distribution between 1991 and 2011. However, Chinese diet quality is still far from optimal (about 50.0 points out of 100 full points) with sociodemographic disparity. From the perspective of public health, web-based key technology of nutrition interventions and government-dominated policy efforts should be given priority to adults with low diet quality who generally had low income, lived in low urbanized communities and lived in southern China.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding

We thank the National Institute for Nutrition and Health, China Center for Disease Control and Prevention; the Carolina Population Center, University of North Carolina at Chapel Hill; the National Institutes of Health (NIH) (R01-HD30880, DK056350, 5 R24 HD050924, R01-HD38700, R01-HL108427 and R01-DK104371); and the Fogarty NIH grant 5 D43 TW009077 for financial support for the CHNS data collection and analysis files from 1989 to 2011 and future surveys and the China-Japan Friendship Hospital, Ministry of Health, for support for the CHNS 2009.

REFERENCES

1. World Health Organization. Global status report on noncommunicable disease 2010. 2011.
2. Zhai FY, Du SF, Wang ZH, Zhang JG, Du WW, Popkin BM. Dynamics of the Chinese diet and the role of urbanicity, 1991–2011. *Obes Rev*. 2014; (15 Suppl 1):16–26. [PubMed: 24341755]
3. Popkin BM. Synthesis and implications: China's nutrition transition in the context of changes across other low- and middle-income countries. *Obes Rev*. 2014; (15 Suppl 1):60–67. [PubMed: 24341759]
4. Xiao YT, Su C, Ouyang YF, Zhang B. Trends of vegetables and fruits consumption among Chinese adults aged 18 to 44 years old from 1991 to 2011. *Zhonghua liu xing bing xue za zhi = Zhonghua liuxingbingxue zazhi*. 2015; 36(3):232–236. [PubMed: 25975399]
5. Moeller SM, Reedy J, Millen AE, Dixon LB, Newby PK, Tucker KL, et al. Dietary patterns: challenges and opportunities in dietary patterns research an Experimental Biology workshop, April

- 1, 2006. Journal of the American Dietetic Association. 2007; 107(7):1233–1239. [PubMed: 17604756]
6. Willett WC, Stampfer MJ. Current evidence on healthy eating. Annual review of public health. 2013; 34:77–95.
7. McCullough ML, Feskanich D, Stampfer MJ, Giovannucci EL, Rimm EB, Hu FB, et al. Diet quality and major chronic disease risk in men and women: moving toward improved dietary guidance. Am J Clin Nutr. 2002; 76(6):1261–1271. [PubMed: 12450892]
8. Chiuve SE, Fung TT, Rimm EB, Hu FB, McCullough ML, Wang M, et al. Alternative dietary indices both strongly predict risk of chronic disease. The Journal of nutrition. 2012; 142(6):1009–1018. [PubMed: 22513989]
9. Hu FB. Dietary pattern analysis: a new direction in nutritional epidemiology. Curr Opin Lipidol. 2002; 13(1):3–9. [PubMed: 11790957]
10. Jacques PF, Tucker KL. Are dietary patterns useful for understanding the role of diet in chronic disease? Am J Clin Nutr. 2001; 73(1):1–2. [PubMed: 11124739]
11. Newby PK, Tucker KL. Empirically derived eating patterns using factor or cluster analysis: a review. Nutr Rev. 2004; 62(5):177–203. [PubMed: 15212319]
12. Willett WC, McCullough ML. Dietary pattern analysis for the evaluation of dietary guidelines. Asia Pac J Clin Nutr. 2008; (17 Suppl 1):75–78. [PubMed: 18296306]
13. Kant AK. Dietary patterns and health outcomes. J Am Diet Assoc. 2004; 104(4):615–635. [PubMed: 15054348]
14. Wang ZH, Siega-riz AM, Gordon-Larsen P, Cai JW, Adair LS, Zhang B, et al. Diet Quality and the Risk of Type 2 Diabetes and Major Cardiometabolic Risk Factors among Adults in China. 2015 (in press).
15. Dai Z, Butler LM, van Dam RM, Ang LW, Yuan JM, Koh WP. Adherence to a vegetable-fruit-soy dietary pattern or the Alternative Healthy Eating Index is associated with lower hip fracture risk among Singapore Chinese. The Journal of nutrition. 2014; 144(4):511–518. [PubMed: 24572035]
16. Wang DD, Leung CW, Li Y, Ding EL, Chiuve SE, Hu FB, et al. Trends in dietary quality among adults in the United States, 1999 through 2010. JAMA internal medicine. 2014; 174(10):1587–1595. [PubMed: 25179639]
17. Popkin BM, Du S, Zhai F, Zhang B. Cohort Profile: The China Health and Nutrition Survey--monitoring and understanding socio-economic and health change in China, 1989–2011. International journal of epidemiology. 2010; 39(6):1435–1440. [PubMed: 19887509]
18. Wang Z, Zhang B, Wang H, Zhang J, Du W, Su C, et al. Study on the multilevel and longitudinal association between meat consumption and changes in body mass index, body weight and risk of incident overweight among Chinese adults. Chin J Epidemiol. 2013; 34(7):1–7.
19. Tee ES. Development and promotion of Malaysian Dietary Guidelines. Asia Pac J Clin Nutr. 2011; 20(3):455–461. [PubMed: 21859667]
20. Popkin BM, Lu B, Zhai F. Understanding the nutrition transition: measuring rapid dietary changes in transitional countries. Public health nutrition. 2002; 5(6A):947–953. [PubMed: 12633520]
21. Ng SW, Howard, Annie-Green, Wang, Su, Chang, Zhang, Bing. The Physical Activity Transition among adults in China: 1991–2011. Obes Rev. 2014; (15 Suppl 1):27–36.
22. Ahuja, J., Montville, J., Omolewa, TG., Heendeniya, K., Martin, C., Steinfeldt, L., et al. USDA Food and Nutrient Database for Dietary Studies, 5.0. USDA. In. Beltsville, MD: USDA Agricultural Research Service, Food Surveys Research Group; 2014.
23. USDA Agricultural Research Service. USDA National Nutrient Database for Standard Reference, release 25. Nutrient Data Laboratory Home Page. 2014. Nutrient Data Laboratory Home Page
24. Jones-Smith JC, Popkin BM. Understanding community context and adult health changes in China: development of an urbanicity scale. Social science & medicine. 2010; 71(8):1436–1446. [PubMed: 20810197]
25. Du S, Batis C, Wang H, Zhang B, Zhang J, Popkin BM. Understanding the patterns and trends of sodium intake, potassium intake, and sodium to potassium ratio and their effect on hypertension in China. Am J Clin Nutr. 2014; 99(2):334–343. [PubMed: 24257724]
26. Ge K. The transition of Chinese dietary guidelines and food guide pagoda. Asia Pac J Clin Nutr. 2011; 20(3):439–446. [PubMed: 21859664]

27. Cole TJ, Freeman JV, Preece MA. British 1990 growth reference centiles for weight, height, body mass index and head circumference fitted by maximum penalized likelihood. *Statistics in medicine*. 1998; 17(4):407–429. [PubMed: 9496720]
28. Koenker R. Quantile Regression for Longitudinal Data. *Journal of Multivariate Analysis*. 2004; 91:74–89.
29. Koenker R, Bassett G. Regression quantile. *Econometrica*. 1978; 46:33–50.
30. Skrondal, SR-HaA. *Multilevel and Longitudinal Modeling Using Stata*. Second. Stata press; 2005.
31. Du S, Mroz TA, Zhai F, Popkin BM. Rapid income growth adversely affects diet quality in China--particularly for the poor! *Social science & medicine*. 2004; 59(7):1505–1515. [PubMed: 15246178]
32. Wang ZH, Zhai FY, Wang HJ, Zhang JG, Du WW, Su C, et al. Secular trends in meat and seafood consumption patterns among Chinese adults, 1991–2011. *European journal of clinical nutrition*. 2015; 69(2):227–233. [PubMed: 25351649]
33. Wang Z, Zhai F, He Y, Wang H. Influence of family income on dietary nutrients intake and dietary structure in China. *Wei sheng yan jiu = Journal of hygiene research*. 2008; 37(1):62–64. [PubMed: 18421867]
34. Wang ZH, Zhai FY, Wang HJ, Zhang JG, Du WW, Su C, et al. Influence of Household Income Level on Food Consumption in Chinese Population. *Food and Nutrition In China*. 2015; 21(3):46–49.
35. Sugiyama T, Shapiro MF. The growing socioeconomic disparity in dietary quality: mind the gap. *JAMA internal medicine*. 2014; 174(10):1595–1596. [PubMed: 25178585]
36. Rehm CD, Peñalvo JL, Afshin A, Mozaffarian D. Dietary Intake Among US Adults, 1999–2012. *Jama the Journal of the American Medical Association*. 2016; 315(23):2542–2553. [PubMed: 27327801]
37. Wang HJ, Zhang B, Du WW, Liu AD, Zhang JG, Wang ZH, et al. Trends of the dietary fiber intake among Chinese aged 18 – 45 in nine provinces (autonomous region) from 1989 to 2006. *Zhonghua yu fang yi xue za zhi [Chinese journal of preventive medicine]*. 2011; 45(4):318–322.
38. Zamora D, Gordon-Larsen P, He K, Jacobs DR Jr, Shikany JM, Popkin BM. Are the 2005 Dietary Guidelines for Americans Associated With reduced risk of type 2 diabetes and cardiometabolic risk factors? Twenty-year findings from the CARDIA study. *Diabetes care*. 2011; 34(5):1183–1185. [PubMed: 21478463]
39. Wallingford JC, Yuhas R, Du S, Zhai F, Popkin BM. Fatty acids in Chinese edible oils: value of direct analysis as a basis for labeling. *Food and nutrition bulletin*. 2004; 25(4):330–336. [PubMed: 15646310]
40. Wang Z, Zhai F, Zhang B, Popkin BM. Trends in Chinese snacking behaviors and patterns and the social-demographic role between 1991 and 2009. *Asia Pac J Clin Nutr*. 2012; 21(2):253–262. [PubMed: 22507613]

Figure 1. Estimated shifts in covariate-adjusted tAHEI component score in Chinese men (a) and women (b) in China, CHNS, 1991–2011 (estimation based on mixed-effect linear regression models adjustment for survey year, baseline age, income, education, geographic region, urbanicity, and total energy intake)

Figure 2. Predicted tAHEI score across demographic and socioeconomic factors in Chinese adults, CHNS, 1991–2011 (estimation based on mixed-effect linear regression models adjustment for baseline age, total energy intake, survey years, income, education, geographic region, urbanicity and corresponding interaction terms with survey years)

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1

Cross-sectional characteristics of the study sample in the CHNS, 1991–2011

	Men								Women							
	1991	1993	1997	2000	2004	2006	2009	2011	1991	1993	1997	2000	2004	2006	2009	2011
N	3,210	3,473	3,608	3,998	3,578	3,496	3,493	3,063	3,342	3,581	3,530	3,967	3,632	3,634	3,544	3,170
Energy-adjusted tAHEI score <i>a</i> *	36.9 ± 0.2	39.3 ± 0.2	39.0 ± 0.2	38.8 ± 0.2	40.4 ± 0.2	41.6 ± 0.2	42.1 ± 0.2	50.3 ± 0.2	35.6 ± 0.2	36.7 ± 0.1	35.9 ± 0.2	35.2 ± 0.1	36.4 ± 0.1	37.6 ± 0.1	38.2 ± 0.2	46.9 ± 0.2
Age (years) <i>a</i> *	38.0 ± 0.2	37.2 ± 0.2	36.1 ± 0.2	35.4 ± 0.2	35.5 ± 0.2	35.3 ± 0.2	34.7 ± 0.2	34.7 ± 0.2	38.3 ± 0.2	37.7 ± 0.2	37.2 ± 0.2	36.4 ± 0.2	36.3 ± 0.2	35.9 ± 0.2	36.0 ± 0.2	36.0 ± 0.2
Income tertile <i>b</i> *																
Low	1180.2	1188.5	1538.2	1617.9	1869.9	2172.3	3560.2	4346.2	1213.5	1188.2	1536.6	1651.1	1858.6	2014.7	3386.2	3834.3
Medium	2659.6	2785.5	3472.7	4339.3	5235.9	6081.4	9000.6	11341.1	2666.3	2785.1	3499.0	4461.4	5178.2	5856.5	8820.5	10678.7
High	4798.5	5689.9	6808.0	8679.0	12248.5	14353.9	19872.6	24033.5	4757.9	5763.0	6901.2	8810.2	12131.5	13982.2	19786.4	22687.7
Education (%) *																
Less than primary school	18.9	19.4	16.2	11.7	8.9	11.5	10.1	10.3	40.5	40.0	39.0	29.2	25.5	27.1	24.2	24.3
Primary school	21.5	23.6	23.5	23.0	22.7	18.8	19.9	19.2	18.5	20.5	21.2	23.3	24.3	19.7	22.4	22.3
Higher than primary school	59.6	57.0	60.3	65.3	68.4	69.8	70.0	70.5	41.0	39.5	39.8	47.5	50.2	53.2	53.4	53.4
Urbanicity tertile <i>b</i> *																
Low	28.8	29.2	31.5	38.0	39.8	40.3	46.5	48.2	28.8	29.3	31.7	40.3	39.9	40.3	46.6	48.2
Medium	44.1	46.8	50.6	53.8	57.8	61.8	63.0	68.0	44.5	46.9	51.3	55.7	59.0	61.9	63.1	68.0
High	63.2	66.2	72.3	79.2	84.2	86.0	89.3	91.0	63.9	66.5	72.8	79.7	85.0	86.0	89.5	91.0
Geographic region (%) *																
Northern	13.3	12.0	5.0	20.3	21.1	22.2	21.5	23.0	12.8	11.8	4.8	21.0	22.9	22.2	22.5	23.9
Central	36.6	36.0	38.3	33.8	32.0	32.3	32.6	31.7	40.6	37.5	41.1	34.5	32.8	33.9	33.3	31.5
Southern	50.2	52.0	56.7	45.9	46.9	45.5	45.9	45.3	46.6	50.7	54.1	44.5	44.3	43.8	44.2	44.6

^aMean ± standard error;

^bMedian values by survey year-specific tertiles;

^{*}Statistically significant difference among survey years based on chi-square tests, analysis of variance tests or nonparametric median test for median of tertiles, p < 0.05.

Table 2

Coefficients (95% CI) from quantile regression versus mixed-effect regression of the tAHEI score on survey year and socioeconomic factors in Chinese men

Mean regression, <i>a</i> Coefficient (95% CI) Mean		Quantile regression, coefficient (95% CI)				
		25th	50th	75th	85th	95th
Model 1						
1993	2.3(1.8,2.7) ***	1.7(1.2,2.1) ***	2.2(1.8,2.6) ***	2.9(2.5,3.4) ***	3.2(2.6,3.9) ***	3.2(2.3,4.0) ***
1997	2.2(1.8,2.7) ***	2.2(1.8,2.7) ***	2.2(1.7,2.6) ***	2.6(2.2,3.1) ***	3.0(2.5,3.5) ***	3.0(1.8,4.2) ***
2000	1.8(1.3,2.2) ***	1.4(1.0,1.8) ***	1.7(1.3,2.1) ***	2.5(2.0,2.9) ***	3.0(2.5,3.6) ***	3.7(2.8,4.6) ***
2004	3.3(2.9,3.8) ***	2.2(1.7,2.7) ***	3.0(2.6,3.4) ***	4.7(4.1,5.3) ***	6.3(5.6,6.9) ***	6.2(5.1,7.3) ***
2006	4.7(4.2,5.1) ***	2.9(2.4,3.5) ***	4.3(3.7,4.9) ***	6.4(5.7,7.0) ***	7.9(7.2,8.7) ***	9.2(7.8,10.6) ***
2009	4.9(4.5,5.4) ***	2.6(2.1,3.0) ***	4.4(3.9,4.9) ***	7.5(6.9,8.2) ***	9.6(8.8,10.3) ***	10.4(8.9,11.9) ***
2011	13.1(12.7,13.6) ***	11.1(10.5,11.8) ***	14.6(13.9,15.2) ***	17.5(16.8,18.2) ***	18.7(17.7,19.7) ***	19.0(17.8,20.2) ***
Model 2						
1993	2.3(1.8,2.7) ***	2.3(0.8,3.8) ***	2.2(1.8,2.7) ***	3.0(2.3,3.6) ***	3.4(2.9,4.0) ***	3.4(2.6,4.3) ***
1997	2.2(1.8,2.7) ***	2.7(1.2,4.2) ***	2.2(1.8,2.6) ***	2.7(2.1,3.2) ***	3.2(2.6,3.8) ***	3.1(2.2,4.1) ***
2000	1.8(1.3,2.2) ***	2.1(0.6,3.5) ***	1.7(1.3,2.1) ***	2.5(2.0,3.0) ***	3.3(2.7,3.9) ***	3.9(2.9,5.0) ***
2004	3.4(2.9,3.8) ***	3.2(1.7,4.8) ***	3.1(2.6,3.6) ***	4.9(4.3,5.4) ***	6.5(5.8,7.1) ***	6.4(5.5,7.3) ***
2006	4.7(4.2,5.1) ***	3.9(2.3,5.5) ***	4.3(3.7,4.8) ***	6.6(5.9,7.2) ***	8.2(7.5,9.0) ***	9.7(8.1,11.2) ***
2009	5.0(4.5,5.4) ***	3.5(1.9,5.1) ***	4.4(3.7,5.0) ***	7.7(6.9,8.4) ***	9.8(9.0,10.6) ***	10.5(9.2,11.9) ***
2011	13.1(12.7,13.6) ***	12.1(10.2,13.9) ***	14.6(14.1,15.1) ***	17.6(17.0,18.2) ***	18.9(18.2,19.7) ***	19.3(18.2,20.4) ***
Model 3						
1993	2.3(1.8,2.7) ***	1.7(0.6,2.8) ***	2.2(1.8,2.6) ***	3.1(2.5,3.7) ***	3.3(2.3,4.4) ***	2.9(1.9,3.8) ***
1997	2.3(1.8,2.7) ***	2.2(1.0,3.3) **	1.9(1.5,2.3) ***	2.8(2.3,3.3) ***	3.1(2.1,4.1) ***	2.6(1.5,3.8) ***
2000	1.8(1.3,2.2) ***	1.3(0.1,2.4) *	1.4(0.9,1.8) ***	2.5(1.9,3.1) ***	2.9(1.7,4.1) ***	3.1(1.9,4.3) ***
2004	3.4(2.9,3.8) ***	2.0(0.7,3.3) **	2.7(2.2,3.3) ***	4.9(4.2,5.6) ***	6.1(5.0,7.2) ***	6.3(4.7,7.9) ***
2006	4.7(4.2,5.1) ***	2.9(1.6,4.2) ***	4.0(3.4,4.5) ***	6.4(5.7,7.0) ***	7.6(6.4,8.9) ***	8.9(7.7,10.1) ***
2009	4.9(4.5,5.4) ***	2.3(1.3,7) **	4.0(3.5,4.6) ***	7.6(7.0,8.3) ***	9.5(8.3,10.7) ***	10.1(8.6,11.6) ***

	Mean regression, <i>a</i> Coefficient (95% CI) Mean	Quantile regression, coefficient (95% CI)				
		25th	50th	75th	85th	95th
2011	13.1(12.6,13.6) ***	10.9(9.3,12.6) ***	14.3(13.8,14.8) ***	17.4(16.7,18) ***	18.6(17.6,19.6) ***	18.6(17.4,19.8) ***

^aMixed-effect linear random intercept regression models.

^bModels adjusted for survey years (dummy variables) and baseline age (model 1), plus individual income (tertile) and education (less than primary, primary, and higher than primary) (model 2), additional urbanicity index (tertile) and geographic region (model3).

* p < 0.05,
** p < 0.01,
*** p < 0.001.

Coefficients (95% CI) from quantile regression versus mixed-effect regression of the tAHEI score on survey year and socioeconomic factors in Chinese women

Mean regression, ^a Coefficient (95% CI) Mean		Quantile regression, coefficient (95% CI)				
		25th	50th	75th	85th	95th
Model 1 ^b						
1993	1.1(0.7,1.5)***	0.5(0.1,0.9)***	1.0(0.7,1.4)***	1.5(1.1,2.0)***	1.8(1.3,2.4)***	1.9(1.3,2.6)***
1997	0.5(0.1,0.9)*	0.1(-0.4,0.5)	0.1(-0.3,0.5)	0.5(0.1,0.9)*	0.7(0.2,1.3)**	1.5(0.4,2.6)**
2000	-0.5(-0.9,-0.1)**	-0.9(-1.3,-0.6)***	-0.8(-1.2,-0.3)***	-0.4(-0.8,0)	-0.1(-0.7,0.4)	0.2(-0.7,1.0)
2004	0.6(0.2,1.0)**	-0.7(-1.0,-0.3)***	0.1(-0.4,0.5)	1.4(0.9,1.9)***	2.8(2.2,3.5)***	3.3(2.3,4.4)***
2006	1.9(1.5,2.3)***	0(-0.5,0.5)	1.1(0.5,1.7)***	3.1(2.5,3.7)***	4.9(4.2,5.7)***	5.3(4.2,6.5)***
2009	2.3(1.9,2.7)***	-0.4(-0.8,0.1)	1.3(0.8,1.7)***	4.4(3.7,5.0)***	6.8(6.0,7.7)***	8.0(6.6,9.3)***
2011	10.7(10.3,11.2)***	8.4(7.7,9.0)***	12.3(11.6,12.9)***	14.8(14.3,15.4)***	16.1(15.3,17.0)***	16.2(15.3,17.1)***
Model 2						
1993	1.1(0.7,1.5)***	0.6(-0.5,1.7)	1.1(0.8,1.5)***	1.5(1.1,2)***	1.8(1.3,2.4)***	1.9(1.2,2.7)***
1997	0.5(0.1,0.9)*	0(-1.2,1.3)	0.2(-0.1,0.6)	0.5(0.1,0)*	0.7(0.1,1.4)*	1.5(0.3,2.6)*
2000	-0.5(-0.9,-0.1)*	-0.8(-2.0,0.5)	-0.6(-1.0,-0.2)**	-0.4(-0.9,0.1)	0(-0.6,0.7)	-0.1(-1.1,1.0)
2004	0.6(0.2,1.0)**	-0.4(-1.8,0.9)	0.4(-0.1,0.8)	1.5(0.9,2.0)***	2.9(2.1,3.6)***	3.2(2.2,4.1)***
2006	1.9(1.5,2.3)***	0.1(-1.2,1.5)	1.2(0.8,1.7)***	3.2(2.5,3.8)***	4.9(4.1,5.8)***	5.3(3.8,6.9)***
2009	2.4(2.0,2.8)***	-0.2(-1.4,1.1)	1.4(0.9,1.8)***	4.6(3.8,5.4)***	7.0(6.0,8.0)***	8.0(6.6,9.3)***
2011	10.8(10.3,11.2)***	8.6(7.0,10.2)***	12.5(12,13)***	14.9(14.3,15.5)***	16.1(15.2,16.9)***	16.3(15.3,17.4)***
Model 3						
1993	1.1(0.7,1.5)***	1.5(0.6,2.5)**	1.0(0.6,1.5)***	1.7(1.2,2.2)***	1.6(1.0,2.3)***	1.8(0.5,3.0)***
1997	0.5(0.1,0.9)*	1.2(0.1,2.2)*	0.3(-0.2,0.7)	0.6(0.1,1.1)*	0.7(0.1,1.4)*	1.4(0.3,2.5)*
2000	-0.5(-0.9,-0.1)**	0(-1.0,9)	-0.7(-1.1,-0.4)	-0.4(-0.8,0.1)	-0.3(-0.9,0.3)	0.3(-0.8,1.4)
2004	0.6(0.2,1.0)**	0.3(-0.8,1.4)	0.1(-0.3,0.5)***	1.6(1.1,2.2)***	2.5(1.8,3.1)***	4.1(2.8,5.4)***
2006	1.9(1.5,2.3)***	1.0(-0.1,2.2)	1.2(0.8,1.6)***	3.3(2.7,3.9)***	4.6(4.0,5.3)***	5.2(3.6,6.8)***

	Mean regression, <i>a</i> Coefficient (95% CI) Mean	Quantile regression, coefficient (95% CI)				
		25th	50th	75th	85th	95th
2009	2.3(1.9,2.8)***	0.8(−0.3,1.9)	1.2(0.7,1.7)***	4.5(3.8,5.2)***	6.6(5.7,7.6)***	8.0(6.4,9.6)***
2011	10.8(10.3,11.2)***	9.8(8.2,11.3)	12.2(11.7,12.7)***	15.0(14.5,15.5)***	15.8(15.1,16.5)***	16.1(14.8,17.3)***

^a Mixed-effect linear random intercept regression models.

^b Models adjusted for survey years (dummy variables) and baseline age (model 1), plus individual income (tertile) and education (less than primary, primary, and higher than primary) (model 2), additional urbanicity index (tertile) and geographic region (model3).

* p < 0.05,
** p < 0.01,
*** p < 0.001.

Table 4
Predicted tAHEI score by demographic and socioeconomic factors, CHNS, 1991–2011

Characteristic	1991	1993	1997	2000	2004	2006	2009	2011
Men^a								
Income								
Low	37.0(0.4)	40.0(0.3)	40.0(0.3)	39.0(0.3)	41.1(0.3)	42.2(0.3)	41.8(0.3)	49.7(0.3)
Medium	37.1(0.3)	39.6(0.3)	39.1(0.3)	39.0(0.3)	40.8(0.3)	41.9(0.3)	42.2(0.3)	50.8(0.3)
High	36.8(0.3)	38.6(0.3)	38.8(0.3)	38.7(0.3)	39.9(0.3)	40.9(0.3)	42.7(0.3)	50.2(0.4)
Education								
Less than primary school	36.9(0.4)	39.9(0.4)	39.6(0.4)	38.4(0.5)	41.6(0.6)	41.7(0.5)	43.3(0.5)	49.6(0.6)
Primary school	37.1(0.4)	40.1(0.4)	39.3(0.4)	38.6(0.4)	40.3(0.4)	41.1(0.4)	42.4(0.4)	51.0(0.4)
Higher than primary school	37.0(0.3)	39.1(0.3)	39.2(0.3)	39.1(0.3)	40.6(0.3)	41.9(0.3)	42.0(0.3)	50.1(0.3)
Geographic region								
Northern	38.1(0.6)	42.2(0.6)	42.2(0.6)	39.9(0.5)	41.0(0.5)	44.6(0.5)	44.2(0.5)	55.8(0.5)
Central	38.3(0.4)	40.8(0.4)	40.8(0.4)	40.7(0.4)	41.5(0.4)	43.6(0.4)	43.5(0.4)	49.7(0.4)
Southern	35.7(0.4)	37.3(0.3)	37.1(0.3)	37.2(0.3)	39.9(0.4)	39.2(0.4)	40.6(0.4)	48.6(0.4)
Urbanicity								
Low	37.7(0.4)	39.0(0.4)	39.1(0.4)	40.1(0.3)	41.7(0.4)	41.4(0.4)	42.2(0.4)	50.0(0.4)
Medium	37.7(0.4)	40.4(0.3)	40.1(0.3)	39.0(0.3)	40.2(0.3)	41.4(0.3)	41.9(0.3)	49.7(0.4)
High	35.5(0.4)	38.8(0.4)	38.6(0.4)	37.6(0.3)	40.0(0.4)	42.2(0.4)	42.6(0.4)	51.1(0.4)
Women								
Income								
Low	35.9(0.3)	37.3(0.3)	36.7(0.3)	35.6(0.3)	36.9(0.3)	38.1(0.3)	38.2(0.3)	46.4(0.3)
Medium	35.6(0.3)	37.0(0.3)	36.1(0.3)	35.3(0.3)	37.1(0.3)	37.4(0.3)	38.5(0.3)	47.1(0.3)
High	35.2(0.3)	36.0(0.3)	35.8(0.3)	35.2(0.3)	36.1(0.3)	37.5(0.3)	38.6(0.3)	46.8(0.3)
Education								
Less than primary school	35.6(0.3)	37.1(0.3)	36.4(0.3)	35.5(0.3)	38.2(0.3)	37.5(0.3)	38.5(0.3)	46.8(0.4)
Primary school	36.0(0.4)	36.7(0.4)	35.8(0.3)	34.8(0.3)	36.1(0.3)	37.5(0.4)	38.7(0.4)	46.6(0.4)
Higher than primary school	35.4(0.3)	36.6(0.3)	36.2(0.3)	35.5(0.3)	36.1(0.3)	37.9(0.3)	38.3(0.3)	46.8(0.3)

Characteristic	1991	1993	1997	2000	2004	2006	2009	2011
Geographic region								
Northern	36.1(0.6)	38.0(0.6)	37.5(0.6)	35.6(0.5)	36.9(0.5)	39.9(0.5)	39.8(0.5)	51.7(0.5)
Central	36.9(0.4)	38.4(0.4)	38.1(0.4)	37.1(0.4)	37.3(0.4)	39.2(0.4)	39.7(0.4)	46.4(0.4)
Southern	34.3(0.3)	35.0(0.3)	34.1(0.3)	33.9(0.3)	36.2(0.3)	35.6(0.3)	36.9(0.3)	45.1(0.3)
Urbanicity								
Low	36.1(0.4)	36.6(0.3)	36.5(0.3)	36.0(0.3)	37.4(0.3)	37.2(0.3)	37.8(0.3)	46.1(0.3)
Medium	36.4(0.3)	37.4(0.3)	36.7(0.3)	35.5(0.3)	36.6(0.3)	37.2(0.3)	38.2(0.3)	46.6(0.3)
High	34.2(0.3)	36.2(0.3)	35.3(0.3)	34.6(0.3)	36.2(0.3)	38.6(0.3)	39.3(0.3)	47.6(0.3)

Abbreviations: CHNS, China Health and Nutrition Survey; tAHEI, tailored Alternative Healthy Eating Index.

^aValues are predicted margins (standard error) estimated by mix-effect linear random intercept models adjusted for baseline age (continuous), total energy intake (continuous), survey years(dummy variables), income, education, geographic region, urbanicity and corresponding interaction terms with survey years.