Published in final edited form as:

Addiction. 2017 May ; 112(5): 884–896. doi:10.1111/add.13734.

Neural responses to negative outcomes predict success in community-based substance use treatment

Sarah E. Forster, PhD^{1,2,3}, Peter R. Finn, PhD¹, and Joshua W. Brown, PhD¹

¹Indiana University, Department of Psychological and Brain Sciences

²VA Pittsburgh Healthcare System

³University of Pittsburgh, Department of Psychiatry

Abstract

Background and aims—Activation in some specific brain regions has demonstrated promise as prognostic indicators in substance dependent individuals (SDIs) but this issue has not yet been explored in SDIs attending typical of community-based treatment. We used a data-driven, exploratory approach to identify brain-based predictors of treatment outcome in a representative community sample of SDIs. The predictive utility of brain-based measures was evaluated against clinical indicators, cognitive-behavioral performance, and self-report assessments.

Design—Prospective clinical outcome design, evaluating baseline functional magnetic resonance imaging data from the Balloon Analogue Risk Task (BART) as a predictor of 3-month substance use treatment outcomes.

Setting—Community-based substance use programs in Bloomington, Indiana, USA.

Participants—Twenty-three SDIs (17 male, ages 18–43) in an intensive outpatient or residential treatment program; abstinent 1–4 weeks at baseline.

Measurements—Event-related brain response, BART performance, and self-report scores at treatment onset, substance use outcome measure (based on days of use)

Findings—Using voxel-level predictive modeling and leave-one-out cross-validation, an elevated response to unexpected negative feedback in bilateral amygdala and anterior hippocampus (Amyg/aHipp) at baseline successfully predicted greater substance use over the 3-month study interval ($p = 0.006$, cluster-corrected). This effect was robust to inclusion of significant non-brain-based covariates. A larger response to negative feedback in bilateral Amyg/aHipp was also associated with faster reward-seeking responses after negative feedback ($r(23) = -0.544$, $p = 0.007$; $r(23) = -0.588$, $p = 0.003$). A model including Amyg/aHipp activation, faster reward-seeking after

Corresponding author: Joshua W. Brown, PhD, Indiana University, 1101 E. 10th St., Bloomington, IN 47405, Tel.: (812) 855-9282; Fax: (812) 855-4691; jwmbrown@indiana.edu.

Competing Interests: None declared

Author Contributions: PRF, JWB, and SEF were responsible for the study design and concept. SEF performed all data preprocessing and analysis. SEF, JWB, and PRF contributed to the interpretation of results. SEF drafted the manuscript. SEF, JWB, and PRF participated in revision of the manuscript. All authors critically reviewed content and approved the final version for publication. SEF had full access to all the data in the study and takes responsibility for the integrity of data and the accuracy of the data analysis.

Conflict of Interest Disclosures: The authors have no conflicts of interest to disclose.

negative feedback, and significant self-report scores accounted for 45% of the variance in substance use outcomes in our sample.

Conclusions—An elevated response to unexpected negative feedback in bilateral amygdala and anterior hippocampus (Amyg/aHipp) appears to predict relapse to substance use in people attending community-based treatment.

The majority of substance dependent individuals (SDIs) resume use within two years of treatment (1, 2) and as few as 4% achieve continuous abstinence (3). Recovery from substance use disorders (SUDs) involves a complex interaction between intrinsic and extrinsic factors including motivation, social environment, and capacity for balanced decision-making – which vary across individuals and over time (4). As demographic, psychosocial, clinical, and cognitive-behavioral predictors have demonstrated limited utility (5), attention has increasingly turned to brain-based “neuromarkers” of relapse risk (6).

Alterations in brain function underlie compulsive use (e.g. incentive sensitization of drug cues) and failures of self-control (7) and these may serve as prognostic indicators in SDIs. For example, aberrant activation (e.g. hyperactivation of reward and sensory processing areas) to substance-related (8–15) and non-substance-related reward cues (16–20) has been associated with earlier and/or more likely relapse, as has impaired control-related processing during risky (21) or probabilistic decision-making (18, 19, 22) and other control-demanding tasks (13, 23–27).

In line with models of stress-induced relapse (28, 29), relapse risk also increases with ventromedial PFC/anterior cingulate cortex (vmPFC/ACC) hyperactivation during relaxation and hypoactivation during stress (30, 31), as well as reduced functional connectivity between vmPFC/ACC and amygdala, which may reflect impaired stress and affect regulation (32). Risk-related structural abnormalities within prefrontal control areas (8, 15, 33–36) and the amygdala (8, 15, 33, 34, 37), are also consistent with this account.

Relapse-related neuromarkers have contributed to enthusiasm for neuroimaging to enhance clinical assessment and inform personalized treatment (6). Several studies have demonstrated that brain-based measures outperform subjective self-reports and other traditional clinical indicators (12, 13, 21, 25, 26, 31, 32). However, as was recently identified by Whelan and Garavan (38), many studies fail to correct for statistical optimism when assessing brain-based predictors and may not generalize. This may be particularly problematic in studies of substance use relapse which have generally focused on a single SUD and excluded comorbid conditions, already limiting generalizability to real-world treatment settings.

SUDs commonly co-occur with internalizing psychopathology, which has also been associated with abnormal reward-processing and stress-reactivity (39, 40). Relapse prediction based on naturalistic samples (i.e., including comorbid SUD and internalizing disorders) may thus reflect transdiagnostic constructs relevant to chronicity. Dysfunction of negative valence systems underlying threat and stress-reactivity, for example, may increase vulnerability to internalizing disorders (40–42) and substance use compounds this effect (43). However, SUDs and internalizing comorbidities may also act in opposition; for

example, with the former decreasing and the latter increasing error signaling (44). It is therefore possible that neural predictors from SUD samples without comorbidity will not effectively translate for clinical use.

The current study represents a first step toward addressing this issue by exploring neural predictors of relapse in a heterogeneous sample of SDIs, typical of community-based treatment settings. We used a Balloon Analogue Risk Task (BART) that concurrently measures several predictive domains (i.e., risky decision-making, reward/negative outcome processing), uses monetary incentives rather than substance-specific stimuli, and has ecological validity (45). Fast event-related imaging further allowed differentiation of decision-making and outcome evaluation phases of the BART (46). Non-brain-based measures were also assessed, against which the predictive utility of brain-based measures could be evaluated. Despite including a mixed SUD sample with representative comorbidity, we hypothesized that previous findings of increased relapse risk with impaired error signaling in ACC (25) would replicate for negative outcomes in the BART and extend to risk-related signaling during decision-making. Due to our novel use of a community-based sample, we additionally conducted data-driven, exploratory analyses of corticolimbic circuitry to inform future work.

Methods

Design

We utilized a prospective clinical outcome design. Neuroimaging, cognitive-behavioral, and self-report measures were collected at the beginning of treatment and examined as candidate predictors of three-month substance use outcomes.

Participants

Twenty-six SDIs were enrolled upon engagement with community-based treatment and assessed within 1–4 weeks of self-reported abstinence; a sample of 23 (ages 18–43) was followed for 3 months (see Table 1 for demographic and recruitment data). Participants received treatment-as-usual through one of two abstinence-oriented, community-based addiction treatment programs (one intensive outpatient and one residential program; see Supporting Materials for additional information) and were not receiving replacement pharmacotherapy. All met DSM-IV criteria for alcohol, drug, or polysubstance dependence, without history of traumatic brain injury, neurocognitive disorders, Bipolar, or psychotic illness. An electronic breathalyzer and 6-panel urine drug screen were conducted at baseline and 3-month follow-up; behavioral signs of intoxication were also evaluated. Assessments were only conducted if blood alcohol content was 0.000 percent by volume, urinalysis was negative for illicit substances, and behavioral signs of intoxication were absent. Participants provided written informed consent; all methods were approved by the Indiana University Institutional Review Board.

fMRI Acquisition and BART Procedure

Imaging data were acquired at baseline and 3-month follow-up using a 32-channel head coil-equipped, Siemens Magnetom Trio 3-Tesla MRI scanner; follow-up data are reported

elsewhere (47). Echo-planar gradient-echo T2*-weighted sequences of 240 whole-brain volumes measured the functional blood-oxygen-level-dependent (BOLD) response during two eight-minute blocks of BART (see Figure 1). Acquisition, task, and preprocessing parameters were identical to previous work (46, 48); details provided in Supporting Information. In brief, the BART involved button-press responses to either ‘Inflate’ a balloon – incrementally and quasi-exponentially increasing its size and value in parallel with the probability of explosion – or ‘Redeem’ it for its current value. Stimuli were presented using E-Prime (Psychology Software Tools, Pittsburgh, PA); a projector was used for display within the scanner bore. Trials began with an image of a balloon. A red rectangle was presented for 1.5–2.5 seconds, indicating to wait before responding and subsequently turned green, indicating to respond when ready. Participants responded with left or right index fingers (response-mapping counterbalanced across subjects).

Each balloon sequence began with a balloon worth \$0.00. After each response, a jittered delay of 0–6 seconds preceded feedback (‘Successful_Gamble’, ‘Explode’, ‘Confirmed_Gain’) such that the BOLD response could be estimated separately for decision and outcome intervals. ‘Inflate’ responses could result in either 1) a ‘Successful_Gamble,’ whereupon the display was updated with a larger balloon and increased wager value, initiating the next decision or 2) ‘Explode’ feedback, indicating loss of the current wager. ‘Redeem’ responses always resulted in ‘Confirmed_Gain’ feedback and addition of the balloon value to block winnings (visible at bottom of screen). Participants completed an average of 95 decision trials and 16 balloon sequences per block.

Additional Measures

The timeline follow-back procedure was administered at 1-, 2-, and 3-month timepoints to document drug and alcohol use during the study interval. Narrative details of use (e.g., subjective intoxication) were also acquired. Details of the timeline follow-back procedure, as well as psychodiagnostic, self-report, and cognitive-behavioral assessments are provided in Supporting Information.

fMRI Analysis

Imaging data were analyzed using SPM5 and Matlab R2013a. BART BOLD responses were estimated using a general linear model with 20 regressors: six motion regressors, two constants, five main effect regressors (‘Redeem’, ‘Inflate’, ‘Explode’, ‘Successful_Gamble’, ‘Confirmed_Gain’), five parametric modulators representing explosion probability for each event, and 2 parametric modulators representing ‘Redeem’ and ‘Inflate’ response times (RTs). A canonical hemodynamic response function was used to model event-related signals during decision (i.e., RT) and outcome (i.e., feedback onset) intervals. Subject was included as a random effect at the second-level.

Contrasts isolated effects associated with negative outcomes over uncertain (‘Explode – Successful_Gamble’) and certain gains (‘Explode – Confirmed_Gain’), decisions to pursue uncertain gains (‘Inflate – Redeem’), and decisions to pursue certain gains (‘Redeem – Inflate’), as well as regions in which trial-to-trial fluctuations in BOLD signal correlated with the probability of explosion (i.e. parametric modulators, designated by ‘*p(Explode)’).

Using a cluster-forming threshold of $p < 0.001$, clusters of 30 or more voxels with a cluster-corrected p -value of < 0.05 , were identified as regions of interest (ROIs). The cluster-extent threshold was determined using AFNI's 3dClustSim to provide a type I error rate of $\alpha = 0.005$. Two participants were excluded from the second-level 'Explode*p(Explode)' contrast because low explosion frequency precluded unique specification in the first-level general linear model.

Whole-brain correlational analyses were conducted for each contrast to identify regions in which task-related activity was associated with substance use during the study interval. A substance use metric (SUM) was calculated from timeline follow-back data for each participant, representing incidents of drug and alcohol use during the study, weighted by presence/absence of intoxication;¹ a log-odds transformation was applied prior to analysis. This approach was chosen to better capture variability in substance use outcomes within the study sample (see Supporting Information). Next, voxel-level predictive modeling and leave-one-out cross-validation were used to identify ROIs predictive of substance use outcomes and compare performance of models based on brain- and/or non-brain-based predictors (see Supporting Information for details). Briefly, linear regression was applied with cross-validation on a voxel-by-voxel basis. Clusters wherein constituent voxels yielded predicted SUM values for left-out participants that strongly correlated with actual SUM values were targeted for further investigation. Leave-one-out cross-validation was selected for optimism-correction because this approach has been recommended for small neuroimaging datasets (49) and used in similar work with a comparable sample size (16). Predictive analysis of binary treatment outcomes was also conducted using Receiver Operating Characteristic curves and is described in Supporting Information and Figure S1.

Analysis of Cognitive-behavioral and Self-Report Measures

Pearson correlations with the substance use outcome variable (i.e., SUM) were calculated for each measure. Paired t-tests and repeated measures analysis of variance (ANOVA) were utilized for comparisons between conditions, as necessary. Statistics were computed in SPSS. Self-report and behavioral findings for cognitive-behavioral tasks other than BART are summarized in Supporting Information and Table S1.

Results

BART Performance

BART performance is summarized in Table 2. Consistent with successful performance, average winnings were \$16 per block, 'Confirmed_Gains' occurred more often than 'Explode' events, and the average number of inflations prior to 'Redeem'/'Confirmed_Gain' events was higher than that for 'Explode' outcomes. The number of completed balloons, proportion of balloons ending in 'Confirmed_Gain' vs. 'Explode,' and number of inflations did not significantly differ between blocks. There were no significant effects of block or response type on RT. A higher number of inflations prior to 'Redeem' responses was

¹(Total Days Monitored/4)*[Days Alcohol Used without Intoxication + 2*(Days Alcohol Used with Intoxication) + Days Drugs Used without Intoxication + 2*(Days Drugs Used with Intoxication)]

associated with increased substance use during the study interval; this effect was significant at $p = 0.046$, two-tailed, ($r(23) = 0.42$). Average RT did not significantly correlate with SUM ($r(23) = -0.23$, $p = 0.291$).

Neural Prediction of Relapse Risk

Basic event-related and parametric modulator contrasts are described in Supporting Information and Table S2a–c. Whole-brain correlational analyses between planned contrasts and SUM values are summarized in Table 3. Contrary to our hypotheses, an elevated response to negative feedback ('Explode – Confirmed_Gain') in right supplementary motor area (SMA), left dorsal posterior cingulate, and a region including right amygdala and anterior hippocampus (Amyg/aHipp) was associated with greater use – as was an increased BOLD response with increasing explosion probability in midline SMA during 'Inflate' events (i.e., $\text{Inflate} * p(\text{Explode})$). A similar effect was observed in left angular gyrus and putamen for 'Successful_Gamble * $p(\text{Explode})$ ' and in right insula and left SMA for 'Explode * $p(\text{Explode})$.' Greater 'Redeem – Inflate' activation in left IFG was associated with less use.

Given our modest sample size, correlational findings may not have external validity (see, 38). To address this, leave-one-out cross-validation was used to test outcome prediction by event-related brain signals on a voxel-by-voxel basis. Only two contrasts ('Explode – Confirmed_Gain' and 'Explode – Successful_Gamble') revealed regions of 30 or more voxels in which predicted outcomes for left-out participants strongly correlated with actual SUM values ($r = 0.70$) and survived cluster-level correction. For both contrasts, peak model performance (indicated by the strongest correlation between actual and predicted values) was identified in bilateral Amyg/aHipp (see Table 3, Figure 2). It is noted that the 'Explode – Successful_Gamble' right Amyg/aHipp cluster was 29 voxels, just below our conservative, *a priori* threshold.

To further investigate these significant clusters, spherical ROIs (radius: 1 voxel; volume: 7 voxels) were defined around voxels associated with peak model performance in bilateral Amyg/aHipp and leave-one-out cross-validation was repeated following addition of non-brain-based measures that significantly correlated with SUM (specifically, DOSPERT Expected Benefit scores), as well as a "null" model including only non-brain-based covariates (NM1). Mean Akaike information criterion (AIC) values for each model and ROI are summarized in Table 4. Model comparison was conducted separately for each ROI; multiple brain-based predictors were not included in the same model due to collinearity concerns. Left and right 'Explode – Successful_Gamble' ROIs and the right 'Explode – Confirmed_Gain' ROI passed the model comparison test – each demonstrating a reduction in AIC following addition of the voxel-based predictor. A second null model (NM2) added covariates with $p = 0.125$ (i.e., DOSPERT Risk Perception and Go/No-Go performance) and intuitive clinical indicators (i.e., craving scores, SUD symptom counts). Here, addition of the voxel-based predictor increased AIC (see Table 4), indicating that the brain-based measure was not supported with this expanded set of non-brain-based covariates.

Behavioral Correlates of Predictive Activation

Although not predicted in our original hypotheses, identification of Amyg/aHipp-based predictive signals may be consistent with the amygdala's putative role in stress-induced reinstatement of drug-seeking (28). If unexpected negative feedback (i.e. stress) increased reward-seeking in the current paradigm, it may be evident in faster inflation responses following 'Explode' versus 'Confirmed_Gain' events. To explore this possibility, the difference in initial inflation RT between post-Explode and post-Confirmed_Gain events was calculated for each participant ($M = -229$ ms, $SD = 225$) and significantly correlated with SUM, such that greater post-explosion speeding corresponded with increased use ($r(23) = -0.537$, $p = 0.008$). In addition, a significant negative correlation between BOLD signal change and post-explosion speeding (i.e. greater activation during 'Explode' events associated with faster post-explosion RTs) was identified in three of four Amyg/aHipp ROIs (see Figure 2).

Leave-one-out cross-validation was again repeated with post-explosion speeding added as a significant non-brain-based covariate (NM3). Resulting AICs were lower for the null model in all four ROIs (see Table 3) but approached zero for both ROIs in right Amyg/aHipp, suggesting that improved model performance with addition of the voxel-based predictor was nearly sufficient to justify increased model complexity. For right Amyg/aHipp ROIs, the full model accounted for 45.2% and 45% of variance in substance use outcomes for 'Explode – Confirmed_Gain' and 'Explode – Successful_Gamble' contrasts, respectively. Equations for these models are provided in Supporting Information.

Discussion

This study extends previous findings that brain-based measures outperform non-brain-based measures (e.g. clinical indicators, self-reports, etc.) in predicting substance use treatment outcomes by demonstrating this effect in a mixed sample of SDIs – generally representative of community-based treatment settings with respect to psychiatric and substance use comorbidities. While our initial hypotheses were not supported, voxel-by-voxel predictive modeling specifically revealed a bilateral effect in Amyg/aHipp for negative feedback contrasts, wherein an elevated Amyg/aHipp response was associated with increased use during early recovery. This effect corresponded with faster inflation responses following negative feedback, possibly reflecting increased reward-seeking in response to negative affect – as predicted by models of stress-induced relapse.

Importantly, consistency of observations between 'Explode – Confirmed_Gain' and 'Explode – Successful_Gamble' contrasts suggests the identified effect is associated with negative feedback processing and is not a consequence of differing feedback certainty in 'Explode' versus 'Confirmed_Gain' conditions. However, our findings require replication and effects of uncertainty versus negative valence should be directly examined in future work. Taken together, a predictive model including right Amyg/aHipp response to negative feedback (from either contrast), the behavioral post-explosion speeding effect, and DOSPERT Expected Benefits scores accounted for approximately 45% of variance in substance use outcomes in our sample. These results support the utility of multimodal

predictive models (including neural, behavioral, and self-report measures) to assess relapse vulnerability in community-based treatment settings.

Consistent with our findings, abnormality of the amygdala and hippocampus has been widely reported in SDIs. Reduced amygdalar and hippocampal volumes have been noted in SDIs (33, 34, 37), and smaller amygdalae have been associated with increased craving and relapse severity (34, 37). Hyperactivation of the amygdala and hippocampus in SDIs has also been observed in response to stress (31) and substance-related cues (31, 50) and increased resting cerebral blood flow to posterior hippocampus has been associated with greater relapse risk (51, 52). Amygdala activation has similarly been associated with poorer treatment outcomes (17), as well as increased cue-induced craving (50, 53–57). Importantly, the amygdala is integral to the aversive experience of monetary loss (58), as well as negative affect more generally (59), suggesting that a stronger experience of negative outcomes may contribute to greater substance use in the current study.

The amygdala has similarly been implicated in animal models of stress-induced relapse (28, 29) and in the “frustration effect” – wherein response speed increases following reward omission (60, 61). With respect to the latter, a similar effect (i.e. increased inflation speed following explosions) was significantly correlated with both substance use outcomes and activation of Amyg/aHipp ROIs in the current study. Behaviorally, this phenomenon is similar to “loss-chasing” in pathological gamblers (62, 63), which has also been linked to amygdala in animal models (64). Gambling disorder was not assessed herein but is commonly comorbid with SUDs and may share similar neurobiological substrates (65).

Amygdala has further been implicated in individual differences in risky reward-seeking by the triadic model of motivated behavior (66), wherein different neurodevelopmental trajectories of the striatal approach system, amygdala-based avoidance system, and prefrontal control system underlie cognitive-behavioral changes during adolescence. In line with predictions of this model, problematic alcohol use has been associated with a striatal-amygdalar imbalance – precipitating alcohol use disorder when relative amygdalar hyperactivity is potentiated by stress (67). Consistent with current findings, stress has also been shown to increase functional connectivity between the amygdala and dorsal striatum which may drive a shift toward faster, more impulsive responding (68). However, while greater dorsal striatal ‘Successful_Gamble*p(Explode)’ signal positively correlated with substance use in the current study, neither ventral nor dorsal striatal ROIs were identified by voxel-wise predictive modeling.

Taken together, the current results add to converging evidence in support of amygdala-based models of stress-induced relapse. Susceptibility to stress-induced relapse is state-dependent, making it challenging to characterize through conventional psychometric approaches. A reliable brain-based predictor of stress-induced relapse would have a significant translational impact but has not previously been described for a naturalistic sample wherein vulnerability to relapse is shaped, in part, by psychiatric comorbidity. Recent evidence suggests that amygdala reactivity predicts a fundamental vulnerability to life stress that may predispose individuals to internalizing disorders (40), as well as SUDs. In effect, previous exclusion of comorbid presentations may have precluded identification of amygdala-based

neuroprognostic indicators. However, by extension, the current findings may not generalize to those with unimorbid SUDs.

Importantly, neural predictors may also inform development of novel interventions and individualized treatment approaches (5, 69, 70). Antagonism of corticotropin-releasing factor (CRF) may dampen the stress response in the hypothalamic-pituitary-adrenal axis and amygdala and has been shown to attenuate stress-induced relapse in rats (71). Evidence-based cognitive-behavioral interventions may also prevent relapse by altering neural pathways associated with stress responsivity and negative affect (72). In addition, interventions that upregulate dlPFC function (e.g., cognitive reappraisal (73), repetitive transcranial magnetic stimulation (rTMS) (74), and transcranial direct current stimulation (75)) may improve regulation of amygdala reactivity (76). Indeed, rTMS over left dlPFC has been shown to reduce subjective craving and improve control of compulsive substance use (77).

To our knowledge, this is the first study to utilize cross-validation of voxel-wise predictive modeling with a continuous substance use outcome variable. This method uniquely identified ROIs in bilateral Amyg/aHipp associated with negative feedback. By comparison, whole-brain correlation analyses identified several additional predictive clusters and only a single ROI in right Amyg/aHipp for 'Explode – Confirmed_Gain.' Small volume correction for bilateral amygdala and hippocampus was sufficient to identify clusters approaching cluster-corrected significance in left Amyg/aHipp for both 'Explode – Confirmed_Gain' ($x = -34$, $y = -20$, $z = -18$; cluster size: 18; $t(21) = 4.30$, $p = 0.057$) and 'Explode – Successful_Gamble' ($x = -32$, $y = -4$, $z = -20$; cluster size: 26; $t(21) = 4.50$, $p = 0.046$) contrasts. However, this bilateral effect would have been overlooked if analyses were limited to whole-brain correlations.

The current study has several limitations. Our modest sample size precluded systematic investigation of sex differences (78–80) which may exist in neuropredictive signals (25). In addition, while we made efforts to identify generalizable predictive signals (e.g. monetary incentives rather than substance-related cues, leave-one-out cross-validation), we were unable to explore SUD subgroups or specific comorbidities. Moreover, we did not have a matched healthy control group, so we cannot conclude that predictive signals localized to Amyg/aHipp are elevated relative to individuals without SUDs, although this view is supported by existing evidence (31, 50).

The current study represents the first effort to identify neuropredictive indicators of relapse risk in a naturalistic sample of SDIs, typical of community-based treatment. We identified ROIs in bilateral Amyg/aHipp as significantly predictive of substance use outcomes, even when controlling for significant non-brain-based covariates. In addition, a novel behavioral correlate of relapse risk – faster reward-seeking responses after negative feedback – was also identified. Results of the current study may have translational relevance to the development of multimodal assessment tools and targeted interventions for individuals at the highest risk of relapse.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors would like to thank William Hetrick, PhD and Brian O'Donnell, PhD for helpful comments on the dissertation of which this work was a part.

Funding Support: This research was supported by NIH R01 grant DA026457 to JWB and NIH R01 grant AA13650 to PRF. SEF was partly supported by NIH TL1 award RR025759 (A. Shekhar, PI) during early stages of the project and by funding from the VISN 4 Mental Illness Research, Education and Clinical Center (MIRECC, Director: D. Oslin; Pittsburgh Site Director: G. Haas), VA Pittsburgh Healthcare System during manuscript preparation. The contents do not represent the views of the Department of Veterans Affairs, Department of Defense, or the United States Government. All financial support for the current work came from governmental or other public sources.

REFERENCES

1. Finney JW, Moos RH. The long-term course of treated alcoholism: II. Predictors and correlates of 10-year functioning and mortality. *Journal of studies on alcohol*. 1992; 53:142–153. [PubMed: 1313935]
2. Hubbard RL, Craddock SG, Anderson J. Overview of 5-year followup outcomes in the drug abuse treatment outcome studies (DATOS). *Journal of substance abuse treatment*. 2003; 25:125–134. [PubMed: 14670518]
3. Taylor C, Brown D, Duckitt A, Edwards G, Oppenheimer E, Sheehan M. Patterns of outcome: drinking histories over ten years among a group of alcoholics. *British journal of addiction*. 1985; 80:45–50. [PubMed: 3856447]
4. Prochaska JO, DiClemente CC. Stages and processes of self-change of smoking: toward an integrative model of change. *Journal of consulting and clinical psychology*. 1983; 51:390–395. [PubMed: 6863699]
5. Reske M, Paulus MP. Predicting treatment outcome in stimulant dependence. *Annals of the New York Academy of Sciences*. 2008; 1141:270–283. [PubMed: 18991963]
6. Gabrieli JD, Ghosh SS, Whitfield-Gabrieli S. Prediction as a humanitarian and pragmatic contribution from human cognitive neuroscience. *Neuron*. 2015; 85:11–26. [PubMed: 25569345]
7. Goldstein RZ, Volkow ND. Dysfunction of the prefrontal cortex in addiction: neuroimaging findings and clinical implications. *Nat Rev Neurosci*. 2011; 12:652–669. [PubMed: 22011681]
8. Beck A, Wustenberg T, Genauck A, Wrase J, Schlagenhauf F, Smolka MN, et al. Effect of brain structure, brain function, and brain connectivity on relapse in alcohol-dependent patients. *Arch Gen Psychiatry*. 2012; 69:842–852. [PubMed: 22868938]
9. Braus DF, Wrase J, Grusser S, Hermann D, Ruf M, Flor H, et al. Alcohol-associated stimuli activate the ventral striatum in abstinent alcoholics. *Journal of neural transmission (Vienna, Austria : 1996)*. 2001; 108:887–894.
10. Grusser SM, Wrase J, Klein S, Hermann D, Smolka MN, Ruf M, et al. Cue-induced activation of the striatum and medial prefrontal cortex is associated with subsequent relapse in abstinent alcoholics. *Psychopharmacology (Berl)*. 2004; 175:296–302. [PubMed: 15127179]
11. Kosten TR, Scanley BE, Tucker KA, Oliveto A, Prince C, Sinha R, et al. Cue-induced brain activity changes and relapse in cocaine-dependent patients. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology*. 2006; 31:644–650. [PubMed: 16123763]
12. Marhe R, Luijten M, van de Wetering BJ, Smits M, Franken IH. Individual differences in anterior cingulate activation associated with attentional bias predict cocaine use after treatment. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology*. 2013; 38:1085–1093. [PubMed: 23303067]

13. Prisciandaro JJ, Myrick H, Henderson S, McRae-Clark AL, Brady KT. Prospective associations between brain activation to cocaine and no-go cues and cocaine relapse. *Drug and alcohol dependence*. 2013; 131:44–49. [PubMed: 23683790]
14. Li Q, Li W, Wang H, Wang Y, Zhang Y, Zhu J, et al. Predicting subsequent relapse by drug-related cue-induced brain activation in heroin addiction: an event-related functional magnetic resonance imaging study. *Addiction biology*. 2015; 20:968–978. [PubMed: 25214465]
15. Seo S, Mohr J, Beck A, Wustenberg T, Heinz A, Obermayer K. Predicting the future relapse of alcohol-dependent patients from structural and functional brain images. *Addiction biology*. 2015; 20:1042–1055. [PubMed: 26435383]
16. Garbusow M, Schad DJ, Sebold M, Friedel E, Bernhardt N, Koch SP, et al. Pavlovian-to-instrumental transfer effects in the nucleus accumbens relate to relapse in alcohol dependence. *Addiction biology*. 2015
17. Jia Z, Worhunsky PD, Carroll KM, Rounsaville BJ, Stevens MC, Pearlson GD, et al. An initial study of neural responses to monetary incentives as related to treatment outcome in cocaine dependence. *Biol Psychiatry*. 2011; 70:553–560. [PubMed: 21704307]
18. Stewart JL, Connolly CG, May AC, Tapert SF, Wittmann M, Paulus MP. Cocaine dependent individuals with attenuated striatal activation during reinforcement learning are more susceptible to relapse. *Psychiatry Res*. 2014; 223:129–139. [PubMed: 24862388]
19. Stewart JL, Connolly CG, May AC, Tapert SF, Wittmann M, Paulus MP. Striatum and insula dysfunction during reinforcement learning differentiates abstinent and relapsed methamphetamine-dependent individuals. *Addiction*. 2014; 109:460–471. [PubMed: 24329936]
20. Gowin JL, Ball TM, Wittmann M, Tapert SF, Paulus MP. Individualized relapse prediction: Personality measures and striatal and insular activity during reward-processing robustly predict relapse. *Drug and alcohol dependence*. 2015; 152:93–101. [PubMed: 25977206]
21. Gowin JL, Harle KM, Stewart JL, Wittmann M, Tapert SF, Paulus MP. Attenuated insular processing during risk predicts relapse in early abstinent methamphetamine-dependent individuals. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology*. 2014; 39:1379–1387. [PubMed: 24317375]
22. Paulus MP, Tapert SF, Schuckit MA. Neural activation patterns of methamphetamine-dependent subjects during decision making predict relapse. *Arch Gen Psychiatry*. 2005; 62:761–768. [PubMed: 15997017]
23. Brewer JA, Worhunsky PD, Carroll KM, Rounsaville BJ, Potenza MN. Pretreatment brain activation during stroop task is associated with outcomes in cocaine-dependent patients. *Biol Psychiatry*. 2008; 64:998–1004. [PubMed: 18635157]
24. Kober H, DeVito EE, DeLeone CM, Carroll KM, Potenza MN. Cannabis abstinence during treatment and one-year follow-up: relationship to neural activity in men. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology*. 2014; 39:2288–2298. [PubMed: 24705568]
25. Luo X, Zhang S, Hu S, Bednarski SR, Erdman E, Farr OM, et al. Error processing and gender-shared and -specific neural predictors of relapse in cocaine dependence. *Brain : a journal of neurology*. 2013; 136:1231–1244. [PubMed: 23485852]
26. Marhe R, van de Wetering BJ, Franken IH. Error-related brain activity predicts cocaine use after treatment at 3-month follow-up. *Biol Psychiatry*. 2013; 73:782–788. [PubMed: 23374638]
27. Charlet K, Beck A, Jorde A, Wimmer L, Vollstadt-Klein S, Gallinat J, et al. Increased neural activity during high working memory load predicts low relapse risk in alcohol dependence. *Addiction biology*. 2014; 19:402–414. [PubMed: 24147643]
28. Koob GF, Le Moal M. Review. Neurobiological mechanisms for opponent motivational processes in addiction, *Philosophical transactions of the Royal Society of London Series B. Biological sciences*. 2008; 363:3113–3123. [PubMed: 18653439]
29. Peters J, Kalivas PW, Quirk GJ. Extinction circuits for fear and addiction overlap in prefrontal cortex. *Learning & memory (Cold Spring Harbor NY)*. 2009; 16:279–288.
30. Blaine SK, Seo D, Sinha R. Peripheral and prefrontal stress system markers and risk of relapse in alcoholism. *Addiction biology*. 2015

31. Seo D, Lacadie CM, Tuit K, Hong KI, Constable RT, Sinha R. Disrupted ventromedial prefrontal function, alcohol craving, and subsequent relapse risk. *JAMA psychiatry*. 2013; 70:727–739. [PubMed: 23636842]
32. McHugh MJ, Demers CH, Salmeron BJ, Devous MD, Stein EA, Adinoff B. Cortico-amygdala coupling as a marker of early relapse risk in cocaine-addicted individuals. *Frontiers in psychiatry*. 2014; 5:16. [PubMed: 24578695]
33. Cardenas VA, Durazzo TC, Gazdzinski S, Mon A, Studholme C, Meyerhoff DJ. Brain morphology at entry into treatment for alcohol dependence is related to relapse propensity. *Biol Psychiatry*. 2011; 70:561–567. [PubMed: 21601177]
34. Durazzo TC, Tosun D, Buckley S, Gazdzinski S, Mon A, Fryer SL, et al. Cortical thickness, surface area, and volume of the brain reward system in alcohol dependence: relationships to relapse and extended abstinence. *Alcoholism. clinical and experimental research*. 2011; 35:1187–1200.
35. Rando K, Hong KI, Bhagwagar Z, Li CS, Bergquist K, Guarnaccia J, et al. Association of frontal and posterior cortical gray matter volume with time to alcohol relapse: a prospective study. *Am J Psychiatry*. 2011; 168:183–192. [PubMed: 21078704]
36. Sorg SF, Taylor MJ, Alhassoon OM, Gongvatana A, Theilmann RJ, Frank LR, et al. Frontal white matter integrity predictors of adult alcohol treatment outcome. *Biol Psychiatry*. 2012; 71:262–268. [PubMed: 22047719]
37. Wrase J, Makris N, Braus DF, Mann K, Smolka MN, Kennedy DN, et al. Amygdala volume associated with alcohol abuse relapse and craving. *Am J Psychiatry*. 2008; 165:1179–1184. [PubMed: 18593776]
38. Whelan R, Garavan H. When optimism hurts: inflated predictions in psychiatric neuroimaging. *Biol Psychiatry*. 2014; 75:746–748. [PubMed: 23778288]
39. Dillon DG, Rosso IM, Pechtel P, Killgore WD, Rauch SL, Pizzagalli DA. Peril and pleasure: an rdoc-inspired examination of threat responses and reward processing in anxiety and depression. *Depression and anxiety*. 2014; 31:233–249. [PubMed: 24151118]
40. Swartz JR, Knodt AR, Radtke SR, Hariri AR. A neural biomarker of psychological vulnerability to future life stress. *Neuron*. 2015; 85:505–511. [PubMed: 25654256]
41. Kujawa A, Hajcak G, Danzig AP, Black SR, Bromet EJ, Carlson GA, et al. Neural Reactivity to Emotional Stimuli Prospectively Predicts the Impact of a Natural Disaster on Psychiatric Symptoms in Children. *Biol Psychiatry*. 2016; 80:381–389. [PubMed: 26526228]
42. Mattson WI, Hyde LW, Shaw DS, Forbes EE, Monk CS. Clinical neuroprediction: Amygdala reactivity predicts depressive symptoms 2 years later. *Social cognitive and affective neuroscience*. 2016; 11:892–898. [PubMed: 26865423]
43. Fosnocht AQ, Briand LA. Substance use modulates stress reactivity: Behavioral and physiological outcomes. *Physiol Behav*. 2016
44. Olvet DM, Hajcak G. The error-related negativity (ERN) and psychopathology: toward an endophenotype. *Clinical psychology review*. 2008; 28:1343–1354. [PubMed: 18694617]
45. Fernie G, Cole JC, Goudie AJ, Field M. Risk-taking but not response inhibition or delay discounting predict alcohol consumption in social drinkers. *Drug and alcohol dependence*. 2010; 112:54–61. [PubMed: 20580495]
46. Fukunaga R, Brown JW, Bogg T. Decision making in the Balloon Analogue Risk Task (BART): anterior cingulate cortex signals loss aversion but not the infrequency of risky choices. *Cognitive, affective & behavioral neuroscience*. 2012; 12:479–490.
47. Forster SE, Finn PR, Brown JW. A preliminary study of longitudinal neuroadaptation associated with recovery from addiction. *Drug and alcohol dependence*. 2016; 168:52–60. [PubMed: 27620345]
48. Bogg T, Fukunaga R, Finn PR, Brown JW. Cognitive control links alcohol use, trait disinhibition, and reduced cognitive capacity: Evidence for medial prefrontal cortex dysregulation during reward-seeking behavior. *Drug and alcohol dependence*. 2012; 122:112–118. [PubMed: 21992873]
49. Price CJ, Ramsden S, Hope TM, Friston KJ, Seghier ML. Predicting IQ change from brain structure: a cross-validation study. *Developmental cognitive neuroscience*. 2013; 5:172–184. [PubMed: 23567505]

50. Schneider F, Habel U, Wagner M, Franke P, Salloum JB, Shah NJ, et al. Subcortical correlates of craving in recently abstinent alcoholic patients. *Am J Psychiatry*. 2001; 158:1075–1083. [PubMed: 11431229]
51. Adinoff B, Gu H, Merrick C, McHugh M, Jeon-Slaughter H, Lu H, et al. Basal Hippocampal Activity and Its Functional Connectivity Predicts Cocaine Relapse. *Biol Psychiatry*. 2015; 78:496–504. [PubMed: 25749098]
52. Adinoff B, Harris TS, Gu H, Stein EA. Posterior hippocampal regional cerebral blood flow predicts abstinence: a replication study. *Addiction biology*. 2016
53. Kilts CD, Schweitzer JB, Quinn CK, Gross RE, Faber TL, Muhammad F, et al. Neural activity related to drug craving in cocaine addiction. *Arch Gen Psychiatry*. 2001; 58:334–341. [PubMed: 11296093]
54. Prisciandaro JJ, McRae-Clark AL, Myrick H, Henderson S, Brady KT. Brain activation to cocaine cues and motivation/treatment status. *Addiction biology*. 2014; 19:240–249. [PubMed: 22458561]
55. Childress AR, Mozley PD, McElgin W, Fitzgerald J, Reivich M, O'Brien CP. Limbic activation during cue-induced cocaine craving. *Am J Psychiatry*. 1999; 156:11–18. [PubMed: 9892292]
56. Wiers CE, Stelzel C, Gladwin TE, Park SQ, Pawelczack S, Gawron CK, et al. Effects of cognitive bias modification training on neural alcohol cue reactivity in alcohol dependence. *Am J Psychiatry*. 2015; 172:335–343. [PubMed: 25526597]
57. Grant S, London ED, Newlin DB, Villemagne VL, Liu X, Contoreggi C, et al. Activation of memory circuits during cue-elicited cocaine craving. *Proc Natl Acad Sci U S A*. 1996; 93:12040–12045. [PubMed: 8876259]
58. Bechara A, Damasio H, Damasio AR, Lee GP. Different contributions of the human amygdala and ventromedial prefrontal cortex to decision-making. *J Neurosci*. 1999; 19:5473–5481. [PubMed: 10377356]
59. Costafreda SG, Brammer MJ, David AS, Fu CH. Predictors of amygdala activation during the processing of emotional stimuli: a meta-analysis of 385 PET and fMRI studies. *Brain research reviews*. 2008; 58:57–70. [PubMed: 18076995]
60. Henke PG. Limbic lesions and the energizing, aversive, and inhibitory effects of non-reward in rats. *Canadian journal of psychology*. 1979; 33:133–140. [PubMed: 519543]
61. Henke PG, Maxwell D. Lesions in the amygdala and the frustration effect. *Physiol Behav*. 1973; 10:647–650. [PubMed: 4575404]
62. Dickerson M, Hinchey J, Fabre J. Chasing, arousal and sensation seeking in off-course gamblers. *British journal of addiction*. 1987; 82:673–680. [PubMed: 3475105]
63. Lesieur HR. The compulsive gambler's spiral of options and involvement. *Psychiatry*. 1979; 42:79–87. [PubMed: 760136]
64. Tremblay M, Cocker PJ, Hosking JG, Zeeb FD, Rogers RD, Winstanley CA. Dissociable effects of basolateral amygdala lesions on decision making biases in rats when loss or gain is emphasized, Cognitive. *affective & behavioral neuroscience*. 2014; 14:1184–1195.
65. Probst CC, van Eimeren T. The functional anatomy of impulse control disorders. *Current neurology and neuroscience reports*. 2013; 13:386. [PubMed: 23963609]
66. Ernst M. The triadic model perspective for the study of adolescent motivated behavior. *Brain Cogn*. 2014; 89:104–111. [PubMed: 24556507]
67. Nikolova YS, Knodt AR, Radtke SR, Hariri AR. Divergent responses of the amygdala and ventral striatum predict stress-related problem drinking in young adults: possible differential markers of affective and impulsive pathways of risk for alcohol use disorder. *Molecular psychiatry*. 2015
68. Vogel S, Klumpers F, Krugers HJ, Fang Z, Oplaat KT, Oitzl MS, et al. Blocking the mineralocorticoid receptor in humans prevents the stress-induced enhancement of centromedial amygdala connectivity with the dorsal striatum. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology*. 2015; 40:947–956. [PubMed: 25355243]
69. Reske M. What neuroimaging has and has not yet added to our understanding of addiction. *Addiction*. 2013; 108:1357–1359. [PubMed: 23574436]
70. Volkow ND, Baler RD. Brain imaging biomarkers to predict relapse in alcohol addiction. *JAMA psychiatry*. 2013; 70:661–663. [PubMed: 23636789]

71. Shaham Y, Erb S, Leung S, Buczek Y, Stewart J. CP-154,526, a selective, non-peptide antagonist of the corticotropin-releasing factor1 receptor attenuates stress-induced relapse to drug seeking in cocaine- and heroin-trained rats. *Psychopharmacology (Berl)*. 1998; 137:184–190. [PubMed: 9630005]
72. Witkiewitz K, Lustyk MK, Bowen S. Retraining the addicted brain: a review of hypothesized neurobiological mechanisms of mindfulness-based relapse prevention. *Psychology of addictive behaviors : journal of the Society of Psychologists in Addictive Behaviors*. 2013; 27:351–365. [PubMed: 22775773]
73. Buhle JT, Silvers JA, Wager TD, Lopez R, Onyemekwu C, Kober H, et al. Cognitive reappraisal of emotion: a meta-analysis of human neuroimaging studies. *Cereb Cortex*. 2014; 24:2981–2990. [PubMed: 23765157]
74. Baeken C, De Raedt R, Van Schuerbeek P, Vanderhasselt MA, De Mey J, Bossuyt A, et al. Right prefrontal HF-rTMS attenuates right amygdala processing of negatively valenced emotional stimuli in healthy females. *Behavioural brain research*. 2010; 214:450–455. [PubMed: 20600336]
75. Andrews SC, Hoy KE, Enticott PG, Daskalakis ZJ, Fitzgerald PB. Improving working memory: the effect of combining cognitive activity and anodal transcranial direct current stimulation to the left dorsolateral prefrontal cortex. *Brain stimulation*. 2011; 4:84–89. [PubMed: 21511208]
76. Ochsner KN, Silvers JA, Buhle JT. Functional imaging studies of emotion regulation: a synthetic review and evolving model of the cognitive control of emotion. *Annals of the New York Academy of Sciences*. 2012; 1251:E1–E24. [PubMed: 23025352]
77. Protasio MI, da Silva JP, Arias-Carrion O, Nardi AE, Machado S, Cruz MS. Repetitive Transcranial Magnetic Stimulation to Treat Substance Use Disorders and Compulsive Behavior. *CNS & neurological disorders drug targets*. 2015; 14:331–340. [PubMed: 25801835]
78. Fox HC, Garcia M Jr, Kemp K, Milivojevic V, Kreek MJ, Sinha R. Gender differences in cardiovascular and corticoadrenal response to stress and drug cues in cocaine dependent individuals. *Psychopharmacology (Berl)*. 2006; 185:348–357. [PubMed: 16514523]
79. Li CS, Kosten TR, Sinha R. Sex differences in brain activation during stress imagery in abstinent cocaine users: a functional magnetic resonance imaging study. *Biol Psychiatry*. 2005; 57:487–494. [PubMed: 15737663]
80. Kilts CD, Gross RE, Ely TD, Drexler KP. The neural correlates of cue-induced craving in cocaine-dependent women. *Am J Psychiatry*. 2004; 161:233–241. [PubMed: 14754771]

Figure 1.
Schematic representation of the Balloon Analogue Risk Task (BART), reproduced, with permission, from Fukunaga et al., 2012.

Figure 2.

Correlation between the BOLD response to negative feedback in predictive regions identified through leave-one-out cross-validation analysis and the post-explosion speeding effect. ROIs identified in the leave-one-out cross-validation analysis for both contrasts include voxels in left and right amygdala and anterior hippocampus. Scatterplots depict mean BOLD response within spherical ROIs (radius = 1 voxel) defined around the maximally predictive voxel for each contrast (38, -16, -16 for 'Explode - Confirmed_Gain' and -34, -4, -22 for 'Explode - Successful_Gamble') versus post-explosion speeding (i.e. Post-Explosion RT - Post-Confirmed_Gain RT). Mean BOLD responses in left and right Amyg/aHipp ROIs for the 'Explode - Confirmed_Gain' contrast significantly correlated with post-explosion speeding ($r(23) = -0.544$, $p = 0.007$ and $r(23) = -0.588$, $p = 0.003$, respectively), indicating that a stronger response to negative feedback was associated with a greater reduction in post-explosion RT. The left ROI for the 'Explode - Successful_Gamble' contrast also significantly correlated with post-explosion speeding ($r(23) = -0.496$, $p = 0.016$) but the right ROI did not ($r(23) = -0.345$, $p = 0.107$).

Table 1

Summary of demographic, recruitment, and clinical characteristics of study sample (n = 23).

<u>Demographic Variables</u> [†]		<u>Mean (SD) or Count</u>
Age		27 (6.5)
Male		17
Total Years of Education		13 (1.8)
Employment Status	<i>Employed Full Time</i>	6
	<i>Employed Part Time</i>	6
	<i>Unemployed</i>	9
	<i>Student</i>	2
<u>Recruitment Information</u>		<u>Count</u>
Treatment Setting	<i>Outpatient</i>	14
	<i>Residential</i>	9
Referral Source	<i>Self</i>	13
	<i>Drug Court</i>	5
	<i>Probation</i>	5
<u>Clinical Variables</u>		<u>Mean (SD) or Count</u>
Times in Previous Treatment		2 (2.4)
Primary DSM-IV Diagnosis	<i>Alcohol dependence</i>	13
	<i>Sedative, Hypnotic, or Anxiolytic Dependence</i>	1
	<i>Cannabis dependence</i>	1
	<i>Amphetamine dependence</i>	1
	<i>Opioid dependence</i>	3
	<i>Polysubstance dependence</i>	4
Additional Psychiatric Diagnoses	<i>Major Depressive Disorder</i>	4
	<i>Dysthymic Disorder</i>	1
	<i>Generalized Anxiety Disorder</i>	4
	<i>Posttraumatic Stress Disorder</i>	1
	<i>Anxiety Disorder NOS</i>	2
	<i>Antisocial Personality Disorder</i>	1
	<i>None</i>	7
<u>Substance Use Outcomes</u>		<u>Mean (SD)</u>
	<i>Substance Use Metric</i>	0.64 (0.12)
	<i>% Days of Use</i>	12.8 (24.2)
	<i>% Days Use to Intoxication</i>	9.6 (16.0)

[†]The Substance Use Metric (SUM) did not significantly correlate with demographic factors, age ($r(23) = -0.171, p = 0.435$) and years of education ($r(23) = -0.332, p = 0.122$), and did not significantly differ with respect to sex ($\chi^2(2) = 2.08, p = 0.222$) or employment status ($F(3,19) = 0.67, p = 0.581$).

Summary of Balloon Analog Risk Task (BART) behavioral performance measures (n = 23).

Table 2

	Confirmed_Gain	Explode	All	Confirmed_Gain vs. Explode t-value (p-value)
# of Balloons	21 (5.4)	11.7 (4)	32.7 (3.1)	4.78 (< 0.001)
Inflation Count (#)	Mean (SD) Range Within Subject SD	5.2 (0.6) 2-9 1.3	5.9 (0.5) 2-10 1.3	9.77 (< 0.001) n/a n/a
	Redeem	Inflate	All	F-value (p-value)
Response Time (ms)	Mean (SD) 1,018 (732)	1,044 (445)	1,037 (463)	n/a
			Main Effect: Block	1.55 (0.226)
			Main Effect: Response Type	0.08 (0.782)
			Block × Response Type Interaction	1.90 (0.182)

Table 3

Regions of interest from whole-brain correlation analysis of substance use outcomes and leave-one-out cross-validation analysis.

Negative Correlation: Greater Activation Associated with Less Use					Peak MNI Coordinates		
Contrast	Region	Size	x	y	z	T-value	p-value (cluster-corrected)
Redeem – Inflate	L Inferior Frontal Gyrus (BA 45/47)	105	-46	24	8	5.23	0.019
Explode – Successful_Gamble	n/a	-	-	-	-	-	-
Explode – Confirmed_Gain	n/a	-	-	-	-	-	-
Redeem*p(Explode)	n/a	-	-	-	-	-	-
Inflate*p(Explode)	n/a	-	-	-	-	-	-
Explode*p(Explode)	n/a	-	-	-	-	-	-
Successful_Gamble*p(Explode)	n/a	-	-	-	-	-	-
Confirmed_Gain*p(Explode)	n/a	-	-	-	-	-	-
Positive Correlation: Greater Activation Associated with More Use					Peak MNI Coordinates		
Contrast	Region	Size	x	y	z	T-value	p-value (cluster-corrected)
Redeem – Inflate	n/a	-	-	-	-	-	-
Explode – Successful_Gamble	n/a	-	-	-	-	-	-
Explode – Confirmed_Gain	R Supplementary Motor Area (BA 6)	222	14	-26	62	5.55	< 0.001
	R Amygdala/Anterior Hippocampus	341	34	-22	-16	5.54	< 0.001
	L Dorsal Posterior Cingulate (subgyral)	148	-24	-30	46	5.37	0.001
Redeem*p(Explode)	n/a	-	-	-	-	-	-
Inflate*p(Explode)	L/R Supplementary Motor Area (BA 6)	132	-8	0	68	5.61	0.003
Explode*p(Explode)	R Insula (BA 13)	56	36	4	12	4.41	0.042
	L Supplementary Motor Area (BA 6)	74	-12	-12	60	3.91	0.010
Successful_Gamble*p(Explode)	L Angular Gyrus (BA 39)	149	-40	-82	30	5.90	0.001
	L Putamen	83	-26	10	-4	3.78	0.028
Confirmed_Gain*p(Explode)	n/a	-	-	-	-	-	-
Leave-One-Out Cross-Validation Analysis					Peak MNI Coordinates		

Negative Correlation: Greater Activation Associated with Less Use									
Peak MNI Coordinates									
Contrast	Region	Size	x	y	z	T-value	p-value (cluster-corrected)		
		Size	x	y	z	Mean T-value	p-value (cluster-corrected)		
Explode – Successful_Gamble	L Amygdala/Anterior Hippocampus	38	-34	-4	-22	0.77	< 0.001		
	R Amygdala/Anterior Hippocampus	29	34	-20	-16	0.78	0.006		
Explode – Confirmed_Gain	L Amygdala/Anterior Hippocampus	38	-34	-8	-26	0.80	< 0.001		
	R Amygdala/Anterior Hippocampus	33	38	-16	-16	0.80	0.002		

[‡]Reported *r*-values reflect the mean of voxel-level correlation coefficients for each cluster, representing the relationship between actual and predicted SUM values for the left-out participant. Correlation coefficients are provided as descriptive rather than inferential statistics and corresponding *p*-values are omitted. Reported *p*-values reflect whole-brain cluster-level correction to account for multiple comparisons.

Summary of predictive model performance in peak ROIs identified through leave-one-out cross-validation analysis (n = 23)

Table 4

Voxel Model Comparison	Contrast	Peak MNI Coordinates	Null Model		Voxel Only Model			Voxel Model with Covariates		
			AIC	R ²	Peak Voxel	ROI Mean	Mean R ²	Mean AIC	Mean Diff [†]	Mean R ^{2†}
NM1: Model with Significant Covariates	Explode – Confirmed_Gain	-34 -8 -26	44.09	0.141	0.93	0.81	0.116	46.78	-2.69	0.113
	Explode – Confirmed_Gain	38 -16 -16	44.09	0.141	0.96	0.80	0.226	42.13	1.96	0.306
	Explode – Successful_Gamble	-34 -4 -22	44.09	0.141	0.92	0.75	0.287	40.59	3.50	0.370
	Explode – Successful_Gamble	34 -20 -16	44.09	0.141	0.88	0.64	0.368	41.11	2.98	0.349
NM2: Model with Significant & Trending Covariates + Clinical Indices	Explode – Confirmed_Gain	-34 -8 -26	52.04	0.144	0.93	0.81	0.116	58.50	-6.46	-0.042
	Explode – Confirmed_Gain	38 -16 -16	52.04	0.144	0.96	0.80	0.226	56.73	-4.69	0.032
	Explode – Successful_Gamble	-34 -4 -22	52.04	0.144	0.92	0.75	0.287	54.63	-2.59	0.119
	Explode – Successful_Gamble	34 -20 -16	52.04	0.144	0.88	0.64	0.368	53.57	-1.53	0.163
NM3: Model with Significant Covariates + Post-Explosion Speeding	Explode – Confirmed_Gain	-34 -8 -26	40.10	0.390	0.93	0.81	0.116	43.68	-3.58	0.325
	Explode – Confirmed_Gain	38 -16 -16	40.10	0.390	0.96	0.80	0.226	40.63	-0.53	0.452
	Explode – Successful_Gamble	-34 -4 -22	40.10	0.390	0.92	0.75	0.287	41.21	-1.11	0.427
	Explode – Successful_Gamble	34 -20 -16	40.10	0.390	0.88	0.64	0.368	40.67	-0.57	0.450

[†] Reported *r*-values represent the correlation between actual and predicted SUM values for the left-out participant.

^{††} A positive mean AIC diff indicates that the addition of a voxel-based predictor was preferred despite constraints to limit model complexity.

^{†††} $R^2 = 1 - (\text{SSEResidual} / \text{SSETotal})$, where SSEResidual reflects the sum of squared differences between actual and predicted values for each left-out participant and SSETotal reflects the sum of squared differences between actual values for each left-out participant and the average of all actual values. Given our leave-one-out cross validation approach, a reduced (or negative) R^2 for the full model reflects overfitting on the basis of training data, which can worsen performance on the independent test set.