CORRESPONDENCE

The Risk of Bacterial Infection After Tattooing

by Dr. rer. nat. Ralf Dieckmann, Ph. D. Ides Boone, Stefan O. Brockmann, Dr. rer. nat. Jens A. Hammerl, Prof. Dr. med. Dr. rer. nat. Annette Kolb-Mäurer, Prof. Dr. med. Matthias Goebeler, Prof. Dr. med. Dr. rer. nat. Dr. med. habil. Andreas Luch, and Prof. Dr. med. M.Sc. Sascha Al Dahouk in issue 40/2016

Additional Risks

We thank Dieckmann and colleagues for their important—and long overdue—article (1). We agree with their conclusion, that people should probably refrain from having tattoos altogether, in order to avoid health risks but also to avoid experiencing socioeconomic problems. We have been issuing warnings of the potential risks associated with tattoos for many years (2).

Further to the more obvious risks of tattooing, such as infections and allergies, other medically relevant aspects of the procedures are almost unknown. Delayed complications that potentially threaten eyesight have included iris bombe, retinal detachment, and macular edema (3). Tattoos can also be medically relevant in the context of diagnostic measures. In MRI investigations, for example, the iron oxides in the tattoo inks may heat up and cause burns. Furthermore, there is a chance of examination artefacts.

Tattoos should also be regarded critically in the socioeconomic context. Professional starter and promotion opportunities may be negatively affected. Vacancies in many lines of work are preferentially given to persons without body modifications, rather than to applicants with tattoos or piercings (4).

Tattoo removal is also problematic. Laser treatment is associated with breakdown and decomposition products that can be toxic or carcinogenic. In our practice we therefore prefer (serial) excision of the affected areas if this is justifiable and practicable from a functional and esthetic perspective.

DOI: 10.3238/arztebl.2017.0177a

REFERENCES

1. Dieckmann R, Boone I, Brockmann SO, Hammerl JA, Kolb-Mäurer A, Goebeler M, Luch A, Al Dahouk S: The risk of bacterial infection after tattooing—a systematic review of the literature. *Dtsch Arztebl Int* 2016; 113: 665–71.
2. Sattler T, Sattler D: Tätowierungen: Getting inked – oder doch lieber Abziehbildchen?! DOI: 10.3205/15dgpae139.
3. Ostheimer TA, Burkholder BM, Leung TG, Butler NJ, Dunn JP, Thorne JE: Tattoo-associated uveitis. *Am J Ophthalmol* 2014; 158: 637–43.
4. Swanger N: Visible body modification (VBM): Evidence from human resource managers and recruiters and the effects on employment. *Hospitality Management*. 2006; 25: 154–8.

Dr. med. Thorsten Sattler
Dr. med. Dalia Sattler
Hamburg
info@klosterstern-praxis.de

Conflict of interest statement

The authors declare that no conflict of interest exists.

In Reply:

We thank Sattler and Sattler for their response to our article on the risk of bacterial infection after tattooing (1). The popularity of tattoos shows that in the meantime, this kind of body modification as an expression of individuality, lifestyle and freedom has made its way into the heart of Western societies. Ultimately, the decision in favor of or against a tattoo is a matter of personal choice. However, in terms of preventive consumer protection, clients should be comprehensively informed about the health risks associated with tattoos, in order to enable them to give informed consent. The responsibility for adequate information provision lies primarily with the tattooist who performs the procedure; but we as medical doctors are also under obligation to our patients and the general public to point out the increased risk of infections and allergic reactions, especially in patients with a relevant predisposition. To further minimize the number of complications resulting from tattoos, supervisory and control authorities will have to increasingly deal with health relevant issues. These include, for example, training requirements for tattooists, the sterility of tattoo inks, and the adherence to hygiene measures in the tattoo studio. The European standard for hygiene requirements for tattooing, which is currently in preparation, aims to create the basis for quality assurance (Comité Européen de Normalisation (CEN)/ Technical Committee 435). We are hoping that these standards for tattooing practice will result in greater safety for tattooists and those being tattooed.

DOI: 10.3238/arztebl.2017.0177b

REFERENCES

1. Dieckmann R, Boone I, Brockmann SO, Hammerl JA, Kolb-Mäurer A, Goebeler M, Luch A, Al Dahouk S: The risk of bacterial infection after tattooing—a systematic review of the literature. *Dtsch Arztebl Int* 2016; 113: 665–71.

On behalf of the authors

Prof. Dr. med. Sascha Al Dahouk

German Federal Institute for Risk Assessment (BfR), Berlin
Sascha.Al-Dahouk@bfr.bund.de

Conflict of interest statement

Prof. Al Dahouk has written a medical expertise report regarding this paper's subject matter.