Preparing Older Adults with Serious Illness To Formulate Their Goals for Medical Care in the Emergency Department

Kei Ouchi, MD, MPH,¹⁻³ Vinicius Knabben, BS,⁴ Laura Rivera-Reyes, MPH,⁴ Niharika Ganta, MD, MPH,⁵
Laura P. Gelfman, MD, MPH,⁶ Rebecca Sudore, MD,⁷ Ula Hwang, MD, MPH^{4,6}
on behalf of the GEDI WISE Investigators

Abstract

Background: Emergency department (ED) clinicians often lack training and resources to conduct advance care planning (ACP) conversations. The use of technology for health education is increasing, yet little is known if it can be used to engage older ED patients in ACP.

Objective: To determine the feasibility of using tablets to provide ACP education (www.prepareforyourcare.org)(PREPARE) to older ED patients.

Design: A feasibility study conducted in late 2014 and early 2015.

Setting/Subjects: Subjects were recruited from a parent cohort of older adults enrolled in a survey about Geriatric ED care. Inclusion criteria were ≥ 65 years age and English speaking; exclusions were hearing or vision impairment or if clinically unstable.

Measurements: Primary outcome was completion of ≥ 1 of 5 PREPARE modules. Secondary outcomes were ease of use (10-point scale; 1 = very hard, 10 = very easy) and the reasons for refusal to participate.

Results: Sixty-one subjects were approached; 24 (39%) were interested in viewing PREPARE after the Geriatric ED survey. Mean age was 75 years (standard deviation [SD] 9); 67% were female and 54% were nonwhite. Seventy-one percent of participants completed ≥ 1 module. Participants rated the website as easy to use for themselves (mean 8.4, SD 2.39) and for others (mean 7.3, SD 2.31). Of the subjects who declined, top reasons cited were fatigue (26%), already feeling prepared (13%), and technology limitations (11%).

Conclusion: PREPARE has the potential to engage older adults who are not acutely ill in ACP during their ED visits. Further studies should explore optimal approaches for ED implementation.

Keywords: advance care planning; emergency department; educational tool; geriatrics

Introduction

ADVANCE CARE PLANNING (ACP) is the process of planning for future medical decisions¹ with the goal of aligning the care patients to receive their preferences.² ACP has been shown to be associated with increased patient and family satisfaction, reduced anxiety, depression,

stress among surrogate decisions makers, and healthcare utilization,¹⁻³ as well as care consistent with patients' wishes.^{4,5}

The framework of ACP has expanded beyond code status, specific treatments,⁶ and completion of statutory documents (i.e., living will) covering hypothetical situations (i.e., advance directive [AD])⁷ to communication-based programs

¹Department of Emergency Medicine, Brigham and Women's Hospital, Boston, Massachusetts.

²Department of Emergency Medicine, Harvard Medical School, Boston, Massachusetts.

³Serious Illness Care Program, Ariadne Labs, Boston, Massachusetts.

⁴Department of Emergency Medicine, Icahn School of Medicine at Mount Sinai, New York, New York.

⁵Department of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania.

⁶Department of Geriatrics and Palliative Medicine, Icahn School of Medicine at Mount Sinai, New York, New York.

⁷Department of Medicine, University of California, San Francisco, San Francisco, California.

Accepted September 12, 2016.

facilitating discussion between patient and surrogate regarding goals, values, and preferences for medical care.⁸ To operationalize this expanded paradigm of ACP, PREPARE was developed (www.prepareforyourcare.org)⁹ and has been shown in a pilot study to engage a diverse set of older adults at senior centers in ACP.⁹ PREPARE has not been tested in other environments, such as the emergency department (ED).

The ED is an increasingly important setting to introduce ACP. ACP conversations occur infrequently in outpatient settings¹⁰ and many patients use the ED for primary care¹¹ often resulting in patients presenting to the ED who have not had the opportunity to identify and document their goals for care. Furthermore, 75% of older adults visit the ED in the last six months of their lives,¹² yet the majority (56%–99%) of older adults in the ED do not have documented goals for care available in the medical record.¹³ Such patients may benefit from revisiting ACP.¹⁴ Even if ACP occurred before ED arrival, it is rarely recorded in the medical records¹⁵ and patients' values and goals may change based on changing health states necessitating ED physicians revisiting patients' goals.¹⁶ Studies show that ED clinicians worry about providing value-concordant care,¹⁷ but also do not feel adequately trained to discuss goals of care with patients.^{17,18}

To help introduce ACP in the ED setting, a patient-centered multimedia ACP engagement tool, such as PREPARE, may be ideal. Patient-oriented decision aids have been used in the ED for a variety of conditions to facilitate shared decision making.¹⁹ However, a multimedia decision aid for ACP in the ED has never been studied. The aim of this feasibility study was to determine the proportion of adults ≥65 years in the ED who would be interested in using PREPARE on a tablet computer.

Methods

Subjects and setting

We conducted a pilot feasibility study at an academic ED in New York City as part of an innovative geriatric emergency care model called GEDI WISE (Geriatric Emergency Department Innovations in Care through Workforce, Informatics, and Structural Enhancements).²⁰ Participants eligible for this substudy were a convenience sample of the prospective GEDI WISE study. The GEDI WISE study was a convenience sample of English speakers, ≥65 years in age, who were cared for in the ED. Subjects were excluded if judged by the research assistant (RA) to have hearing or vision impairment or by the treating clinician to be clinically unstable (e.g., acute MI). After completing the GEDI WISE survey (median duration of 48 minutes), the RA subsequently invited subjects to participate in the ED PREPARE substudy. Data were collected for 2 one-month periods (October to November 2014 and January to February 2015) (Fig. 1). These surveys were approved by the institutional review boards and all subjects provided written informed consent.

Baseline assessment

Baseline demographic data included age, gender, race/ethnicity, emergency severity index (ESI),²¹ and level of education.

Procedures

Once the GEDI WISE study was completed, our study team approached a subset of individuals to view PREPARE and the study was described using a scripted text. If subjects expressed interest, the RA presented PREPARE on a tablet

FIG. 1. Study enrollment process.

computer. The RA instructed subjects to choose a module most pertinent to their situation and encouraged them to view as many modules as desired. Subjects then viewed modules independently. The RA returned in 20-minute intervals to ascertain module completion or if patients required further assistance. Subjects who declined to participate were asked about reason(s) for declining.

PREPARE

PREPARE consists of five interactive modules with videos that model ACP behaviors: (1) choosing a medical decision maker, (2) deciding what matters most in life, (3) deciding on leeway for surrogate decision makers, (4) communicating wishes with others, and (5) asking doctors questions to make informed decisions.⁹ PREPARE is designed to be easy to use with large font, text written at a fifth grade reading level, and closed captioning of all videos. Prior study showed a mean of 57 ± 16 minutes to complete all five modules or ~ 10 minutes for each module.⁹

Measures and analysis

Primary outcomes included the proportions of subjects who expressed interest in PREPARE and the proportion able to complete ≥ 1 modules. Secondary outcomes were ease of use on a 10 point scale, number of modules completed, and reasons for refusal to participate. For analysis, ANOVA and *t*-test were completed for continuous outcomes and chi-square for binary outcomes. A *p*-value of <0.05 was used as the threshold for statistical significance.

Results

During the study period, a total of 223 subjects were enrolled in the primary GEDI WISE study. Of these subjects, 61 were approached about PREPARE (Fig. 1). The mean age of the participants was 75 years (standard deviation [SD] 9), 67% were female, 54% were nonwhite, and 79% reported having more than high school education. There were no statistical differences between groups interested and not interested in PREPARE (Table 1).

Twenty-four subjects (39%) expressed interest in PREPARE. All interested participants engaged in at least some PREPARE content while in the ED, and 17 (71%) participants completed ≥ 1 module. The breakdown of modules watched were as follows: module 1 (41%), module 2 (41%), module 3 (18%), module 4 (18%), and module 5 (29%). The users found the website easy to use for themselves (mean score 8.4 out of 10, SD 2.39) and thought to be so for others (mean score 7.3, SD 2.31).

Thirty-seven (61%) participants declined to view PREPARE. The reported reasons included feeling fatigued after the 45-minute GEDI WISE survey (26%); already feeling prepared about ACP (13%); having technology limitations, such as difficulty using tablets in general (11%); and their current clinical situation (11%), such as experiencing anxiety while awaiting test results (e.g., a cancer patient awaiting CT scan results).

Discussion

Findings from this study indicate that it is feasible to engage older ED patients in ACP with a tablet-based version of

TABLE 1. DEMOGRAPHICS

Characteristics	Parent geriatric ED study (n = 223)	PREPARE (n = 24)	Refused (n = 37)	p
Age, mean (SD), years	76 (± 8.5)	74 (± 9)	77 (± 10)	0.42
No. of women (%)	125 (54.8)	16 (67)	20 (53)	0.78
Race/ethnicity (%)				
White	47	46	47	0.98
Black	30	28	32	0.97
Hispanic/Latino	10	11	11	0.94
Missing/other	11	11	11	0.98
ESI, mean (SD) (1 = acute, 5 = nonacute)	2.7 (0.6)	2.6 (0.5)	2.6 (0.5)	0.49
$\geq$ High school education	79	75	81	0.97

ESI, emergency severity index; SD, standard deviation.

PREPARE.⁹ Even after completing a lengthy Geriatric ED survey for another study, nearly 40% of 61 older adults were interested and willing to view PREPARE. This rate is similar to AD completion rates (18–36%) in the outpatient settings.²² Subjects also rated PREPARE as easy to use for themselves and for other patients.

Our findings are complimentary to prior studies and may help shape future strategies to engage older adults in ACP and health education technology in the ED. First, it is important to identify appropriate ED patients to engage in ACP. Participation may be challenging for patients who are fatigued from clinical and research activities while in the ED. Recruitment may have been higher if this were not a substudy conducted after the lengthy parent study. Second, older adults found a technology-based intervention in the ED easy to use. The result was consistent with recent findings of increasing internet use among older adults,²³ familiarity with tablet computers,²⁴ and high prevalence of mobile technology in the ED.²⁵ Third, the participation rate was higher than expected despite the fatigue from the primary GEDI WISE survey without clinician involvement. Well-established ED interventions for alcohol abuse suggest the importance of, at minimum, some clinician involvement (≤ 10 minutes) and motivational interviewing techniques, which may improve participation rate.²⁶ Such interventions may serve as a model to make it easier for busy EM clinicians to introduce ACP at the important time for the patient's illness trajectory, which may improve prognostic awareness and prepare patients for ACP conversation after leaving the ED. Fourth, prior studies suggest that an algorithm to tailor the intervention for specific stages of patient readiness²⁷ may optimally target the intervention to the most appropriate population. Finally, after engagement with PREPARE, it is important to consider ways to encourage patients to document their wishes and to discuss ACP with their primary outpatient providers. With increasing informational technologies for ACP,²⁸ seamless communication and documentation may be possible when patients are prepared to discuss ACP.

Limitations

Our study had several important limitations. First, our subjects were participants of an already lengthy geriatric ED survey. Because subjects completed the primary survey immediately before our study, it is likely that this added to fatigue and unwillingness to participate in our substudy. Second, the convenience sampling with only English speakers limits the generalizability of our findings. Third, we did not ask whether our participants had prior AD or pre-existing serious illness. Some patients declined participating in our study (13%) as they already felt prepared about ACP, and prior AD or absence of preexisting serious illness may have been one such reason. Targeted subject selection, based on prior engagement in ACP and preexisting serious illness, may be critical factors to achieve targeted participation for future studies. Finally, we did not record the time participants spent using PREPARE or why not all modules were completed in participants who completed ≥ 1 module, which are important considerations for feasibility.

Conclusion

With appropriate implementation, PREPARE has the potential to engage older adults who are not acutely ill in ACP during their ED visits. Further studies should explore optimal approaches for ED implementation.

The project described was supported by Grant Number 1C1CMS331055-01-00 of the Department of Health and Human Services, Centers for Medicare & Medicaid Services. Its contents are solely the responsibility of the authors and have not been approved by the Department of Health and Human Services, Centers for Medicare & Medicaid Services. The research presented in this study was conducted by the awardee. Findings might or might not be consistent with or confirmed by the independent evaluation contractor.

Author Disclosure Statement

No competing financial interests exist.

References

1. Detering KM, Hancock AD, Reade MC, Silvester W. The impact of advance care planning on end of life care in elderly patients: Randomised controlled trial. *BMJ* 2010; 340:c1345.
2. Leung JM, Udris EM, Uman J, Au DH. The effect of end-of-life discussions on perceived quality of care and health status among patients with COPD. *Chest* 2012;142:128–133.
3. Teno JM, Gruneir A, Nanda A, Wetle T. Association between advance directives and quality of end-of-life care: A national study. *J Am Geriatr Soc* 2007;55:189–194.
4. Houben CH, Spruit MA, Groenen MT, et al. Efficacy of advance care planning: A systematic review and meta-analysis. *J Am Med Dir Assoc* 2014;15:477–489.
5. Silveira MJ, Kim SY, Langa KM. Advance directives and outcomes of surrogate decision making before death. *NEJM*. 2010;362:1211–1218.
6. Sudore RL, Fried TR. Redefining the “planning” in advance care planning: Preparing for end-of-life decision making. *Ann Intern Med* 2010;153:256–261.
7. Brett AS. Limitations of listingspecific medical interventions in advance directives. *JAMA* 1991;266:825–828.
8. McMahan RD, Knight SJ, Fried TR, Sudore RL. Advance care planning beyond advance directives: Perspectives from patients and surrogates. *J Pain Symptom Manage* 2013;46: 355–365.
9. Sudore RL, Knight SJ, McMahan RD, et al. A novel website to prepare diverse older adults for decision making and advance care planning: A pilot study. *J Pain Symptom Manage* 2014;47:674–686.
10. Heiman H, Bates DW, Fairchild D, et al. Improving completion of advance directives in the primary care setting: A randomized controlled trial. *Am J Med* 2004;117:318–324.
11. Weinick RM, Burns RM, Mehrotra A. How many emergency department visits could be managed at urgent care centers and retail clinics? *Health Aff* 2010;29:1630–1636.
12. Smith AK, et al. Half of older Americans seen in emergency department in last month of life; most admitted to hospital, and many die there. *Health Aff (Millwood)* 2012; 31:1277–1285.
13. Oulton J, et al. Advance directives for older adults in the emergency department: A systematic review. *J Palliat Med* 2015;18:500–505.
14. Respecting Choices [Online]. www.gundersenhealth.org/respecting-choices/about-us/stages-of-planning (Last accessed May 16, 2016).
15. Grudzen CR, et al. Concordance of advance care plans with inpatient directives in the electronic medical record for older patients admitted from the emergency department. *J Pain Symptom Manage* 2016;51:647–651.
16. Fried TR, O’Leary J, Ness PV, Fraenkel L. Inconsistency over time in the preferences of older persons with advanced illness for life-sustaining treatment. *J Am Geriatr Soc* 2007; 55:1007–1014.
17. Smith AK, et al. Am I doing the right thing? Provider perspectives on improving palliative care in the emergency department. *Ann Emerg Med* 2009;54:86–93.
18. Stone SC, et al. Emergency medicine physicians’ perspectives of providing palliative care in an emergency department. *J Palliat Med* 2011;14:1333–1338.
19. Hess EP, et al. The chest pain choice decision aid: A randomized trial. *Circ Cardiovasc Qual Outcomes* 2012;5: 251–259.
20. Center for Medicare and Medicaid Innovation. Health care Innovation Award Project Profiles, 2012. [Online]. <http://innovation.cms.gov/Files/x/HCIA-Project-Profiles.pdf>. (Last accessed December 23, 2015).
21. Gilboy N, Tanabe T, Travers D, Rosenau AM. *Emergency Severity Index (ESI): A Triage Tool for Emergency Department Care, Version 4*. Implementation Handbook 2012 Edition. Rockville, MD: Agency for Healthcare Research and Quality, 2011.
22. Advance Directives and Advance Care Planning: Report to Congress, 1 August 2008. [Online]. <http://aspe.hhs.gov/basic-report/advance-directives-and-advance-care-planning-report-congress> (Last accessed November 22, 2015).
23. Greysen SR, Garcia CC, Sudore RL, et al. Functional impairment and internet use among older adults: Implications for meaningful use of patient portals. *JAMA Intern Med* 2014;174:1188–1190.
24. Smith A. Older adults and technology use, 3 April 2014. [Online]. www.pewinternet.org/2014/04/03/older-adults-and-technology-use/?utm_expid=53098246-2.Lly4CFSVQG2l

- phsg-KopIg.0&utm_referrer=https%3A%2F%2Fwww.google.com%2F (Last accessed December 3, 2015).
25. Post LA, Vaca FE, Biroscak BJ, et al. The prevalence and characteristics of emergency medicine patient use of new media. *JMIR Mhealth Uhealth* 2015;3:72.
 26. D'Onofrio GD, Pantalon MV, Degutis LC, et al. Development and implementation of an Emergency Practitioner-performed brief intervention for hazardous and harmful drinkers in the emergency department. *Acad Emerg Med* 2005;12:249–256.
 27. Fried TR, Redding CA, Robbins ML, et al. Stages of change for the component behaviors of advance care planning. *J Am Geriatr Soc* 2010;58:2329–2336.
 28. Ostherr K, Killoran P, Shegog R, Bruera E. Death in the digital age: A systematic review of information and communication technologies in end-of-life care. *J Palliat Med* 2016;19:408–420.

Address correspondence to:

Kei Ouchi, MD, MPH
Department of Emergency Medicine
Brigham and Women's Hospital
75 Francis Street, Neville 200
Boston, MA 02125

E-mail: kouchi@partners.org