CORRESPONDENCE

Refractive Errors—Epidemiology, Effects and Treatment Options

by Prof. Dr. med. Ulrich Schiefer, Christina Kraus, B.Sc. Augenoptik, Prof. Dr. rer. nat. Peter Baumbach, Judith Ungewiß, M.Sc., and Dipl.-Ing. (FH) Ralf Michels in issue 41/2016

Prof. Dr. med. Dieter Schmidt
Klinik für Augenheilkunde der Universität Freiburg, Germany
dieter.schmidt@uniklinik-freiburg.de

Conflict of interest statement
The author declares that no conflict of interest exists.

Some Additions

Schiefer et al. provided a comprehensive and instructive overview of the most important refractive errors, highlighting typical functional impairments of relevance to physicians not specialized in ophthalmology (1).

I would like to contribute the following additions:

In her handbook article (p. 972), Aulhorn (2) already reported that the shortsighted often experience poor vision at night, especially while driving. In her opinion, some ophthalmologists do not completely correct myopia in some cases. Due to high luminance, pupils are small during the day. In this situation, slightly under-corrected glasses are generally well tolerated, but at night, when the pupils are wider, any loss in visual acuity may be experienced as troublesome. In addition, a true night myopia of about 1.0 to 1.5 dpt may occur in some people. These patients require glasses for night driving.

Schiefer et al. introduced the internationally renowned and diagnostically important Amsler test for the detection of metamorphopsia. In addition to the Amsler grid, Weiss (3) pointed out the diagnostic importance of a so far little known test which involves the assessment of lines running in a spoke-like fashion to a center point. Binder et al. (4) highlighted the existence of “labile” metamorphopsia detectable with this spoke test.

Schiefer et al. introduced the computerized visual acuity testing which plays an important role in clinical practice and is available online. For the sake of completeness, I would like to add here that Crossman et al. and Timberlake et al. already emphasized the importance of the computerized acuity testing in 1970 and 1980, respectively.

DOI: 10.3238/arztebl.2017.0195a

REFERENCES

1. Schiefer U, Kraus C, Baumbach P, Ungewiß J, Michels R: Refractive errors—epidemiology, effects and treatment options. *Dtsch Arztebl Int* 2016; 113: 693–702.
2. Aulhorn E: Funktionsprüfung des Auges zur Feststellung der Kraftfahreignung. In: Straub W (ed.): *Die ophthalmologischen Untersuchungsmethoden*, Band II. Stuttgart: F. Enke Verlag 1976; 922–90.
3. Weiss H: Untersuchungen zur Bestimmung und Bewertung von Metamorphopsien. *Graefes Arch Klin Exp Ophthalmol* 1971; 183: 105–36.
4. Binder S, Kutschera E, Weiß H: Die Makulafunktion nach postoperativer Wiederanlegung der Netzhaut. *Klin Monatsbl Augenheilkd* 1977; 171: 606–10.

Caution Required in Cases With Vitreous Opacities

Following radiological, otolaryngological, neurological and other diagnostic investigations, countless patients beyond 40 years of age finally present to an ophthalmologist with characteristic symptoms, including fluctuating vision, eye pain and headache, as well as dizziness—in severe cases with nausea.

In most of these cases, the medically appropriate and, from the perspective of health economy, required approach is to perform ophthalmological tests to confirm or exclude suspected refractive errors first, before undertaking costly further examinations at the expense of the community of the insured. In urgent cases, an ophthalmological diagnosis is typically established or excluded on the same day.

It is certainly advisable that non-ophthalmologists perform pinhole testing, if the results are conclusive, especially when there is no better alternative diagnostic test available at that location and time. This test can easily be performed with just a little practice, even without previous training in ophthalmology.

But caution is required: In patients experiencing a reduction in visual acuity due to vitreous opacities, a pinhole occluder can reduce visual acuity at one occasion, yet increase it at another time, which may surprise an inexperienced examiner who follows the recommendations made in the authors' article (1). Vitreous opacities can be a harmless symptom associated with the aging of the vitreous body, but may also be a sign of retinal detachment or even vitritis (inflammation of the vitreous body) in cases of uveitis (inflammation of the uvea).

Thus, the presence of a harmless refractive error should only be assumed in cases where a clear and reproducible improvement of central visual acuity is achieved with a pinhole occluder; only in these cases, the diagnostic workup of the problem can be rightfully postponed to a later date.

An example: How often do ophthalmologists assess the visual acuity in children who did not provide meaningful information when examined by a pediatrician—despite the fact that this is a standard part of the pediatric screening examinations? However, there is no point in using the pinhole test in children, as from experience we know that this leads to strong accommodation.

DOI: 10.3238/arztebl.2017.0195b

REFERENCES

1. Schiefer U, Kraus C, Baumbach P, Ungewiß J, Michels R: Refractive errors—epidemiology, effects and treatment options. *Dtsch Arztebl Int* 2016; 113: 693–702.

PD Dr. med. Dipl.-Phys. Jean-Cyriaque Barry
Ophthalmologist
Solingen, Germany
drbarry.augen@arcor.de

Conflict of interest statement

The author declares that no conflict of interest exists.

In Reply:

The authors would like to thank Dr. J. C. Barry for his comments on our article (1). The authors' aim was to prevent the patients' odyssey from specialist to specialist described in the opening part of his letter. The diagnostic tests mentioned are performed easily and quickly. With the help of these tests, a physician who is not practicing in ophthalmology can adequately differentiate between the described visual disturbances and refer a patient to an ophthalmological-optical specialist, if necessary—avoiding the long journeys described in the letter.

There seems to be general agreement about the fundamental value of the pinhole test. In the opinion of the authors, the vitreous opacities mentioned by Dr. Barry are not included in the refractive errors addressed in this publication. (Circumscribed) vitreous opacities typically result in fluctuating local visual impairments. Those affected note the characteristic symptom of vitreous floaters, in particular along with eye movements while looking at a homogenous bright background: Typically, the shadows of such vitreous opacities continue moving even after the eye movement has stopped. This is explained by the inertia of the vitreous body (and circumscribed opacities contained therein) in relation to the wall of the eyeball.

In young individuals (and especially in children), viewing through a pinhole could in principle trigger accommodation. However, the authors think that this is unlikely, for the following reasons: In contrast to the so-called “instrument myopia“, which occurs, for example, when looking through an optical instrument, the overall depth of a pinhole is negligibly small. In

addition, when a sieve-like pinhole occluder is held in front of a patient's eyes, the impression of looking into an instrument does simply not occur (due to the numerous holes available to look through).

The authors would like to thank Prof. Schmidt for his additions. Due to the limitations imposed by the publisher with regard to text length and number of references, a CME article can never be exhaustive.

The condition of night myopia mentioned by Prof. Schmidt can indeed be a reason for nocturnal visual disturbances. Night myopia can be caused by numerous factors, including but not restricted to the presence of aberrations along with a large pupil diameter, e.g. in the form of spherical aberrations (due to an increased refractive power of peripheral parts of the lens) (2, 3).

With regard to the examination of metamorphopsia, the authors are grateful for the comments pointing out further methods available for the assessment of this visual disturbance. In this context, the method of the working group led by C. Matsumoto is worth mentioning as it allows to quantify this visual disturbance (4).

The authors did not intend to present the historical development of computerized visual acuity testing. They rather wanted to present the design of the Freiburg Visual Acuity Test (FrACT), a widely adopted and well-documented acuity testing tool which is available online.

DOI: 10.3238/arztebl.2017.0196

REFERENCES

1. Schiefer U, Kraus C, Baumbach P, Ungewiß J, Michels R: Refractive errors—epidemiology, effects and treatment options. *Dtsch Arztebl Int* 2016; 113: 693–702.
2. López-Gil N, Peixoto-de-Matos SC, Thibos LN, González-Méjome JM: Shedding light on night myopia. *J Vis* 2012; 12: 4, 1–9.
3. Hope GM, Rubin ML: Night myopia. *Surv Ophthalmol* 1984; 29: 129–36.
4. Nomoto H, Matsumoto C, Arimura E, et al.: Quantification of changes in metamorphopsia and retinal contraction in eyes with spontaneous separation of idiopathic epiretinal membrane. *Eye (Lond)* 2013; 27: 924–30.

Corresponding author:

Prof. Dr. med. Ulrich Schiefer
Hochschule Aalen, Germany
ulrich.schiefer@hs-aalen.de

Conflict of interest statement

Prof. Schiefer has received honoraria for consultancy from Servier, Pharm-Allergan, and Haag-Streit and payments for lectures from Alcon Pharma, Pharm-Allergan, MSD-Chibret, Pfizer, and Oculus.