Published in final edited form as:

Cell Mol Life Sci. 2017 May ; 74(10): 1819–1834. doi:10.1007/s00018-016-2439-z.

Microfabricated tissues for investigating traction forces involved in cell migration and tissue morphogenesis

Bryan A. Neger¹, Michael J. Siedlik¹, and Celeste M. Nelson^{2,*}

¹Department of Chemical & Biological Engineering, Princeton University, Princeton, NJ 08544

²Department of Molecular Biology, Princeton University, Princeton, NJ 08544

Abstract

Cell-generated forces drive an array of biological processes ranging from wound healing to tumor metastasis. Whereas experimental techniques such as traction force microscopy are capable of quantifying traction forces in multidimensional systems, the physical mechanisms by which these forces induce changes in tissue form remain to be elucidated. Understanding these mechanisms will ultimately require techniques that are capable of quantifying traction forces with high precision and accuracy *in vivo* or in systems that recapitulate *in vivo* conditions, such as microfabricated tissues and engineered substrata. To that end, here we review the fundamentals of traction forces, their quantification, and the use of microfabricated tissues designed to study these forces during cell migration and tissue morphogenesis. We emphasize the differences between traction forces in two- and three-dimensional systems, and highlight recently developed techniques for quantifying traction forces.

Keywords

traction force microscopy; extracellular matrix; mechanotransduction; morphodynamics

Introduction

Mechanical forces play an integral role in cellular behaviors as diverse as proliferation [1,2], differentiation [3,4], gene expression [5,6], wound healing [7], tumor metastasis and invasion [8–10], collective migration [11,12], and tissue morphogenesis [13]. In particular, traction stress, which refers to the force per unit area acting on the deformed cell-matrix interface, is a key mechanical cue in cellular systems. Traction forces originate via the contractile actomyosin network and can alter cellular behavior through a number of mechanisms collectively referred to as mechanotransduction [14]. However, the biophysical and biochemical mechanisms by which these forces induce changes in tissue form are not fully understood. The ability to accurately and precisely calculate traction forces is therefore important and will strengthen our quantitative understanding of their role in cell- and tissue-level processes.

* Address correspondence to C.M.N., 303 Hoyt Laboratory, William Street, Princeton, NJ 08544, Tel: 609-258-8851, Fax: 609-258-1247, celesten@princeton.edu.

Possible reviewers: Eric R. Dufresne; Alexander R. Dunn; Ben Fabry; Jeffrey J. Fredberg

The idea that the laws of mechanics can be applied to cells was first postulated in the late 1800s by Wilhelm His [15,16]. Using the developing chicken embryo as a model, His proposed that differential growth rates in the germ layers gave rise to mechanical forces that are responsible for producing the final tissue form [15,17]. Yet, despite this early insight, quantifying cell-generated mechanical forces such as traction forces—either *in vivo* or in culture—presents a tremendous challenge. The inherently small size of cells (~10 μm in diameter) means that traction forces act over a small area. When coupled with the fact that the magnitude of the corresponding forces is ~10 nN [18], it becomes clear that traction forces are inaccessible to macroscopic force measurements. Furthermore, mechanical forces act over both cell- and tissue-level length scales, which further complicates any attempts at measurement since the scale at which forces should be measured is a non-trivial consideration [19]. Despite these challenges, a diverse group of experimental techniques enable the quantification of traction forces in two (2D) and three dimensions (3D) [20].

Several excellent reviews [21,20,22,23] have summarized the role of traction forces in various biological processes and the corresponding tools available to quantify them. Here, we focus specifically on the advancement of traction force microscopy (TFM) and microfabricated tissues for quantifying traction forces in the context of cell migration and tissue morphogenesis. Throughout the review, we discuss the differences between traction forces in 2D and 3D systems and highlight recent developments for traction force quantification. It should be noted that we use the term tractions in cases where traction stress and traction force can be used interchangeably or for consistency when referring to published work that used the term tractions. TFM has emerged as the most widely accepted approach for quantifying traction forces owing to a number of advantages [21]. Above all, TFM can be performed without specialized equipment, and tractions can be calculated using MATLAB code available online [21], which makes the technique readily accessible to most research labs. TFM is also remarkably versatile owing to the fact that force calculations are not inherently limited to any length scale [21].

We begin with a description of experimental progress for traction force quantification. Next, we discuss computational approaches to calculate traction forces, and provide specific examples for the application of TFM to cell migration and morphogenesis. Finally, we discuss the design and use of TFM in microfabricated tissues as well as strategies for overcoming limitations of conventional TFM. While both TFM [20–22] and micropatterning [24–26] have been previously reviewed, here we focus specifically on traction force quantification in the context of multidimensional cell migration and tissue morphogenesis.

Traction forces: multiple decades of experimental progress

The quantification of cell-generated traction forces was pioneered by Harris and co-workers in 1980 [27]. In this seminal study, traction forces were determined by measuring substratum deformations in the form of cell-induced wrinkles at the surface of a thin flexible silicone membrane with known elastic properties. This technique built off two similar approaches that had previously been used to study distortions and birefringence in gelatin [28] and thin plasma clots [29], but which were limited by substratum stability [27]. Quantifying traction forces from wrinkles in the substratum is inherently challenging because the wrinkles are

often larger than the cells and form gradually over time in a non-linear and chaotic manner [30]. Harris's technique was subsequently improved by introducing a stretchable non-wrinkling silicone substratum that incorporated beads as fiducial markers [31,32]. The 2D displacement of these beads was measured and used to calculate traction forces with higher accuracy than could be obtained by measuring wrinkles in the substratum [31,32].

Modifications to the deformable substratum were also investigated in order to obtain additional improvements in traction force quantification. Burton and Taylor developed a new silicone polymer substratum with UV-tunable stiffness in order to control the size of wrinkles such that the length could be measured and the movement of the substratum could be minimized [33]. An alternative approach quantified traction forces from the deformation of collagen gels, which were assumed to approximate an elastic material and were more representative of the ECM encountered by cells in an *in vivo* microenvironment [34]. After polyacrylamide (PA) gels were introduced as a cell culture platform [35,36], these substrata were used to investigate the mechanical forces exerted by cells [37]. Compared to silicone substrata, PA gels exhibit a number of advantages: the stiffness can be tuned by changing the relative concentrations of acrylamide and bisacrylamide; substratum deformation is proportional to force applied over a range of forces; the optical properties of the material enable high-magnification imaging; substratum porosity allows media penetration such that cells can be embedded within the gel; and PA is relatively inert [37,38]. Thick PA gels have been used to produce spatial maps of traction forces exerted by fibroblasts during steady locomotion [30]. Recently, work using PA gels has improved computational and spatial resolution by using fiducial markers that can be imaged in separate fluorescence channels [39]. Lastly, tunable viscoelastic substrata have been used to study cell tractions in an environment that more closely represents the human body [40].

The spatial resolution of traction force quantification was also improved with the creation of micropatterned substrata such as arrays of microposts. Similar to PA gels, micropatterned substrata allow the mechanical properties and surface chemistry to be tuned independently. These systems typically consist of elastic pillars that deform from cell-generated forces [41]. Alternative approaches have also used discrete arrays of cantilever sensors [42] to quantify traction forces at different positions under the cell. Provided that the density of pillars or cantilevers is high enough, the traction forces can be quantified at different focal adhesions within the same cell [43]. Arrays of microposts have also been used to measure the velocity and power of migrating cells [44]. In using these approaches, the density of pillars needs to be sufficiently high such that the surface can be approximated as a continuum [45]. In addition, any calculations performed must account for substratum warping that can occur if the pillars are made from the same elastic material as their underlying support [46].

Initially, 2D traction force calculations relied on an in-plane assumption that there was no stress normal to the substratum beneath the cells [47]. However, this assumption was found to be invalid under certain experimental conditions because cellular contractility can induce traction forces normal to the underlying substratum [47,48]. As a result, 2.5D TFM methods were developed to calculate both in-plane and normal traction forces [18,47,49]. Here, 2.5D refers to the ability of the TFM technique to quantify 3D traction forces—both in-plane and normal—for cells on a 2D planar surface. Both in-plane and normal traction forces were

found to be of the same order of magnitude during the migration of many types of individual cells [50,47]. The first quantification of traction forces in a purely 3D setting used the displacements of beads generated by cells embedded in gels of polyethylene glycol (PEG), from which the corresponding tractions were determined using linear elastic theory [51]. Additional studies have since quantified traction forces for individual cells embedded in highly nonlinear biopolymer networks consisting of collagen, fibrin, and Matrigel [52]. Traction force calculations have also been performed for single cells [53] and tissues [54–56] embedded in 3D matrices of collagen and Matrigel. Taken together, these techniques are well suited to quantify cell-induced tractions in many different contexts and provide the user with the flexibility to choose the setup that best matches experimental requirements.

Quantifying traction forces

As described previously, traction forces are not measured explicitly. Rather, TFM relies on measuring the physical displacement of fiducial markers or the deformation of a substratum. For single isolated cells, local displacement data are combined with the appropriate constitutive equations in order to calculate local traction forces (Figure 1A and C). For multicellular systems, global forces can be calculated from these local tractions using Newton's laws of motion (Figure 1B and D). In this section, we outline a previously described computational approach [21] for calculating local traction forces directly from experimentally-measured displacement data. We further discuss the limitations associated with traction force calculations as well as some recent developments that aim to improve computational efficiency.

For sufficiently small deformations of elastic substrata, discrete traction forces (F) can be calculated using Hooke's law (Eq. 1):

$$F=kx \quad (1)$$

where k is the spring constant and x is the displacement [57]. The accuracy of the calculation is a function of the material properties of the substratum, the resolution of the displacement measurements, and the validity of the assumption that the deformations are sufficiently small [16]. Although Hooke's law provides a simple approach to calculate traction forces at discrete locations, traction forces are actually distributed continuously throughout the substratum [21]. As a result, it is convenient to represent traction forces in terms of traction stress, which represents the traction force per unit contact area between the cell and the underlying substratum or the surrounding matrix [21]. Furthermore, if the substratum is assumed to be a linearly elastic solid, then the traction stress, $\sigma(\mathbf{x})$, is related to the displacement, $\mathbf{u}(\mathbf{x})$, by Hooke's law in tensor form [58,21] (Eq. 2):

$$\sigma = \frac{E}{1-\nu} \left(\frac{1}{2}(\nabla \mathbf{u} + \nabla \mathbf{u}^T) + \frac{\nu \nabla \cdot \mathbf{u}}{1-2\nu} \mathbf{I} \right) \quad (2)$$

where ν is Poisson's ratio, E is the elastic modulus, and $\mathbf{I}$ is the identity tensor. It is important to note that boundary conditions at the top and bottom surfaces of the substratum

and the assumption of mechanical equilibrium ($\nabla \cdot \sigma = 0$) are both required to solve Eq. 2 [58]. These assumptions give Eq. 3 [21]:

$$(1 - 2\nu)\nabla^2 \mathbf{u} + \nabla(\nabla \cdot \mathbf{u}) = 0 \quad (3)$$

which can be solved using a variety of mathematical manipulations, including Fourier transforms [21,59] and finite element analysis [49,60]. Using similar approaches, traction forces have been mapped for single cells and populations of cells cultured on a planar substratum (Figure 2A and B) as well as embedded in a matrix (Figure 2C and D).

One drawback of the aforementioned approach is that calculating the traction stresses requires knowledge about the properties of the underlying matrix, such as the elastic modulus. This presents a challenge for *in vivo* systems where it may be difficult to characterize the mechanical properties of the ECM owing to its complex composition and because cells continuously remodel the ECM as they migrate. One potential solution was recently proposed whereby mean deformation metrics could be used to quantify cell-matrix interactions without the mechanical details of the matrix [61]. However, this method does not provide any information about cell tractions and cannot describe the effects of matrix stiffness without first knowing the mechanical properties of the matrix [61]. A second drawback is that the inverse calculations required to obtain traction stresses can be computationally expensive. One potential solution to this problem is digital volume correlation, which uses volume subsets to track 3D displacement fields. This method allows for the direct construction of traction stresses [62,63] and therefore avoids the need for inverse calculations, allowing one to determine traction fields directly from volumetric confocal image stacks [18].

Computational advances in TFM calculations were realized with the understanding that explicitly quantifying traction stresses throughout the substratum requires displacement information to be processed using a statistical procedure [64]. Building on the work of Oliver [32] and Lee [31], Dembo used a Bayesian method to map traction forces from measurements of substratum displacement [64]. However, as pointed out by Schwarz [65], the solution obtained from the Bayesian method depends strongly on fluctuations in the experimental data. Alternative approaches, such as that of Tichonov [66] and one based on classical functional analysis [60], have also been proposed to circumvent the limitations of the Bayesian method. For more information, the interested reader is directed to recent reviews on the limitations associated with these approaches [67,68].

Application 1: forces exerted during cell migration

The study of mechanical forces in cell migration was first performed qualitatively by examining changes in cell shape during the migration of isolated epithelial cells and cell sheets in 2D cell cultures [69]. While qualitative observations provide relative information about the magnitude of mechanical forces, these approaches are inherently limited by their inability to explicitly quantify forces in a reproducible manner. To that end, a variety of experimental tools have been developed to study the role of mechanical forces in migration

[20] and the associated signaling pathways [70–72]. Studies using TFM-based analysis of 2D and 3D cell migration have generally reported traction stresses with orders of magnitude ranging from 10 Pa to 10^3 Pa [48,52,8]. Depending on the experimental conditions and cell type, the magnitude of traction stresses either remains constant or fluctuates during cell migration.

The role of traction forces in the migration of individual cells has been studied in fibroblasts [8], epithelial cells [45], *Dictyostelium* cells [50], cancer cells [9,73], myoblasts [74], neutrophils [75], dendritic cells [76], keratinocytes [77], and bovine aortic endothelial cells [47], among others. Cells on 2D planar surfaces were found to exert both in-plane and normal tractions as they migrated (Figure 1A). The relative magnitude of these tractions depends on a number of factors including the cell type, the thickness and mechanical properties of the substratum [78], ligand density [79], chemotactic gradients [75], the cell disease state [8,9], and the calculation approach. For example, a study of *Dictyostelium* cells on thick gel substrata [50] reported higher in-plane traction stress while a similar study of *Dictyostelium* cells on thin gel substrata reported higher normal traction stress [48]. The contradictory results in these studies can likely be attributed to the fact that thick isotropic gels show a similar stiffness in response to in-plane and normal traction stresses whereas thin gels do not. This discrepancy arises because thin gels violate the macroscopic isotropy condition, meaning that properties such as gel stiffness vary in magnitude in the normal and in-plane directions within the gel [80,48]. In general, in-plane traction forces are observed throughout the cell with the largest traction forces near the edge, while normal traction forces are concentrated near the center of the cell. The distribution of in-plane traction forces for individual cells is consistent with the idea that protrusions at the leading edge push forward while pulling the rest of the cell during migration. On the other hand, the distribution of normal traction forces is believed to arise from the compression of the nucleus, which is pushed down into the soft substratum by the cell cortex [50].

Far less is known about traction forces during the migration of cells in 3D microenvironments because of challenges associated with imaging and sample preparation. Specifically, measuring bead displacement in 3D gels can be problematic because of working distance limitations associated with high magnification objectives [81]. Moreover, precisely controlling the 3D microenvironment surrounding migrating cells is difficult [82]. Nevertheless, a number of studies have investigated tractions for single isolated cells in 3D matrices [51,52,10]. Legant and co-workers reported that cells embedded in a 3D matrix mainly exerted shear tractions, but there were also small normal tractions close to the body of the cell [51]. Similar to single cells on planar surfaces, tractions were reported to increase with increasing distance from the cell's center of mass [51]. Breast cancer cells were found to generate traction forces independent of collagen concentration and matrix stiffness in 3D, which is in contrast to the observations for cells on planar surfaces [52]. Moreover, 3D tractions were found to be considerably higher for invasive lung and breast cancer cells, and polarized traction fields were found to facilitate invasion [10].

In addition to studying individual cell migration, traction forces have been investigated during the collective migration of endothelial [83] and epithelial [54,11,12,83,45] cells. Reflecting the multicellular nature of collective migration, these migrating cellular cohorts

exert larger tractions than cells migrating individually [45]. A study of the traction forces exerted by monolayers of Madin–Darby canine kidney (MDCK) epithelial cells found that in-plane traction stresses were smaller than normal traction stresses [83], though no clear trend between the magnitude of normal and in-plane stresses has been reported. For 2D systems consisting of sheets of cells on planar surfaces, the largest tractions were typically observed at the leading edge of the cell monolayer. This observation is consistent with the idea that leader cells—cells at the front of a migrating cell cohort—exert larger tractions than do follower cells [84]. For example, in sheets of MDCK epithelial cells, in-plane traction stress calculations revealed a decrease in average traction stress with increasing distance from the edge of the cell sheet [45]. In contrast, Trepap and co-workers found that collective migration of cell sheets on planar surfaces occurs through a global tug-of-war mechanism, and not as a result of cell pulling by leader cells or self-propelled cells [12]. A TFM technique known as monolayer stress microscopy [85] was recently created in order to generate stress maps throughout an entire cell sheet [83]. This approach revealed that local intercellular shear stress was minimized during collective migration [83]. During collective migration within 3D matrices, traction stresses were also found to be highest at the leading edge of the migrating cohort [55,54,56].

Taken together, the aforementioned studies demonstrate the remarkable utility of TFM for quantifying traction forces during the migration of individual cells and collectives in both 2D and 3D systems. That being said, the fact that TFM studies use different experimental conditions and analysis techniques makes it difficult to draw concrete comparisons and make conclusions about the role of traction forces during cell migration. Future work should continue to focus on the creation of generalized experimental protocols for investigating traction forces. This would allow researchers to determine the extent to which the magnitude and spatiotemporal distribution of traction forces present during migration are conserved across different cell types and experimental conditions.

Application 2: forces exerted during tissue morphogenesis

Given their role in cell migration, it is apparent that traction forces are also likely to be important in various morphogenetic processes [86–89]. Indeed, the role of mechanical forces in embryonic development has been recognized since the early 20th century [90]. Moreover, the relationship between mechanical forces, cell growth and death [3,91,92], and tissue geometry is also well appreciated [93,94]. Nonetheless, the physical mechanisms by which traction forces induce changes in tissue form during development remain largely unknown because it is difficult to explicitly quantify traction forces *in vivo*.

The primary challenge facing attempts to quantify traction forces *in vivo* arises from the complex heterogeneous microenvironment surrounding cells, which has poorly defined mechanical properties. To date, the only techniques available to determine forces or stresses *in vivo* are laser ablation [95] and the use of mechanically deformable injected microdroplets [96]. Laser ablation uses thermal energy to selectively destroy tissue at the cellular or sub-cellular level [97] and induce relaxation of the ablated region, yielding values for the tensile forces present within the tissue. In contrast, the microdroplet technique is performed by injecting fluorescent oil microdroplets with defined mechanical properties near cell-cell or

cell-ECM interfaces [96]. Deformation of the microdroplets, which are coated with ligands for a mechanical link to cell surface adhesion receptors, can be used to determine anisotropic stresses *in vivo*. The forces that drive tissue morphogenesis have been investigated in various contexts, including epithelial spreading during zebrafish gastrulation [98], elongation of *Drosophila melanogaster* embryos [99], and differentiation of muscle cells before and after myotube formation [74].

Several excellent reviews discuss morphogenesis in a broader context [88,89,100], as well as the role of mechanical forces in morphogenesis [101,102,13,103,87]. Quantifying traction forces *in vivo* remains a key challenge that must be addressed in order to fully understand the role of these forces in morphogenesis. Future work should focus on the improvement of existing *in vivo* traction force techniques as well as the creation of more precise and accurate approaches. As these approaches proceed into and through their infancy, it is imperative to develop and use culture models that faithfully represent *in vivo* processes. Such models, discussed below, offer a unique avenue for precisely controlling the form of multicellular structures and the properties of the surrounding matrix. In particular, we highlight the ability of microfabrication-based methods to control 2D and 3D tissue form to more accurately represent *in vivo* conditions, standardize results to tissue shape, provide statistical analysis, and make heterogeneous environments by spatially controlling substratum properties.

Designing microfabricated tissues to study traction forces

Traditionally associated with materials science and microelectronics, microfabrication-based methods in biology rely on top-down schemes to dictate the placement and organization of ECM proteins [24–26]. For example, investigators in the 1990s used photolithography to define regions of protein adsorption on a silicon surface [104]. The advent of soft lithography in the same decade simplified the printing of proteins onto 2D surfaces [105,106] and enabled the formation of micromolded 3D ECM structures [107]. Custom 3D ECM gels and scaffolds have also been generated using sacrificial molds [108], as well as inkjet and 3D printing [109,110]. 2D or 3D microfabricated tissues can be formed when cells are added to the matrix framework fashioned by these methods and allowed to form multicellular structures.

Microfabricated tissues offer several advantages for investigating traction stresses. The engineering control provided by microfabrication enables investigators to parse the relationship between signaling, geometry, mechanics, and changes in tissue shape during culture processes that mimic *in vivo* behavior; one can control matrix cross-linking, stiffness, and ligand composition. Furthermore, fiducial markers can be incorporated into the 2D or 3D substratum. As described above, the displacement of these markers near the cell-matrix interface provides insight into the forces generated by cells and tissues. Microfabrication-based techniques can also be used to produce large arrays of identical 2D or 3D tissues with well-defined shapes and sizes. This feature of microfabrication-based systems ensures that experiments are particularly high-throughput, a valuable attribute for efficient statistical analyses and for investigating multiple conditions simultaneously.

Microfabricated tissues both recreate aspects of *in vivo* systems and also offer several advantages traditionally associated with cell culture. Tissues can be created from cells that have genetic modifications introduced via well-established transfection and transduction methods. Coculture is readily included within these systems, thereby enabling investigation of the physical and biochemical interplay between multiple cell types [111–113]. Finally, the motion of cells and matrix is easily imaged, providing investigators with a straightforward approach for quantifying live cell and tissue behavior. Taken together, microfabricated tissues have and will continue to provide valuable insight into multicellular processes. The following subsections detail some 2D- and 3D-microfabricated tissue systems that have been used, paying particular attention to those systems that quantify forces generated by individual cells and multicellular tissues.

Using 2D microfabricated tissues to investigate traction forces

2D tissues with tunable shapes and sizes are formed by exposing cells to surfaces that contain well-defined regions of matrix proteins surrounded by non-adhesive regions. The cells adhere to these regions and proliferate over time to fill the available matrix template. One common technique to produce a surface patterned with adhesive regions (the critical design requirement here) is to use stamps coated with ECM protein. This process, known as microcontact printing, transfers proteins [3,114] or other ligands [115,91] to a surface in a manner defined by the bas-relief of the stamp (Figure 3A). The regions surrounding the deposited protein are then passivated, ensuring that cells adhere only in the desired pattern [116].

Microcontact printing can be used with either PA gels or flexible microposts to investigate traction forces generated by cells and tissues of specific geometry or configuration (Figure 3A). PA gels directly stamped with inorganic oxidizing agents produce surface patterns presenting free aldehydes that can then bind to the desired matrix protein [117]. Alternatively, matrix micropatterns can be incorporated onto the gels during the PA polymerization step [118], which occurs between glass slides. One of the glass slides can be printed with the desired proteins and placed in contact with the polymerizing solution to transfer the protein from the slide to the gel. This technique provides a potential advantage when using very soft hydrogels, which are much more sensitive to unintentional mechanical agitation. Microcontact-printed PA gels have been used to quantify how the composition of ECM ligands influences the magnitude of traction forces [119]. These systems have revealed that traction forces also depend on cell shape and correlate with the differentiation of mesenchymal stem cells [117]. In micropatterned multicellular sheets, high levels of traction stress correlate with regions of elevated proliferation [93].

In a similar approach, distinct adhesive structures can be prepared by microcontact printing proteins directly onto elastic microposts [120–122]. The mechanical properties of the posts can be tuned by varying the ratio of silicone monomer to crosslinking agent or by adjusting the height of the pillars to provide a 1000-fold dynamic range of available post rigidities [121]. The protein of choice is stamped directly onto the surface of the microposts and the side regions are passivated. To date, experiments using microposts printed with distinct regions of protein have revealed that projected cell area, substratum stiffness, and post

density dictate the strengthening of focal adhesions in a manner that closely matches the trends observed for traction forces [122]. In the context of TFM, this system has been used to quantify traction forces within epithelial monolayers migrating along tracks of printed ECM ligands [120]. Microcontact printing offers well-established methods for investigating the traction forces generated by multicellular tissues.

In addition to microcontact printing, micropatterned PA gels can also be produced by stenciling islands of protein onto the surface by adsorbing solution through the holes in microfabricated membranes (Figure 3A) [123]. In this way, the chemical conjugator (*e.g.* sulfo-SANPAH) binds only to regions defined by the holes in the membrane stencil, thereby enabling the production of well-defined islands of ECM protein on the PA surface. This technique has revealed that in epithelial tissues, epithelial-to-mesenchymal transition takes place in regions that experience high traction stress [124].

One drawback of the 2D microfabrication-based strategies described above is that they produce tissues with shapes defined in a photolithography step that occurs well in advance of the actual experiment. One strategy for overcoming this limitation and providing greater experimental flexibility may involve inkjet printing either the adhesive ligand or the cells themselves onto a surface [109]. More interestingly, switchable surface coatings [125–127] may be combined with the techniques described here to both define the initial adhesive regions and to provide temporal control of the shape of microfabricated tissues. In these studies, light-mediated cleavage or electrochemical desorption converts initially passivated regions into domains that support cell adhesion. Incorporating switchable surfaces with microposts or PA gels may enable quantification of the traction forces exerted in 2D as monolayers of cells migrate collectively from a confined region and along a dynamically-defined path. 2D microfabricated tissues are currently well suited for investigating traction forces, migration, differentiation, and proliferation, though there will undoubtedly be many more advances in the coming years.

Using 3D microfabricated tissues to investigate traction forces

When deciding between 2D and 3D microfabricated tissues, the investigator is faced with tradeoffs. On one hand, 2D microfabricated tissues are often simpler and easier to produce. On the other hand, 3D microfabricated tissues exist in a 3D ECM microenvironment that more accurately mimics *in vivo* conditions. 3D microfabricated tissues are generated by adding cells to 3D matrix scaffolds engineered into a desired shape and size. As described in the following section, these 3D scaffolds are often produced via micromolding of an ECM gel or through microfabrication of silicone elastomer support structures.

Replica micromolding can be used to produce 3D substrata within gels of ECM protein (Figure 3B) [56,107,128,129]. As with microcontact printing, this method uses silicone stamps with surface features defined via photolithography. In this case, a solution of ECM protein is allowed to gel around the stamp, and removing the stamp produces the inverse features in the gel. Cells can then be added to these scaffolds and a subsequent layer of solution can be allowed to gel on top of the cells, thereby producing 3D microfabricated tissues fully immersed within an ECM microenvironment.

3D microfabricated tissues produced via replica micromolding have been used to relate cell-generated traction forces to changes in gene expression, biochemical signaling, and 3D collective migration [130–133,55,134]. These tissues often consist of tubular epithelial structures embedded within a gel of type I collagen. Recently, it was demonstrated that 3D tensile forces drive migration of invasive multicellular cohorts from these microfabricated tissues [54]. These tensile forces activate mechanically sensitive signaling pathways and facilitate matrix remodeling at the leading edge of the cohort. This was the first study to investigate the spatiotemporal nature of physical forces exerted by a migrating cohort fully embedded in 3D matrix.

Microfabricated tissue gauges provide another opportunity to quantify forces from 3D microscale constructs of cells (Figure 3B) [135,136]. Micrometer-scale cantilevers are constructed from silicone elastomer in a manner defined by a multilayer photolithographic process. A solution of collagen and cells can be added to the cantilevers and allowed to gel. The cells compact the collagen gel, forming 3D tissues suspended between multiple cantilevers. By monitoring changes in cantilever deflection, one can calculate the tension generated by the tissue during processes such as 3D wound healing [137]. These tissues have been especially useful for investigating relationships between physical forces generated by a tissue and ECM protein deposition and remodeling [135–137].

To date, most TFM-based studies involving microfabricated tissues have used replica micromolding or other similar techniques. Tissues may also be formed by polymerizing an ECM gel around a sacrificial mold, such as a glass micropipette [108]. Alternatively, 3D printing can be used to produce custom substrata via layer-by-layer additive manufacturing [110]. Given that this technology has been used to produce scaffolds for artificial organs, it seems plausible that 3D printing could generate TFM-accessible microenvironments that more accurately mimic *in vivo* conditions. Taken together, there are many exciting opportunities for using engineering-based approaches to study the traction forces associated with cell migration and tissue morphogenesis. A summary of microfabrication techniques that have been used to quantify traction forces is provided in Table 1.

Additional biological insights enabled by microfabricated tissues

Beyond applications for TFM, microfabricated tissues can offer insight into an array of biological processes in development and disease. For example, microfabricated tissues have been used to elucidate the physical and chemical mechanisms involved in a developmental process called branching morphogenesis. Specifically, 3D microfabricated mammary epithelial tissues were used to reveal the interplay between tissue geometry and factors such as mechanical stress [55], autocrine inhibitory morphogen concentration [133], and epithelial-mesenchymal transition [124]. Microfabricated tissues have also been used as models to study diseases including cancer, where they revealed the relationship between mechanical stress and breast cancer cell invasiveness [113] as well as the interplay between interstitial fluid pressure, gene expression, and collective invasion of engineered breast tumors [108]. Additional biological processes such as collective cell migration [54], wound healing [137] and cell sorting [138] have also been studied using microfabricated tissues.

Interested readers can find a more comprehensive discussion of tissue micropatterning in literature [139].

Strategies for overcoming the limitations of conventional TFM

The spatiotemporal accuracy of quantified traction forces has been significantly improved since the original work of Harris and colleagues. Nevertheless, the accuracy of TFM is inherently limited because force is calculated implicitly from measured displacement data. Moreover, this calculation procedure can be computationally expensive, which limits the complexity of the biological systems that can be studied. Additional factors, including the resolution of the fiducial markers, substratum properties, and the validity of the underlying constitutive model used in force calculations limit the accuracy of any traction forces that may be reported [16]. Limitations also arise from the difficulty associated with creating systems that recapitulate the *in vivo* microenvironment and that have well-defined mechanical properties. For example, a combination of spatial heterogeneities and large pore sizes in collagen gels can result in a breakdown of the continuous medium assumption [140]. Thus, the relevant mechanical properties of collagen are difficult to determine because of discrepancies between bulk rheology and micromechanics of the collagen network [140]. Furthermore, discrete fiducial markers cannot fully capture the deformation of a continuous substratum, and any deformations of the matrix in between markers remain uncertain [64,65]. Lastly, obtaining high resolution images can be expensive and may limit the accessibility of the technology.

A recent alternative for measuring substratum deformation involves imaging deformations of fibrous matrix materials as opposed to embedded beads [141,142]. These studies offer a new approach for calculating 3D displacement fields that may be advantageous in systems where bead displacement techniques are not feasible or have limited accuracy. For example, during remodeling of a fibrous matrix, reflection microscopy can be used to calculate matrix displacements and obtain qualitative information on cell tractions [141,142]. Quantitative cell traction calculations rely on the assumption that the fibrous network is isotropic and can be approximated as linearly elastic [141]. In addition, molecular force sensors offer one strategy for improving the spatial resolution of quantified traction forces [143–147]. In contrast to TFM-based approaches that can only quantify traction forces exerted by the cell on its external microenvironment, molecular force sensors can be used to probe internal cellular forces [100]. One example is molecular tension-based fluorescence microscopy, which uses immobilized DNA hairpins that have a tunable force threshold to measure tension across integrins [148]. Another example uses a combination of super-resolution light microscopy and polypeptide fluorescence resonance energy transfer (FRET)-based tension sensors to visualize forces across individual molecules that are present within focal adhesions [149]. Myosin I has been used as a sensor for molecular force [150], and DNA hairpins have been used to probe traction forces with high spatial resolution [151]. A final example consists of a molecular tension sensor that uses a PEG polymer to create spatiotemporal maps of mechanical forces exerted by cell-surface receptors [152]. This tension sensor has well characterized mechanical properties, is biocompatible, and shows minimal nonspecific interactions [152]. That being said, a number of challenges must be addressed, including *in vivo* calibration [145], improved understanding of the relationship

between measurements in culture and *in vivo*, and improved understanding of the appropriate simplifying assumptions [100].

Additional information about cell-generated mechanical forces can also be obtained by combining TFM with other experimental techniques. For example, AFM, which can be used to directly measure the mechanical properties of cells, has been combined with TFM to study how cells respond to applied forces [153,154]. In a similar approach, incorporating magnetic particles within silicone micropillars enables one to study the response of cells to precisely applied forces [155]. Cellular traction forces have also been studied by combining TFM with optical tweezers. In particular, optical tweezers have been used to dictate the placement of beads so that traction forces can be quantified locally [156,157]. It has also been suggested that integrating TFM with molecular force sensors may provide a way to understand the interplay between extrinsic and intrinsic cellular forces [100]. While the aforementioned techniques have made improvements to conventional TFM, developing a complete understanding of traction forces in biological systems will require versatile techniques that can produce spatial maps of forces *in vivo* with high accuracy and precision.

Concluding remarks

Insight into the biophysical and biochemical mechanisms through which traction forces act is required for understanding a vast array of biological processes. Here, we describe experimental techniques and computational methods to quantify traction forces. Despite considerable progress and improvement in experimental and computational approaches, current TFM techniques cannot be used to quantify traction forces exerted within *in vivo* microenvironments, nor can they account for the complex heterogeneity present in native ECMs. As described here, microfabricated tissues provide an experimentally tractable means to quantify the traction forces exerted by and within multicellular cohorts. Coupled with improved computational approaches, microfabricated tissues will shed light on the tractions exerted by cells in heterogeneous nonlinear microenvironments and provide valuable insight into processes that resemble *in vivo* migrational and morphogenetic movements.

Acknowledgments

Work from the authors' group was supported in part by grants from the NIH (GM083997, HL110335, HL118532, HL120142, CA 187692), the NSF (CMMI-1435853), the David & Lucile Packard Foundation, the Alfred P. Sloan Foundation, the Camille & Henry Dreyfus Foundation, and the Burroughs Wellcome Fund. M.J.S. was supported in part by the NSF Graduate Research Fellowship Program. C.M.N. was supported in part by a Faculty Scholars Award from the Howard Hughes Medical Institute.

Abbreviations

2D	two-dimensional
3D	three-dimensional
AFM	atomic force microscopy
ECM	extracellular matrix
FRET	fluorescence resonance energy transfer

MDCK	Madin–Darby canine kidney
PA	polyacrylamide
PEG	polyethylene glycol
TFM	traction force microscopy

References

1. Maniotis AJ, Chen CS, Ingber DE. Demonstration of mechanical connections between integrins, cytoskeletal filaments, and nucleoplasm that stabilize nuclear structure. *Proc Natl Acad Sci USA*. 1997; 94(3):849–854. [PubMed: 9023345]
2. Thery M, Jimenez-Dalmaroni A, Racine V, Bornens M, Julicher F. Experimental and theoretical study of mitotic spindle orientation. *Nature*. 2007; 447(7143):493–496. [PubMed: 17495931]
3. Chen CS, Mrksich M, Huang S, Whitesides GM, Ingber DE. Geometric Control of Cell Life and Death. *Science*. 1997; 276(5317):1425–1428. [PubMed: 9162012]
4. Engler AJ, Sen S, Sweeney HL, Discher DE. Matrix Elasticity Directs Stem Cell Lineage Specification. *Cell*. 2006; 126(4):677–689. [PubMed: 16923388]
5. Desprat N, Supatto W, Pouille P-A, Beaurepaire E, Farge E. Tissue Deformation Modulates Twist Expression to Determine Anterior Midgut Differentiation in *Drosophila* Embryos. *Dev Cell*. 2008; 15(3):470–477. [PubMed: 18804441]
6. Farge E. Mechanical Induction of Twist in the *Drosophila* Foregut/Stomodaeal Primordium. *Curr Biol*. 2003; 13(16):1365–1377. [PubMed: 12932320]
7. Hinz B, Mastrangelo D, Iselin CE, Chaponnier C, Gabbiani G. Mechanical Tension Controls Granulation Tissue Contractile Activity and Myofibroblast Differentiation. *Am J Pathol*. 2001; 159(3):1009–1020. [PubMed: 11549593]
8. Munevar S, Wang Y, Dembo M. Traction force microscopy of migrating normal and H-ras transformed 3T3 fibroblasts. *Biophys J*. 2001; 80(4):1744–1757. [PubMed: 11259288]
9. Peschetola V, Laurent VM, Duperray A, Michel R, Ambrosi D, Preziosi L, Verdier C. Time-dependent traction force microscopy for cancer cells as a measure of invasiveness. (1949-3592 (Electronic)). 2013
10. Koch TM, Münster S, Bonakdar N, Butler JP, Fabry B. 3D Traction Forces in Cancer Cell Invasion. *PLoS ONE*. 2012; 7(3):e33476. [PubMed: 22479403]
11. Reffay M, Parrini MC, Cochet-Escartin O, Ladoux B, Buguin A, Coscoy S, Amblard F, Camonis J, Silberzan P. Interplay of RhoA and mechanical forces in collective cell migration driven by leader cells. *Nat Cell Biol*. 2014; 16(3):217–223. [PubMed: 24561621]
12. Treppe X, Wasserman MR, Angelini TE, Millet E, Weitz DA, Butler JP, Fredberg JJ. Physical forces during collective cell migration. *Nat Phys*. 2009; 5(6):426–430.
13. Heisenberg C-P, Bellaïche Y. Forces in Tissue Morphogenesis and Patterning. *Cell*. 2013; 153(5):948–962. [PubMed: 23706734]
14. Gov NS. Traction forces during collective cell motion. *HFSP Journal*. 2009; 3(4):223–227. [PubMed: 20119479]
15. His, W. Unsere Körperform und das physiologische Problem ihrer Entstehung Briefe an einen befreundeten Naturforscher / von Wilhelm His. Leipzig: F.C.W. Vogel; 1874.
16. Paluch EK, Nelson CM, Biais N, Fabry B, Moeller J, Pruitt BL, Wollnik C, Kudryasheva G, Rehfeldt F, Federle W. Mechanotransduction: use the force(s). *BMC Biol*. 2015; 13(1):1–14. [PubMed: 25555396]
17. Richards, RJ. The Tragic Sense of Life: Ernst Haeckel and the Struggle over Evolutionary Thought. Chicago: University of Chicago Press; 2008.
18. Maskarinec SA, Franck C, Tirrell DA, Ravichandran G. Quantifying cellular traction forces in three dimensions. *Proc Natl Acad Sci USA*. 2009; 106(52):22108–22113. [PubMed: 20018765]

19. Mok S, Moraes C. Thinking big by thinking small: advances in mechanobiology across the length scales. *Integr Biol.* 2016; 8(3):262–266.
20. Polacheck WJ, Chen CS. Measuring cell-generated forces: a guide to the available tools. *Nat Meth.* 2016; 13(5):415–423.
21. Style RW, Boltyskiy R, German GK, Hyland C, MacMinn CW, Mertz AF, Wilen LA, Xu Y, Dufresne ER. Traction force microscopy in physics and biology. *Soft Matter.* 2014; 10(23):4047–4055. [PubMed: 24740485]
22. Ribeiro AJS, Denisin AK, Wilson RE, Pruitt BL. For whom the cells pull: Hydrogel and micropost devices for measuring traction forces. *Methods.* 2016; 94:51–64. [PubMed: 26265073]
23. Gupta, M., Kocgozlu, L., Sarangi, BR., Margadant, F., Ashraf, M., Ladoux, B. Chapter 16 - Micropillar substrates: A tool for studying cell mechanobiology. In: Ewa, KP., editor. *Methods in Cell Biology.* Vol. 125. Academic Press; 2015. p. 289-308.
24. Nelson CM, Tien J. Microstructured extracellular matrices in tissue engineering and development. *Curr Opin Biotechnol.* 2006; 17(5):518–523. [PubMed: 16971111]
25. Tien J, Nelson CM. Microstructured Extracellular Matrices in Tissue Engineering and Development: An Update. *Ann Biomed Eng.* 2013; 42(7):1413–1423. [PubMed: 24030067]
26. Zorlutuna P, Annabi N, Camci-Unal G, Nikkhah M, Cha JM, Nichol JW, Manbachi A, Bae H, Chen S, Khademosseini A. Microfabricated Biomaterials for Engineering 3D Tissues. *Adv Mater.* 2012; 24(14):1782–1804. [PubMed: 22410857]
27. Harris AK, Wild P, Stopak D. Silicone rubber substrata: a new wrinkle in the study of cell locomotion. *Science.* 1980; 208(4440):177–179. [PubMed: 6987736]
28. Harris JK. A photoelastic substrate technique for dynamic measurements of forces exerted by moving organisms. *J Microsc.* 1978; 114(2):219–228. [PubMed: 739538]
29. Harris, AK. Locomotion of Tissue Cells, Ciba Foundation Symposium. Amsterdam: Elsevier; 1973. Cell surface movements related to cell locomotion; p. 3-26.
30. Dembo M, Wang Y-L. Stresses at the Cell-to-Substrate Interface during Locomotion of Fibroblasts. *Biophys J.* 1999; 76(4):2307–2316. [PubMed: 10096925]
31. Lee J, Leonard M, Oliver T, Ishihara A, Jacobson K. Traction forces generated by locomoting keratocytes. *J Cell Biol.* 1994; 127(6):1957–1964. [PubMed: 7806573]
32. Oliver T, Jacobson K, Dembo M. Traction forces in locomoting cells. *Cell Motil Cytoskeleton.* 1995; 31(3):225–240. [PubMed: 7585992]
33. Burton K, Taylor DL. Traction forces of cytokinesis measured with optically modified elastic substrata. *Nature.* 1997; 385(6615):450–454. [PubMed: 9009194]
34. Roy P, Petroll WM, Cavanagh HD, Chuong CJ, Jester JV. An in Vitro Force Measurement Assay to Study the Early Mechanical Interaction between Corneal Fibroblasts and Collagen Matrix. *Exp Cell Res.* 1997; 232(1):106–117. [PubMed: 9141627]
35. Brandley BK, Weisz OA, Schnaar RL. Cell attachment and long-term growth on derivatizable polyacrylamide surfaces. *J Biol Chem.* 1987; 262(13):6431–6437. [PubMed: 3571264]
36. Pelham RJ, Wang Y-L. Cell locomotion and focal adhesions are regulated by substrate flexibility. *Proc Natl Acad Sci USA.* 1997; 94(25):13661–13665. [PubMed: 9391082]
37. Wang, Y-L., Pelham, RJ, Jr. *Methods in Enzymology.* Vol. 298. Academic Press; 1998. Preparation of a flexible, porous polyacrylamide substrate for mechanical studies of cultured cells; p. 489-496.
38. Beningo KA, Wang Y-L. Flexible substrata for the detection of cellular traction forces. *Trends Cell Biol.* 2002; 12(2):79–84. [PubMed: 11849971]
39. Sabass B, Gardel ML, Waterman CM, Schwarz US. High Resolution Traction Force Microscopy Based on Experimental and Computational Advances. *Biophys J.* 2008; 94(1):207–220. [PubMed: 17827246]
40. Lautscham LA, Lin CY, Auernheimer V, Naumann CA, Goldmann WH, Fabry B. Biomembrane-mimicking lipid bilayer system as a mechanically tunable cell substrate. *Biomaterials.* 2014; 35(10):3198–3207. [PubMed: 24439398]
41. Tan JL, Tien J, Pirone DM, Gray DS, Bhadriraju K, Chen CS. Cells lying on a bed of microneedles: An approach to isolate mechanical force. *Proc Natl Acad Sci USA.* 2003; 100(4):1484–1489. [PubMed: 12552122]

42. Galbraith CG, Sheetz MP. A micromachined device provides a new bend on fibroblast traction forces. *Proc Natl Acad Sci USA*. 1997; 94(17):9114–9118. [PubMed: 9256444]
43. Balaban NQ, Schwarz US, Riveline D, Goichberg P, Tzur G, Sabanay I, Mahalu D, Safran S, Bershadsky A, Addadi L, Geiger B. Force and focal adhesion assembly: a close relationship studied using elastic micropatterned substrates. *Nat Cell Biol*. 2001; 3(5):466–472. [PubMed: 11331874]
44. Rodriguez ML, Graham BT, Pabon LM, Han SJ, Murry CE, Sniadecki NJ. Measuring the Contractile Forces of Human Induced Pluripotent Stem Cell-Derived Cardiomyocytes With Arrays of Microposts. *J Biomech Eng*. 2014; 136(5):051005–051005. [PubMed: 24615475]
45. du Roure O, Saez A, Buguin A, Austin RH, Chavrier P, Siberzan P, Ladoux B. Force mapping in epithelial cell migration. *Proc Natl Acad Sci USA*. 2005; 102(7):2390–2395. [PubMed: 15695588]
46. Schoen I, Hu W, Klotzsch E, Vogel V. Probing Cellular Traction Forces by Micropillar Arrays: Contribution of Substrate Warping to Pillar Deflection. *Nano Lett*. 2010; 10(5):1823–1830. [PubMed: 20387859]
47. Hur SS, Zhao Y, Li Y-S, Botvinick E, Chien S. Live Cells Exert 3-Dimensional Traction Forces on Their Substrata. *Cell Mol Bioeng*. 2009; 2(3):425–436. [PubMed: 19779633]
48. del Álamo JC., Meili, R., Álvarez-González, B., Alonso-Latorre, B., Bastounis, E., Firtel, R., Lasheras, JC. Three-Dimensional Quantification of Cellular Traction Forces and Mechanosensing of Thin Substrata by Fourier Traction Force Microscopy. *PLoS ONE*. 2013; 8(9):e69850. [PubMed: 24023712]
49. Legant WR, Choi CK, Miller JS, Shao L, Gao L, Betzig E, Chen CS. Multidimensional traction force microscopy reveals out-of-plane rotational moments about focal adhesions. *Proc Natl Acad Sci USA*. 2013; 110(3):881–886. [PubMed: 23277584]
50. Delanoë-Ayari H, Rieu JP, Sano M. 4D Traction Force Microscopy Reveals Asymmetric Cortical Forces in Migrating Dictyostelium Cells. *Phys Rev Lett*. 2010; 105(24):248103. [PubMed: 21231559]
51. Legant WR, Miller JS, Blakely BL, Cohen DM, Genin GM, Chen CS. Measurement of mechanical tractions exerted by cells in three-dimensional matrices. *Nat Meth*. 2010; 7(12):969–971.
52. Steinwachs J, Metzner C, Skodzek K, Lang N, Thievensen I, Mark C, Munster S, Aifantis KE, Fabry B. Three-dimensional force microscopy of cells in biopolymer networks. *Nat Meth*. 2016; 13(2):171–176.
53. Hall MS, Long R, Feng X, Huang Y, Hui C-Y, Wu M. Toward single cell traction microscopy within 3D collagen matrices. *Exp Cell Res*. 2013; 319(16):2396–2408. [PubMed: 23806281]
54. Gjorevski N, Piotrowski AS, Varner VD, Nelson CM. Dynamic tensile forces drive collective cell migration through three-dimensional extracellular matrices. *Sci Rep*. 2015; 5:11458. [PubMed: 26165921]
55. Gjorevski N, Nelson CM. Mapping of Mechanical Strains and Stresses around Quiescent Engineered Three-Dimensional Epithelial Tissues. *Biophys J*. 2012; 103(1):152–162. [PubMed: 22828342]
56. Piotrowski, AS., Varner, VD., Gjorevski, N., Nelson, CM. Three-Dimensional Traction Force Microscopy of Engineered Epithelial Tissues. In: Nelson, CM., editor. *Tissue Morphogenesis: Methods and Protocols*. New York, NY: Springer; 2015. p. 191-206.
57. Landau, LD., Lifshitz, EM. *Course of Theoretical Physics*. 3. Vol. 7. Moscow USSR: Elsevier Butterworth-Heinemann, Institute of Physical Problems, USSR Academy of Sciences; 1986. Theory of Elasticity.
58. Landau, LD., Lifshitz, EM. *Course of Theoretical Physics*. 3. Vol. 1. Moscow USSR: Elsevier Butterworth-Heinemann, Institute of Physical Problems, USSR Academy of Sciences; 1976. Mechanics.
59. Butler JP, Toli -Nørrelykke IM, Fabry B, Fredberg JJ. Traction fields, moments, and strain energy that cells exert on their surroundings. *Am J Physiol Cell Physiol*. 2002; 282(3):C595–C605. [PubMed: 11832345]
60. Ambrosi D. Cellular Traction as an Inverse Problem. *SIAM J Numer Anal*. 2006; 66(6):2049–2060.

61. Stout DA, Bar-Kochba E, Estrada JB, Toyjanova J, Kesari H, Reichner JS, Franck C. Mean deformation metrics for quantifying 3D cell-matrix interactions without requiring information about matrix material properties. *Proc Natl Acad Sci USA*. 2016; 113(11):2898–2903. [PubMed: 26929377]
62. Franck C, Hong S, Maskarinec SA, Tirrell DA, Ravichandran G. Three-dimensional Full-field Measurements of Large Deformations in Soft Materials Using Confocal Microscopy and Digital Volume Correlation. *Exp Mech*. 2007; 47(3):427–438.
63. Bar-Kochba E, Toyjanova J, Andrews E, Kim KS, Franck C. A Fast Iterative Digital Volume Correlation Algorithm for Large Deformations. *Exp Mech*. 2015; 55(1):261–274.
64. Dembo M, Oliver T, Ishihara A, Jacobson K. Imaging the traction stresses exerted by locomoting cells with the elastic substratum method. *Biophys J*. 1996; 70(4):2008–2022. [PubMed: 8785360]
65. Schwarz US, Balaban NQ, Riveline D, Bershadsky A, Geiger B, Safran SA. Calculation of Forces at Focal Adhesions from Elastic Substrate Data: The Effect of Localized Force and the Need for Regularization. *Biophys J*. 2002; 83(3):1380–1394. [PubMed: 12202364]
66. Hansen P. Rank-Deficient and Discrete Ill-Posed Problems. *Mathematical Modeling and Computation*. Society for Industrial and Applied Mathematics. 1998
67. Schwarz US, Soiné JRD. Traction force microscopy on soft elastic substrates: A guide to recent computational advances. *Biochim Biophys Acta*. 2015; 1853(11, Part B):3095–3104. [PubMed: 26026889]
68. Plotnikov, SV., Sabass, B., Schwarz, US., Waterman, CM. Chapter 20 - High-Resolution Traction Force Microscopy. In: Jennifer, CW., Torsten, W., editors. *Methods in Cell Biology*. Vol. 123. Academic Press; 2014. p. 367-394.
69. Vaughan RB, Trinkaus JP. Movements of Epithelial Cell Sheets In Vitro. *J Cell Sci*. 1966; 1(4): 407–413. [PubMed: 5956715]
70. Siedlik MJ, Varner VD, Nelson CM. Pushing, pulling, and squeezing our way to understanding mechanotransduction. *Methods*. 2016
71. Borghi N, Farge E, Lavelle C. Experimental approaches in mechanotransduction: From molecules to pathology. *Methods*. 2016; 94:1–3. [PubMed: 26896045]
72. Miller CJ, Davidson LA. The interplay between cell signalling and mechanics in developmental processes. *Nat Rev Genet*. 2013; 14(10):733–744. [PubMed: 24045690]
73. Molino D, Quignard S, Gruget C, Pincet F, Chen Y, Piel M, Fattaccioli J. On-Chip Quantitative Measurement of Mechanical Stresses During Cell Migration with Emulsion Droplets. *Sci Rep*. 2016; 6:29113. [PubMed: 27373558]
74. Mann C, Leckband D. Measuring Traction Forces in Long-Term Cell Cultures. *Cell Mol Bioeng*. 2010; 3(1):40–49.
75. Jannat RA, Dembo M, Hammer DA. Traction Forces of Neutrophils Migrating on Compliant Substrates. *Biophys J*. 2011; 101(3):575–584. [PubMed: 21806925]
76. Ricart BG, Yang MT, Hunter CA, Chen CS, Hammer DA. Measuring Traction Forces of Motile Dendritic Cells on Micropost Arrays. *Biophys J*. 2011; 101(11):2620–2628. [PubMed: 22261049]
77. Soon FC, Tee SK, Youseffi M, Denyer CM. Tracking Traction Force Changes of Single Cells on the Liquid Crystal Surface. *Biosensors*. 2015; 5(1):13–24. [PubMed: 25808839]
78. Engler A, Bacakova L, Newman C, Hategan A, Griffin M, Discher D. Substrate Compliance versus Ligand Density in Cell on Gel Responses. *Biophys J*. 2004; 86(1):617–628. [PubMed: 14695306]
79. Reinhart-King CA, Dembo M, Hammer DA. The Dynamics and Mechanics of Endothelial Cell Spreading. *Biophys J*. 2005; 89(1):676–689. [PubMed: 15849250]
80. Maloney JM, Walton EB, Bruce CM, Van Vliet KJ. Influence of finite thickness and stiffness on cellular adhesion-induced deformation of compliant substrata. *Phys Rev E*. 2008; 78(4):041923.
81. Zhou DW, García AJ. Measurement Systems for Cell Adhesive Forces. *J Biomech Eng*. 2015; 137(2):020908–020908. [PubMed: 25416835]
82. Ruder WC, LeDuc PR. Cells gain traction in 3D. *Proc Natl Acad Sci USA*. 2012; 109(28):11060–11061. [PubMed: 22778421]

83. Tambe DT, Corey Hardin C, Angelini TE, Rajendran K, Park CY, Serra-Picamal X, Zhou EH, Zaman MH, Butler JP, Weitz DA, Fredberg JJ, Trepas X. Collective cell guidance by cooperative intercellular forces. *Nat Mater.* 2011; 10(6):469–475. [PubMed: 21602808]
84. Friedl P, Wolf K, Zegers MM. Rho-directed forces in collective migration. *Nat Cell Biol.* 2014; 16(3):208–210. [PubMed: 24576897]
85. Tambe DT, Crouette U, Trepas X, Park CY, Kim JH, Millet E, Butler JP, Fredberg JJ. Monolayer Stress Microscopy: Limitations, Artifacts, and Accuracy of Recovered Intercellular Stresses. *PLoS ONE.* 2013; 8(2):e55172. [PubMed: 23468843]
86. Murray JD, Oster GF. Cell traction models for generating pattern and form in morphogenesis. *J Math Biol.* 1984; 19(3):265–279. [PubMed: 6470581]
87. Szabó A, Mayor R. Cell traction in collective cell migration and morphogenesis: The chase and run mechanism. *Cell Adh Migr.* 2015; 9(5):380–383. [PubMed: 26267782]
88. Varner VD, Nelson CM. Cellular and physical mechanisms of branching morphogenesis. *Development.* 2014; 141(14):2750–2759. [PubMed: 25005470]
89. Siedlik MJ, Nelson CM. Regulation of tissue morphodynamics: an important role for actomyosin contractility. *Curr Opin Genet Dev.* 2015; 32:80–85. [PubMed: 25748251]
90. Thompson, DAW. *On Growth and Form.* Cambridge: Cambridge University Press; 1917.
91. Singhvi R, Kumar A, Lopez GP, Stephanopoulos GN, Wang DI, Whitesides GM, Ingber DE. Engineering cell shape and function. *Science.* 1994; 264(5159):696–698. [PubMed: 8171320]
92. Huang S, Ingber DE. The structural and mechanical complexity of cell-growth control. *Nat Cell Biol.* 1999; 1(5):E131–E138. [PubMed: 10559956]
93. Nelson CM, Jean RP, Tan JL, Liu WF, Sniadecki NJ, Spector AA, Chen CS. Emergent patterns of growth controlled by multicellular form and mechanics. *Proc Natl Acad Sci USA.* 2005; 102(33):11594–11599. [PubMed: 16049098]
94. Chanet, S., Martin, AC. Chapter Thirteen - Mechanical Force Sensing in Tissues. In: Adam, JE., Sanjay, K., editors. *Progress in Molecular Biology and Translational Science.* Vol. 126. Academic Press; 2014. p. 317-352.
95. Hutson MS, Tokutake Y, Chang M-S, Bloor JW, Venakides S, Kiehart DP, Edwards GS. Forces for Morphogenesis Investigated with Laser Microsurgery and Quantitative Modeling. *Science.* 2003; 300(5616):145–149. [PubMed: 12574496]
96. Campas O, Mammoto T, Hasso S, Sperling RA, O'Connell D, Bischof AG, Maas R, Weitz DA, Mahadevan L, Ingber DE. Quantifying cell-generated mechanical forces within living embryonic tissues. *Nat Meth.* 2014; 11(2):183–189.
97. Rauzi, M., Lenne, PF. Probing Cell Mechanics with Subcellular Laser Dissection of Actomyosin Networks in the Early Developing *Drosophila* Embryo. In: Nelson, CM., editor. *Tissue Morphogenesis: Methods and Protocols.* New York, NY: Springer; 2015. p. 209-218.
98. Behrndt M, Salbreux G, Campinho P, Hauschild R, Oswald F, Roensch J, Grill SW, Heisenberg C-P. Forces Driving Epithelial Spreading in Zebrafish Gastrulation. *Science.* 2012; 338(6104):257–260. [PubMed: 23066079]
99. Rauzi M, Verant P, Lecuit T, Lenne P-F. Nature and anisotropy of cortical forces orienting *Drosophila* tissue morphogenesis. *Nat Cell Biol.* 2008; 10(12):1401–1410. [PubMed: 18978783]
100. Varner VD, Nelson CM. Let's push things forward: disruptive technologies and the mechanics of tissue assembly. *Integr Biol.* 2013; 5(9):1162–1173.
101. Farge, E. Chapter eight - Mechanotransduction in Development. In: Michel, L., editor. *Current Topics in Developmental Biology.* Vol. 95. Academic Press; 2011. p. 243-265.
102. Mammoto T, Ingber DE. Mechanical control of tissue and organ development. *Development (Cambridge, England).* 2010; 137(9):1407–1420.
103. Rauzi, M., Lenne, P-F. Chapter four - Cortical Forces in Cell Shape Changes and Tissue Morphogenesis. In: Michel, L., editor. *Current Topics in Developmental Biology.* Vol. 95. Academic Press; 2011. p. 93-144.
104. Mooney JF, Hunt AJ, McIntosh JR, Libenko CA, Walba DM, Rogers CT. Patterning of functional antibodies and other proteins by photolithography of silane monolayers. *Proc Natl Acad Sci USA.* 1996; 93(22):12287–12291. [PubMed: 8901573]

105. Alom Ruiz S, Chen CS. Microcontact printing: A tool to pattern. *Soft Matter*. 2007; 3(2):168–177.
106. Shen K, Qi J, Kam LC. Microcontact Printing of Proteins for Cell Biology. *JoVE*. 2008; (22):e1065.
107. Nelson CM, Inman JL, Bissell MJ. Three-dimensional lithographically defined organotypic tissue arrays for quantitative analysis of morphogenesis and neoplastic progression. *Nat Protocols*. 2008; 3(4):674–678. [PubMed: 18388950]
108. Piotrowski-Daspit AS, Tien J, Nelson CM. Interstitial fluid pressure regulates collective invasion in engineered human breast tumors via Snail, vimentin, and E-cadherin. *Integr Biol*. 2016; 8(3): 319–331.
109. Xu T, Jin J, Gregory C, Hickman JJ, Boland T. Inkjet printing of viable mammalian cells. *Biomaterials*. 2005; 26(1):93–99. [PubMed: 15193884]
110. Murphy SV, Atala A. 3D bioprinting of tissues and organs. *Nat Biotech*. 2014; 32(8):773–785.
111. Tien J, Nelson CM, Chen CS. Fabrication of aligned microstructures with a single elastomeric stamp. *Proc Natl Acad Sci USA*. 2002; 99(4):1758–1762. [PubMed: 11842197]
112. Pavlovich AL, Manivannan S, Nelson CM. Adipose Stroma Induces Branching Morphogenesis of Engineered Epithelial Tubules. *Tissue Eng Part A*. 2010; 16(12):3719–3726. [PubMed: 20649458]
113. Boghaert E, Gleghorn JP, Lee K, Gjorevski N, Radisky DC, Nelson CM. Host epithelial geometry regulates breast cancer cell invasiveness. *Proc Natl Acad Sci USA*. 2012; 109(48):19632–19637. [PubMed: 23150585]
114. Bernard A, Renault JP, Michel B, Bosshard HR, Delamarche E. Microcontact Printing of Proteins. *Adv Mater*. 2000; 12(14):1067–1070.
115. Wilbur JL, Kumar A, Kim E, Whitesides GM. Microfabrication by microcontact printing of self-assembled monolayers. *Adv Mater*. 1994; 6(7-8):600–604.
116. Nelson CM, Raghavan S, Tan JL, Chen CS. Degradation of Micropatterned Surfaces by Cell-Dependent and -Independent Processes. *Langmuir*. 2003; 19(5):1493–1499.
117. Lee J, Abdeen AA, Tang X, Saif TA, Kilian KA. Geometric guidance of integrin mediated traction stress during stem cell differentiation. *Biomaterials*. 2015; 69:174–183. [PubMed: 26285084]
118. Polio SR., Smith, ML. Chapter 2 - Patterned Hydrogels for Simplified Measurement of Cell Traction Forces. In: Matthieu, P., Manuel, T., editors. *Methods in Cell Biology*. Vol. 121. Academic Press; 2014. p. 17-31.
119. Polio SR, Parameswaran H, Canovic EP, Gaut CM, Aksyonova D, Stamenovic D, Smith ML. Topographical control of multiple cell adhesion molecules for traction force microscopy. *Integr Biol*. 2014; 6(3):357–365.
120. Vedula SRK, Leong MC, Lai TL, Hersen P, Kabla AJ, Lim CT, Ladoux B. Emerging modes of collective cell migration induced by geometrical constraints. *Proc Natl Acad Sci USA*. 2012; 109(32):12974–12979. [PubMed: 22814373]
121. Yang MT, Fu J, Wang Y-K, Desai RA, Chen CS. Assaying stem cell mechanobiology on microfabricated elastomeric substrates with geometrically modulated rigidity. *Nat Protocols*. 2011; 6(2):187–213. [PubMed: 21293460]
122. Han SJ, Bielawski KS, Ting LH, Rodriguez ML, Sniadecki NJ. Decoupling Substrate Stiffness, Spread Area, and Micropost Density: A Close Spatial Relationship between Traction Forces and Focal Adhesions. *Biophys J*. 2012; 103(4):640–648. [PubMed: 22947925]
123. Wang N, Ostuni E, Whitesides GM, Ingber DE. Micropatterning tractional forces in living cells. *Cell Motil Cytoskeleton*. 2002; 52(2):97–106. [PubMed: 12112152]
124. Gomez EW, Chen QK, Gjorevski N, Nelson CM. Tissue geometry patterns epithelial-mesenchymal transition via intercellular mechanotransduction. *J Cell Biochem*. 2010; 110(1):44–51. [PubMed: 20336666]
125. Rolli CG, Nakayama H, Yamaguchi K, Spatz JP, Kemkemer R, Nakanishi J. Switchable adhesive substrates: Revealing geometry dependence in collective cell behavior. *Biomaterials*. 2012; 33(8): 2409–2418. [PubMed: 22197568]

126. Raghavan S, Desai RA, Kwon Y, Mrksich M, Chen CS. Micropatterned Dynamically Adhesive Substrates for Cell Migration. *Langmuir*. 2010; 26(22):17733–17738. [PubMed: 20886900]
127. Jiang X, Ferrigno R, Mrksich M, Whitesides GM. Electrochemical Desorption of Self-Assembled Monolayers Noninvasively Releases Patterned Cells from Geometrical Confinements. *J Am Chem Soc*. 2003; 125(9):2366–2367. [PubMed: 12603104]
128. Gomez, EW., Nelson, CM. Lithographically Defined Two- and Three-Dimensional Tissue Microarrays. In: Khademhosseini, A.Suh, KY., Zourob, M., editors. *Biological Microarrays: Methods and Protocols*. Totowa, NJ: Humana Press; 2011. p. 107-116.
129. Manivannan, S., Gleghorn, JP., Nelson, CM. Engineered Tissues to Quantify Collective Cell Migration During Morphogenesis. In: Michos, O., editor. *Kidney Development: Methods and Protocols*. Totowa, NJ: Humana Press; 2012. p. 173-182.
130. Zhu W, Nelson CM. PI3K regulates branch initiation and extension of cultured mammary epithelia via Akt and Rac1 respectively. *Dev Biol*. 2013; 379(2):235–245. [PubMed: 23665174]
131. Lee K, Gjorevski N, Boghaert E, Radisky DC, Nelson CM. Snail1, Snail2, and E47 promote mammary epithelial branching morphogenesis. *EMBO J*. 2011; 30(13):2662–2674. [PubMed: 21610693]
132. Pavlovich AL, Boghaert E, Nelson CM. Mammary branch initiation and extension are inhibited by separate pathways downstream of TGF β in culture. *Exp Cell Res*. 2011; 317(13):1872–1884. [PubMed: 21459084]
133. Nelson CM, VanDuijn MM, Inman JL, Fletcher DA, Bissell MJ. Tissue Geometry Determines Sites of Mammary Branching Morphogenesis in Organotypic Cultures. *Science*. 2006; 314(5797):298–300. [PubMed: 17038622]
134. Gjorevski N, Nelson CM. Endogenous patterns of mechanical stress are required for branching morphogenesis. *Integr Biol*. 2010; 2(9):424–434.
135. Legant WR, Chen CS, Vogel V. Force-induced fibronectin assembly and matrix remodeling in a 3D microtissue model of tissue morphogenesis. *Integr Biol*. 2012; 4(10):1164–1174.
136. Legant WR, Pathak A, Yang MT, Deshpande VS, McMeeking RM, Chen CS. Microfabricated tissue gauges to measure and manipulate forces from 3D microtissues. *Proc Natl Acad Sci USA*. 2009; 106(25):10097–10102. [PubMed: 19541627]
137. Sakar MS, Eyckmans J, Pieters R, Eberli D, Nelson BJ, Chen CS. Cellular forces and matrix assembly coordinate fibrous tissue repair. *Nat Commun*. 2016; 7:11036. [PubMed: 26980715]
138. Mori H, Gjorevski N, Inman JL, Bissell MJ, Nelson CM. Self-organization of engineered epithelial tubules by differential cellular motility. *Proc Natl Acad Sci USA*. 2009; 106(35):14890–14895. [PubMed: 19706461]
139. Théry M. Micropatterning as a tool to decipher cell morphogenesis and functions. *J Cell Sci*. 2010; 123(24):4201. [PubMed: 21123618]
140. Jones CAR, Cibula M, Feng J, Krnacik EA, McIntyre DH, Levine H, Sun B. Micromechanics of cellularized biopolymer networks. *Proc Natl Acad Sci USA*. 2015; 112(37):E5117–E5122. [PubMed: 26324923]
141. Kim J, Jones CAR, Groves NS, Sun B. Three-Dimensional Reflectance Traction Microscopy. *PLoS ONE*. 2016; 11(6):e0156797. [PubMed: 27304456]
142. Notbohm J, Lesman A, Tirrell DA, Ravichandran G. Quantifying cell-induced matrix deformation in three dimensions based on imaging matrix fibers. *Integr Biol*. 2015; 7(10):1186–1195.
143. Cost AL, Ringer P, Chrostek-Grashoff A, Grashoff C. How to Measure Molecular Forces in Cells: A Guide to Evaluating Genetically-Encoded FRET-Based Tension Sensors. *Cell Mol Bioeng*. 2015; 8(1):96–105. [PubMed: 25798203]
144. Gayrard C, Borghi N. FRET-based Molecular Tension Microscopy. *Methods*. 2016; 94:33–42. [PubMed: 26210398]
145. de Souza N. Pulling on single molecules. *Nat Meth*. 2012; 9(9):873–877.
146. Liu B, Kim T-J, Wang Y. Live cell imaging of mechanotransduction. *J R Soc Interface*. 2010; 7(Suppl 3):S365–S375. [PubMed: 20356874]
147. Leidel C, Longoria RA, Gutierrez FM, Shubeita GT. Measuring Molecular Motor Forces In Vivo: Implications for Tug-of-War Models of Bidirectional Transport. *Biophys J*. 2012; 103(3):492–500. [PubMed: 22947865]

148. Zhang Y, Ge C, Zhu C, Salaita K. DNA-based digital tension probes reveal integrin forces during early cell adhesion. *Nat Commun.* 2014; 5
149. Morimatsu M, Mekhdjian AH, Chang AC, Tan SJ, Dunn AR. Visualizing the Interior Architecture of Focal Adhesions with High-Resolution Traction Maps. *Nano Lett.* 2015; 15(4):2220–2228. [PubMed: 25730141]
150. Laakso JM, Lewis JH, Shuman H, Ostap EM. Myosin I can act as a molecular force sensor. *Science.* 2008; 321(5885):133–136. [PubMed: 18599791]
151. Blakely BL, Dumelin CE, Trappmann B, McGregor LM, Choi CK, Anthony PC, Duisterberg VK, Baker BM, Block SM, Liu DR, Chen CS. A DNA-based molecular probe for optically reporting cellular traction forces. *Nat Meth.* 2014; 11(12):1229–1232.
152. Stabley DR, Jurchenko C, Marshall SS, Salaita KS. Visualizing mechanical tension across membrane receptors with a fluorescent sensor. *Nat Meth.* 2012; 9(1):64–67.
153. Zeinab A-R, Andrew EP. Cross talk between matrix elasticity and mechanical force regulates myoblast traction dynamics. *Phys Biol.* 2013; 10(6):066003. [PubMed: 24164970]
154. Haase K, Pelling AE. Investigating cell mechanics with atomic force microscopy. *J R Soc Interface.* 2015; 12(104):20140970. [PubMed: 25589563]
155. Sniadecki NJ, Anguelouch A, Yang MT, Lamb CM, Liu Z, Kirschner SB, Liu Y, Reich DH, Chen CS. Magnetic microposts as an approach to apply forces to living cells. *Proc Natl Acad Sci USA.* 2007; 104(37):14553–14558. [PubMed: 17804810]
156. Mejean CO, Schaefer AW, Millman EA, Forscher P, Dufresne ER. Multiplexed force measurements on live cells with holographic optical tweezers. *Opt Express.* 2009; 17(8):6209–6217. [PubMed: 19365444]
157. van der Horst A, Forde NR. Calibration of dynamic holographic optical tweezers for force measurements on biomaterials. *Opt Express.* 2008; 16(25):20987–21003. [PubMed: 19065239]
158. Ladoux B. Biophysics: Cells guided on their journey. *Nat Phys.* 2009; 5(6):377–378.
159. Franck C, Maskarinec SA, Tirrell DA, Ravichandran G. Three-Dimensional Traction Force Microscopy: A New Tool for Quantifying Cell-Matrix Interactions. *PLoS ONE.* 2011; 6(3):e17833. [PubMed: 21468318]
160. Rape AD, Guo W-H, Wang Y-L. The regulation of traction force in relation to cell shape and focal adhesions. *Biomaterials.* 2011; 32(8):2043–2051. [PubMed: 21163521]
161. Polio SR, Rothenberg KE, Stamenovi D, Smith ML. A micropatterning and image processing approach to simplify measurement of cellular traction forces. *Acta Biomater.* 2012; 8(1):82–88. [PubMed: 21884832]
162. Cvetkovic C, Raman R, Chan V, Williams BJ, Tolish M, Bajaj P, Sakar MS, Asada HH, Saif MTA, Bashir R. Three-dimensionally printed biological machines powered by skeletal muscle. *Proc Natl Acad Sci USA.* 2014; 111(28):10125–10130. [PubMed: 24982152]
163. Sung JH, Yu J, Luo D, Shuler ML, March JC. Microscale 3-D hydrogel scaffold for biomimetic gastrointestinal (GI) tract model. *Lab Chip.* 2011; 11(3):389–392. [PubMed: 21157619]

Figure 1. Single cell and multicellular traction forces in 2D and 3D systems

Traction forces exerted by A) single cell migrating on a 2D substratum, B) population of cells migrating collectively on a 2D substratum, C) single cell migrating in a 3D matrix, and D) population of cells migrating collectively in a 3D matrix. Cell-generated traction forces displace fiducial markers embedded in the substratum or matrix as cells migrate. Traction forces can be quantified by measuring the displacement of beads. Inspired by [158,20].

Figure 2. Visualizing traction force quantification

A) In-plane (top) and normal (bottom) traction forces calculated for a single cell migrating on a 2D substratum. Figure from [159], licensed under Creative Commons. B) In-plane (top) and normal (bottom) traction stresses for population of cells migrating collectively on a 2D substratum. Note that the image size is 750 μm by 750 μm. Reprinted by permission from Macmillan Publishers Ltd: Nature Physics [12], copyright (2009). C) Traction stress map for single cell migrating in a 3D matrix. Reprinted by permission from Macmillan Publishers Ltd: Nature Methods [51], copyright (2010). D) Traction stress map for population of cells

migrating collectively in a 3D matrix. Figure from [54], licensed under Creative Commons.
Scale bar is 50 μm .

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 3. Summary of methods to produce 2D and 3D microfabricated tissues

A) 2D microfabricated tissues can be produced by i) microcontact printing or ii) stenciling. When fabricated on micropost arrays or PA gels, these tissues can be used to quantify the traction forces involved in migratory and morphogenetic processes. The top-right image depicts microcontact-printed fibronectin on micropillars during a study of collective migration [120], the bottom-right image depicts an engineered epithelial monolayer produced on a PA gel via stenciling [124]. Top-right image adapted with permission from Vedula et al [120]. Copyright 2012, the National Academy of Sciences. Bottom-right image

adapted from Gomez et al [124] and reprinted with permission from Wiley-Liss Inc, copyright 2010. B) Microfabricated tissues fully encapsulated within a 3D ECM microenvironment can be produced by iii) replica micromolding or iv) self-assembly into tissue gauges. Replica micromolding produces wells within an ECM gel, into which cells are added and allowed to form 3D tissues. Traction forces exerted by these tissues are calculated from the movement of fluorescent beads that have been incorporated into the surrounding gel (iii, right image). Additionally, tissue gauges may be used to elucidate how levels of tension relate to tissue-level processes. These 3D microfabricated tissues self-assemble as a cell-laden ECM contracts against silicone micropillars. Top-right image adapted from Gjorevski et al [54], licensed under Creative Commons. Bottom-right image adapted with permission from Legant et al [136]. Copyright 2009, the National Academy of Sciences. Scale bars represent i) 20 μm , ii) 25 μm , iii) 50 μm , iv) 100 μm .

Table 1

Summary of microfabricated structures for quantifying traction forces.

Microfabrication technique	Microstructure	Dimension	References
Microcontact printing	Islands of ECM	2D	<ul style="list-style-type: none"> - NIH 3T3 fibroblasts [119,160,161] - MDCK epithelial cells [120] - Mesenchymal stem cells [117]
Stenciling	Islands of ECM	2D	<ul style="list-style-type: none"> - Human airway smooth muscle cells [123] - SCp2 mouse mammary epithelial cells [124]
Replica micromolding	Multidimensional structures	3D	<ul style="list-style-type: none"> - EpH4 mouse mammary epithelial cells [55,54,134] - EpH4 mouse mammary epithelial cells and primary organoids [133]
Tissue gauges	Patterned surfaces and structures	3D	<ul style="list-style-type: none"> - NIH 3T3 fibroblasts and neonatal rat cardiomyocytes [136] - NIH 3T3 fibroblasts [135,137]
3D printing	Patterned surfaces and structures	3D	<ul style="list-style-type: none"> - C2C12 skeletal muscle myoblasts [162]
Sacrificial molds	Multidimensional structures	3D	<ul style="list-style-type: none"> - Caco-2 colon carcinoma cells [163]