Difficult urethral catheterisation made safer: visualising the 'blind' hydrophilic wire

S Tadtayev, H Gresty, D Manson-Bahr

Royal Free London NHS Foundation Trust, UK

CORRESPONDENCE TO

Sergey Tadtayev, E: s.tadtayev@gmail.com
doi 10.1308/rcsann.2016.0213

Mandibular sagittal split osteotomy – A modified technique to reduce postoperative labiomental paraesthesia

M Manisali, FB Naini

St George's Hospital Medical School, London, UK

CORRESPONDENCE TO

Farhad B. Naini, E: Farhad.Naini@yahoo.co.uk
doi 10.1308/rcsann.2016.0220

BACKGROUND

Difficult urethral catheterisation in a male, non-trauma patient with urinary retention is often due to a false passage, urethral stricture or bladder neck contracture. When simple measures¹ have failed, the usual choice is between use of a suprapubic catheter and cystoscopic catheterisation over a guidewire. However, suprapubic catheterisation has its contraindications and the wait for theatre can be long. An alternative strategy is the blind passage of the hydrophilic wire into the bladder in the emergency department,^{2–5} which is not without risk as the guidewire can end up in the false passage, ejaculatory duct or ureter. We advocate ultrasonography visualisation of the "blind" wire, which makes this approach safer.

TECHNIQUE

Following the instillation of lubricating gel, a Sensor™ (Boston Scientific, Marlborough, MA, US) wire is passed in the urethra, its advance monitored using ultrasonography (abdominal transducer pointing over the bladder). Once the sufficient length of the wire is in the bladder, the operator can either pass an open tip 16–18Fr Foley catheter over the wire (if false passage is suspected) or perform dilation with S-Curve™ (Cook Medical, Bloomington, IN, US) urethral dilators (in the case of urethral stricture or bladder neck contracture) prior to catheterisation. If dilation is not tolerated, a temporary 6Fr ureteric catheter or 5/8Fr infant feeding tube can be passed instead.

DISCUSSION

The ultrasonography machine is now an indispensable part of the equipment of many emergency departments. The echogenicity of the guidewires makes it easy to ultrasonically visualise their passage into the full bladder. In our hands, this technique proved to be safe and effective.

References

1. Tadtayev S, Mazaris E. Difficult urethral catheterisation made easy: validation and the new application of the established technique. *Ann R Coll Surg Engl* 2013; **95**: 300.
2. Abbott JE, Heinemann A, Badalament R, Davalos JG. A clever technique for placement of a urinary catheter over a wire. *Urol Ann* 2015; **7**: 367–370.
3. Olson L, Esler R, Gunendran T. Blind hydrophilic guidewire for difficult urethral catheterisation. *Ann R Coll Surg Engl* 2014; **96**: 629.
4. Blitz BF. A simple method using hydrophilic guide wires for the difficult urethral catheterization. *Urology* 1995; **46**: 99–100.
5. Zammit PA, German K. The difficult urethral catheterization: use of a hydrophilic guidewire. *BJU Int* 2004; **93**: 883–884.

Sagittal split mandibular ramus osteotomy is one of the most frequently employed surgical procedures in orthognathic surgery, and is used to advance, set-back and rotate the distal (tooth-bearing) segment of the mandible. The original technique, described by Trauner and Obwegeser,¹ has seen a number of modifications.^{2,3}

The principal complication associated with the technique and its described modifications remains temporary or permanent neurosensory deficit of the lower lip and chin. Reports in the literature on the frequency of this complication vary quite significantly, but reach up to 85%.⁴

TECHNIQUE

The suggested modification in technique is that the vertical osteotomy along the anterior border of the mandibular ramus is made lateral to

Figure 1 Oblique view of the right mandibular ramus, demonstrating the more lateral position of the modified sagittal split osteotomy (solid line) in relation to the traditional position (dotted line)

Figure 2 Anterior view of the right mandibular ramus

the traditionally described position. It is essential to make the osteotomy as far lateral as possible, usually approximately 3–4mm more lateral than the traditional position (Figs 1 and 2). The mandibular nerve is thereby much more likely to remain in the distal segment of the osteotomised mandible.

When creating the sagittal split, the anterior aspect of the proximal (ramus) segment will be quite thin following this modified osteotomy. It is essential to use light force, while gently spreading the segments apart, using two thin osteotomes to avoid fracturing the antero-lateral aspect of the mandibular ramus.

DISCUSSION

This technical modification has reduced the incidence of postoperative labiomental paraesthesia by over 60% (at postoperative day one) in our practice. The efficacy of the technique now needs to be assessed in a randomised controlled clinical trial.

References

1. Trauner R, Obwegeser H. The surgical correction of mandibular prognathism and retrognathia with consideration of genioplasty. I. Surgical procedures to correct mandibular prognathism and reshaping of the chin. *Oral Surg Oral Med Oral Pathol* 1957; **10**: 677–89; contd.
2. Hunsuck EE. A modified intraoral sagittal splitting technic for correction of mandibular prognathism. *J Oral Surg* 1968; **26**: 250–253.
3. Epker BN. Modifications in the sagittal osteotomy of the mandible. *J Oral Surg* 1977; **35**: 157–159.
4. August M, Marchena J, Donady J *et al*. Neurosensory deficit and functional impairment after sagittal ramus osteotomy: a long-term follow-up study. *J Oral Maxillofac Surg* 1998; **56**: 1,231–1,235.

Open intraperitoneal onlay mesh repair for medium–large lateral abdominal wall herniae

L Hendra¹, A Shrestha², M Joshi², S Basu²

¹Isle of Wight Healthcare NHS Trust, UK

²East Kent Hospitals University NHS Foundation Trust

CORRESPONDENCE TO

Sanjoy Basu, E: sanjoy.basu@nhs.net

doi 10.1308/rcsann.2016.0206

The surgical management of medium-to-large (transverse defect >5cm) lateral incisional abdominal wall herniae can be technically challenging due to the absence of sheath and lack of ‘give’ for a tension-free closure. The results can be disappointing, especially when the defect lies in close proximity to bony landmarks, such as the iliac crest or costal margin.^{1,2} Here we describe our approach to these challenging cases.

Figure 1 The introduction of a composite mesh over the defect with a 4/5cm overlap and a 4/5cm rolled-over edge on each side