Published in final edited form as:

ACS Catal. 2017 January 6; 7(1): 150–153. doi:10.1021/acscatal.6b03217.

Rh(III)-Catalyzed Aryl and Alkenyl C–H Bond Addition to Diverse Nitroalkenes

Tyler J. Potter, David N. Kamber, Brandon Q. Mercado, and Jonathan A. Ellman

Department of Chemistry, Yale University, New Haven, Connecticut 06520, United States

Abstract

The transition metal catalyzed C–H bond addition to nitroalkenes has been developed. Very broad nitroalkene scope was observed for this Rh(III)-catalyzed method, including for aliphatic, aromatic and β,β -disubstituted derivatives. Additionally, various directing groups and both aromatic and alkenyl C–H bonds were effective in this transformation. Representative nitroalkane products were converted to dihydroisoquinolones and dihydropyridones in a single step and in high yield by iron mediated reduction and *in situ* cyclization. Moreover, preliminary success in enantioselective Rh(III)-catalyzed C–H bond addition to nitroalkenes was achieved as was X-ray structural characterization of a nitronate intermediate.

Abstract

Keywords

C-H functionalization; Rh(III) catalysis; nitroalkene; enantioselective catalysis; dihydroisoquinolone

Nitroalkenes are valued and extensively employed electrophiles for the addition of a broad range of different nucleophiles to give nitroalkanes, which are versatile intermediates to amines, carbonyl compounds and other useful chemotypes (Scheme 1A).¹ Many heteroatom and carbon nucleophiles have been added to nitroalkenes, including additions catalyzed by Lewis acids and organocatalysts. Nitroalkenes have also been used as Michael acceptors in

Correspondence to: Jonathan A. Ellman.

Supporting Information: Procedures, spectral data, and crystal data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Notes: The authors declare no competing financial interests.

Rh- and Pd-catalyzed additions of stable organometallic reagents such as aryl boronic acids.² However, despite dramatic recent advances in C-H bond functionalization, to our knowledge,^{3,4,5} transition metal catalyzed C-H bond additions to nitroalkenes have not been reported.

Herein, we report on the Rh(III)-catalyzed addition of aryl and alkenyl C(sp²)-H bonds to nitroalkenes (Scheme 1B). A broad range of C-H functionalization substrates with different directing groups as well as branched and unbranched aliphatic and aromatic nitroalkenes are effective inputs. Moreover, β,β -disubstituted nitroalkenes also coupled efficiently and represent the first examples of Rh(III)-catalyzed intermolecular C-H bond additions to trisubstituted alkenes.⁶ The utility of representative products was demonstrated by their conversion in a single step and in high yield to dihydroisoquinolones and dihydropyridones that cannot be accessed by alternative Rh(III)-catalyzed methods. Finally, promising initial results for enantioselective catalytic C-H bond addition to nitroalkenes was accomplished.

After evaluating a number of reaction parameters for coupling pyrrolidine benzamide **1a** and nitroalkene **2a**, optimal conditions were identified to be [Cp*RhCl₂]₂ as the precatalyst, AgSbF₆ as the chloride abstractor, dichloroethane as the solvent and an 80 °C reaction temperature (entry 1, Table 1). Importantly, product **3a** was obtained in high yield without the formation of any bis-addition byproduct.⁷ Both [Cp*RhCl₂]₂ and AgSbF₆ are essential as is apparent by the complete absence of product obtained when either is excluded (entries 2 and 3). Replacing AgSbF₆ with the less coordinating AgB(C₆F₅)₄ did not improve the yield of the reaction (entry 4).⁸ The addition of 20 mol % of AgOAc, an additive that has been reported to be successful for other Rh(III)-catalyzed transformations,⁹ decreased the yield to 65% for this reaction (entry 5). However, this additive was found to be beneficial for other C-H functionalization substrates (*vide infra*). The reaction proceeded equally well in 1,4-dioxane as solvent (entry 6), but a significantly lower yield was observed in AcOH despite its previous use as solvent in other Rh(III)-catalyzed additions to Michael acceptors.^{10,11} When the reaction was performed at room temperature, a slightly diminished yield was observed (entry 8). Because 80 °C proved to be the optimal temperature for more challenging C-H functionalization substrate/nitroalkene pairs, this reaction temperature was chosen for evaluating substrate scope. Use of the bench stable preformed catalyst Cp*Rh(CH₃CN)₃(SbF₆)₂ resulted in a comparable yield of 80% (entry 9), and when the reaction was performed on the bench top with the preformed catalyst, a 78% yield was observed (entry 10). While not as effective as [Cp*RhCl₂]₂, [Cp*IrCl₂]₂ and [RuCl₂(p-cymene)]₂ precatalysts did afford product **3a** in 53% and 62% yield, respectively (entries 11 and 12). No conversion was observed for the cobalt precatalyst [Cp*CoCl₂]₂ under these conditions (entry 13).

Very broad scope for the nitroalkene coupling partner was observed (Table 2). Product **3a** was isolated by silica gel chromatography in 85% yield using the standard conditions, and when the reaction was conducted on the bench top with the preformed catalyst Cp*Rh(CH₃CN)₃(SbF₆)₂, **3a** was isolated in 75% yield. This result highlights the tolerance of this reaction to air and ambient conditions. Additional branched and unbranched aliphatic nitroalkenes afforded products **3b**, **3c**, and **3d** in high yields. A broad range of aromatic β -nitrostyrenes were also effective coupling partners (**3e-3h**), including electron rich (**3f**),

electron poor (**3g**) and ortho-substituted (**3h**) derivatives. Nitroalkenes containing indole (**3i**) and furan (**3j**) heterocycles also coupled in acceptable yields.

We next explored β,β -disubstituted nitroalkenes. Although acyclic 2-methyl-1-nitroprop-1-ene and *trans*- α -methyl- β -nitrostyrene did not couple under these reaction conditions, we were pleased to find modest to good yields of the coupled products when employing azetidine (**3k** and **3l**) and cyclobutane (**3m**) nitroalkenes. Significantly, these products represent the first examples of a Rh(III)-catalyzed intermolecular C-H bond addition to trisubstituted alkenes.⁶

After establishing broad nitroalkene scope, we next explored the scope of the C-H functionalization substrate (Table 3). In addition to pyrrolidine benzamide, other amide directing groups such as *N,N*-diisopropylbenzamide (**3n**) and *N*-methylbenzamide (**3o**) were also successful in this reaction. Electron rich (**3p** and **q**) and electron poor (**3s**) substituents on the C-H bond functionalization substrate were well tolerated, and when the substituents were placed at the meta-position, high regioselectivity was observed for the least hindered site (**3q-s**). Heteroaromatic C-H bond activation substrates such as thiophene (**3t**), furan (**3u**), and indoles (**3v** and **3w**) afforded the addition products in good to excellent yields. In addition to aryl substrates, alkenyl amides were also effective C-H bond substrates with products **3x**, **3y**, and **3z** obtained in good yields. We also expanded the scope of the directing group beyond amides. The electron rich urea directing group afforded product **3aa** in a high yield. Additionally, *N*-heterocyclic directing groups such as pyrazole (**3ab**), triazole (**3ac**), and tetrazole (**3ad**) provided the addition products in good yields. Surprisingly, 2-phenylpyridine resulted in <5% of the desired addition product **3ae**, instead giving multiple unassigned decomposition products. The poor results obtained for the archetypal 2-pyridyl directing group, which is more basic than all of the other directing groups used, perhaps clarify why C-H bond additions to nitroalkenes had not previously been reported.

The weakly coordinating acetophenone C-H functionalization substrate was also evaluated, and required more forcing conditions similar to those reported for the Rh(III)-catalyzed synthesis of indenenes from acetophenones and enones.⁹ Under these conditions, addition and cyclization with concomitant loss of water afforded nitroindene **4** (eq 1).

(1)

To demonstrate the utility of the nitroalkene addition products **3**, representative derivatives were subjected to a single step iron-mediated nitro group reduction/cyclization sequence (Table 4). Dihydroisoquinolones have previously been synthesized by Rh(III)-catalyzed annulations of *O*-pivaloyl or *O*-Boc benzhydroxamic acids and alkenes, but controlling

selectivity is often a challenge, and a mixture of isomers is typically observed.¹² Dihydroisoquinolone **5a** was obtained in high yield with the structure confirmed by X-ray crystallographic analysis. This is a different isomer than that obtained by the Rh(III)-catalyzed annulation of *O*-Boc benzhydroxamic acid and styrene.¹³ Moreover, both dihydroisoquinolone **5b** and dihydropyridone **5c** were also obtained in good yields as single isomers and represent structural types that have not previously been prepared by Rh(III)-catalyzed alkene annulation approaches and would likely be very difficult to access.

A plausible mechanism for the Rh(III)-catalyzed C–H bond addition to nitroalkenes is shown in Scheme 2. Concerted metalation/deprotonation of **2** generates the well-documented rhodacycle **6**.¹⁴ Coordination of the nitroalkene **1** provides complex **7**, which upon insertion to the nitroalkene provides the Rh(III) nitronate **8**. Coordination of **2** followed by concerted metallation/deprotonation then concomitantly releases product **3** and regenerates rhodacycle **6**.

To investigate the mechanism of this transformation, we sought to characterize the proposed Rh(III) nitronate intermediate by X-ray crystallography (Scheme 3). Subjecting the preformed rhodacycle **9** to stoichiometric AgSbF₆ followed by excess β-nitrostyrene gave the Rh(III) nitronate **10** in 72% yield after trituration. The structure of **10** was rigorously characterized by X-ray crystallographic analysis as a C-bound nitronate dimer and to our knowledge represents the first X-ray structure of a rhodium nitronate.¹⁵ While **10** exists as the η¹-C-bound rhodacycle nitronate dimer in crystal form, it is possible that an η³-bound monomer exists in solution.¹⁶

Finally, in preliminary results using the Rh-diiodo dimer **11** employing Cramer's elegantly designed chiral ligand,¹⁷ we observed promising selectivities in the catalytic enantioselective synthesis of various alkyl (**3a**) and aromatic (**3e**, **3f**, **3g**) nitroalkene addition products (eq 2). Notably, electron-poor, neutral and rich aromatic nitroalkenes all coupled with 90:10 er. These results suggests a new general strategy for catalytic asymmetric additions to nitroalkenes.^{1b}

(2)

In conclusion, we have developed the first transition metal catalyzed C–H bond addition to nitroalkenes. Broad substrate scope was realized for both the C–H bond and nitroalkene coupling partners. Representative addition products were converted to dihydroisoquinolone and dihydropyridone heterocycles in a single step and in high yield by iron-mediated

reduction/cyclization. Additionally, preliminary success with the catalytic enantioselective addition of C-H bonds to nitroalkenes was achieved as was X-ray structural characterization of a Rh(III) nitronate intermediate.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by the NIH (GM069559). We gratefully acknowledge Dr. James Phelan (Yale) for important discussions contributing to the inception of this project, Dr. Michael Witten (Yale) for assistance with the acquisition of high-resolution mass spectrometric data, and Vlad Bacauanu (Yale) for providing an initial batch of chiral catalysts.

References

1. For select reviews on nitroalkenes and their use as Michael acceptors, see: Barrett AGM, Graboski GG. *Chem Rev.* 1986; 86:751–762. Berner OM, Tedeschi L, Enders D. *Eur J Org Chem.* 2002:1877–1894. Doyle AG, Jacobsen EN. *Chem Rev.* 2007; 107:5713–5743. [PubMed: 18072808]
2. For select examples of Rh- and Pd-catalyzed additions of aryl boronic acids to nitroalkenes, see: Hayashi T, Senda T, Ogasawara M. *J Am Chem Soc.* 2000; 122:10716–10717. Lu X, Lin S. *J Org Chem.* 2005; 70:9651–9653. [PubMed: 16268654] Wang ZQ, Feng CG, Zhang SS, Xu MH, Lin GQ. *Angew Chem Int Ed.* 2010; 49:5780–5783. Xue F, Wang D, Li X, Wan B. *J Org Chem.* 2012; 77:3071–3081. [PubMed: 22443387] He Q, Xie F, Fu G, Quan M, Shen C, Yang G, Gridnev ID, Zhang W. *Org Lett.* 2015; 17:2250–2253. [PubMed: 25879921]
3. For recent reviews on C-H functionalization, see: Yamaguchi J, Yamaguchi AD, Itami K. *Angew Chem Int Ed.* 2012; 51:8960–9009. Chen Z, Wang B, Zhang J, Yu W, Liu Z, Zhang Y. *Org Chem Front.* 2015; 2:1107–1295. Yang L, Huang H. *Chem Rev.* 2015; 115:3468–3517. [PubMed: 25749375] Gensch T, Hopkinson MN, Glorius F, Wencel-Delord J. *Chem Soc Rev.* 2016; 45:2900–2936. [PubMed: 27072661]
4. For reviews on Rh(III)-catalyzed C–H functionalization, see: Satoh T, Miura M. *Chem Eur J.* 2010; 16:11212–11222. [PubMed: 20740508] Patureau FW, Wencel-Delord J, Glorius F. *Aldrichimica Acta.* 2012; 45:31–41. Song G, Wang F, Li X. *Chem Soc Rev.* 2012; 41:3651–3678. [PubMed: 22377832]
5. For the Pd-catalyzed synthesis of C₁-benzoyl isoquinolines from isoquinoline *N*-oxides and β -nitrostyrenes via oxidative alkene cleavage, see: Li JL, Li WZ, Wang YC, Ren Q, Wang HS, Pan YM. *Chem Commun.* 2016; 52:10028–10031.
6. For examples of intramolecular Rh(III)-catalyzed C-H bond additions to trisubstituted alkenes, see: Davis TA, Hyster TK, Rovis T. *Angew Chem Int Ed.* 2013; 52:14181–14185. Davis TA, Wang C, Rovis T. *Synlett.* 2015; 26:1520–1524. [PubMed: 27540274] . For examples of intramolecular Rh(III)-catalyzed C-H bond additions to disubstituted alkenes, see: Shi Z, Bouladakis-Arapinis M, Koester DC, Glorius F. *Chem Commun.* 2014; 50:2650–2652. Chabaud L, Raynal Q, Barre E, Guillou C. *Adv Synth Catal.* 2015; 357:3880–3884.
7. For early examples of Rh-catalyzed transformations employing pyrrolidine amides as directing groups, see: Shibata Y, Otake Y, Hirano M, Tanaka K. *Org Lett.* 2009; 11:689–692. [PubMed: 19132939] Park J, Park E, Kim A, Lee Y, Chi KW, Kwak JH, Jung YH, Kim IS. *Org Lett.* 2011; 13:4390–4393. [PubMed: 21793534] Hesp KD, Bergman RG, Ellman JA. *Org Lett.* 2012; 14:2304–2307. [PubMed: 22500854] Wang F, Qi Z, Sun J, Zhang X, Li X. *Org Lett.* 2013; 15:6290–6293. [PubMed: 24274184]
8. For select examples where AgB(C₆F₅)₄ was more effective than AgSbF₆ in Rh(III)-catalyzed reactions, see: (a) ref 7c Otley KD, Ellman JA. *Org Lett.* 2015; 17:1332–1335. [PubMed: 25715077]
9. Shi XY, Li CJ. *Org Lett.* 2013; 15:1476–1479. [PubMed: 23470191]
10. Yang L, Qian B, Huang H. *Chem Eur J.* 2012; 18:9511–9515. [PubMed: 22744877]

11. Boerth JA, Ellman JA. Chem Sci. 2016; 7:1474–1479. [PubMed: 26918112]
12. (a) Guimond N, Gorelsky SI, Fagnou K. J Am Chem Soc. 2011; 133:6449–6457. [PubMed: 21452842] (b) Rakshit S, Grohmann C, Besset T, Glorius F. J Am Chem Soc. 2011; 133:2350–2353. [PubMed: 21275421] (c) Wodrich MD, Ye B, Gonthier JF, Corminboeuf C, Cramer N. Chem Eur J. 2014; 20:15409–15418. [PubMed: 25283798] (d) Hyster TK, Dalton DM, Rovis T. Chem Sci. 2015; 6:254–258. [PubMed: 25489470]
13. Although compound **5a** was reported to be the product of Rh(III)-catalyzed annulation using styrene,^{12c} our spectral data for **5a**, whose structure was rigorously confirmed by X-ray crystallographic analysis, is very different than the reported data. We conclude that the previously reported compound was incorrectly assigned.
14. For select relevant mechanistic studies for Rh(III)-catalysis, see: Li L, Brennessel WW, Jones WD. J Am Chem Soc. 2008; 130:12414–12419. [PubMed: 18714995] Brasse M, Campora J, Ellman JA, Bergman RG. J Am Chem Soc. 2013; 135:6427–6430. [PubMed: 23590843] Walsh AP, Jones WD. Organometallics. 2015; 34:3400–3407.
15. For examples of Co(III) and Ir(III) nitronates, see: Murata K, Konishi H, Ito M, Ikariya T. C-bound Ir. Organometallics. 2002; 21:253–255. Beckmann U, Ewing JD, Brooker S. C-bound Co. Chem Commun. 2003:1690–1691. Zhang X, Emge TJ, Ghosh R, Krogh-Jespersen K, Goldman AS. O- and C-bound Ir. Organometallics. 2006; 25:1303–1309.
16. A related η^3 -bound Cp*Rh(III) enolate was recently characterized by X-ray crystallography, see: ref 11.
17. For an outstanding recent Perspective on the application of Cramer's chiral ligands, see: Newton CG, Kossler D, Cramer N. J Am Chem Soc. 2016; 138:3935–3941. [PubMed: 26863546]

A. Prior work: Addition of nucleophiles to nitroalkenes **B. This work: First catalytic C-H bond addition to nitroalkenes** **Scheme 1. Nitroalkenes as Versatile Electrophiles**

Scheme 2. Proposed Mechanism for the Transformation

Scheme 3. Preparation of a Rh(III) Nitronate

Table 1

Control Reactions

Entry ^a	Variation from the standard conditions	Yield 3a (%) ^b
1	none	84
2	no [Cp [*] RhCl ₂] ₂	0
3	no AgSbF ₆	0
4	AgB(C ₆ F ₅) ₄ instead of AgSbF ₆	71
5	with 20 mol % AgOAc	65
6	in 1,4-dioxane	82
7	in AcOH	14
8	r.t. instead of 80 °C	66
9	Cp [*] Rh(CH ₃ CN) ₃ (SbF ₆) ₂ instead of [Cp [*] RhCl ₂] ₂	80
10 ^c	Cp [*] Rh(CH ₃ CN) ₃ (SbF ₆) ₂ instead of [Cp [*] RhCl ₂] ₂	78
11	[Cp [*] IrCl ₂] ₂ instead of [Cp [*] RhCl ₂] ₂	53
12	[RuCl ₂ (p-cymene)] ₂ instead of [Cp [*] RhCl ₂] ₂	62
13	[Cp [*] CoCl ₂] ₂ instead of [Cp [*] RhCl ₂] ₂	0

^aStandard conditions: 0.15 mmol of **1a**, 0.10 mmol of **2a** with [Cp^{*}RhCl₂]₂, AgSbF₆ in dichloroethane at 80 °C for 20 h.

^bYields determined by ¹H NMR spectroscopic analysis relative to SiMe₃Ph as an external standard.

^cReaction conducted on the bench top.

Table 2

Nitroalkene Scope^{a,b}

^a Conditions: 0.30 mmol of **1a**, 0.20 mmol of **2a** with $[\text{Cp}^*\text{RhCl}_2]_2$, AgSbF_6 in dichloroethane at 80 °C for 20 h.

^b Isolated yields after silica gel chromatography.

^c Reaction conducted on the bench top using 10 mol % of $\text{Cp}^*\text{Rh}(\text{CH}_3\text{CN})_3(\text{SbF}_6)_2$.

Table 3
C–H Functionalization Substrate Scope^{a,b}

^a Conditions: 0.30 mmol of **1a**, 0.20 mmol of **2a** with [Cp*RhCl₂]₂, AgSbF₆ in dichloroethane at 80 °C for 20 h.

^b Isolated yields after silica gel chromatography.

^c 20 mol % of AgOAc was added.

Table 4
Product Reduction and Cyclization^{a,b}

^a Conditions: **3** (1 equiv), Fe (6 equiv) in 1:1 EtOH/AcOH (0.2 M) at 80 °C for 3-6 h.

^b Isolated yields after aqueous workup.