CORRESPONDENCE

Fine Needle Aspiration in the Investigation of Thyroid Nodules—Indications, Procedures, and Interpretation

by PD Dr. med. Joachim Feldkamp, Prof. Dr. med. Dagmar Führer, Prof. Dr. med. Markus Luster, Prof. Dr. med. Thomas J. Musholt, Prof. Dr. med. Christine Spitzweg, and Prof. Dr. med. Matthias Schott in issue 20/2016

Increasing Importance

Feldkamp et al. emphasized the importance of the cytological investigation of thyroid nodules for their potential surgical treatment and define indications for and the validity of fine needle aspiration biopsy with a degree of clarity that is much welcomed, and while taking into consideration ultrasonographic and scintigraphic criteria (1). It is to be hoped that their article will find a wide readership among outpatient primary care providers too. I wish to comment on two issues.

- The statement that methoxyisobutylisonitrile (MIBI) scintigraphy is not useful in the routine work-up of thyroid nodules is too much of a generalization. At least this statement should be qualified by adding that this investigation can make a great contribution to reaching a decision in favor of or against surgery in cases where the cytology finding is non-diagnostic. According to recent relevant publications (2–4) it would not be surprising if in future MIBI scintigraphy increased in importance, since the accuracy of cytopathology within the outpatient setting is moderate at best.
- Because of the close correlation between tumor volume and calcitonin concentrations on the one hand and the now low-threshold access to molecular genetic diagnostic evaluation on the other hand, the benefit of fine needle aspiration cytology for the primary diagnostic evaluation of medullary thyroid cancer (MTC) has become mostly undetectable. For example, cytology in a scenario of a known germline mutation in the RET gene does not contribute to the decision on the extent or timing of thyroid surgery. The same is true for typical cases of sporadic MTCs, which become noticeable because of massively increased serum concentrations of calcitonin when investigating solitary or dominant hypoechoic nodules >1 cm in diameter. In this setting too, the surgeon who is familiar with the tumor entity will not make his or her strategy dependent on the results of fine needle aspiration cytology (5).

DOI: 10.3238/arztebl.2017.0025a

REFERENCES

1. Feldkamp J, Führer D, Luster M, Musholt TJ, Spitzweg C, Schott M: Fine needle aspiration in the investigation of thyroid nodules—indications, procedures and interpretation. *Dtsch Arztebl Int* 2016; 113: 353–9.
2. Wale A, Miles KA, Young B, et al.: Combined (99m)Tc-methoxyisobutylisonitrile scintigraphy and fine-needle aspiration cytology offers an accurate and potentially cost-effective investigative strategy for the assessment of solitary or dominant thyroid nodules. *Eur J Nucl Med Mol Imaging* 2014; 41: 105–15.
3. Riaz A, Kalantarhormozi M, Nabipour I, et al.: Technetium-99m methoxyisobutylisonitrile scintigraphy in the assessment of cold thyroid nodules: is it time to change the approach to the management of cold thyroid nodules? *Nucl Med Commun* 2014; 35: 51–7.
4. Giovanella L, Suriano S, Maffioli M, Ceriani L, Spriano G: (99m)Tc-sestamibi scanning in thyroid nodules with nondiagnostic cytology. *Head Neck* 2010; 32: 607–11.
5. Machens A, Dralle H: Biomarker-based risk stratification for previously untreated medullary thyroid cancer. *J Clin Endocrinol Metab* 2010; 95: 2655–63.

Dr. med. Tim Scholz
Brandenburg an der Havel
t.scholz@klinikum-brandenburg.de

Critical Assessment Is Missing

We wish to add the following to the review article on fine needle aspiration (FNA) biopsy (1):

- 1 In the second line of the *Figure* the large group of thyroid nodules >1 cm with normal findings is missing. A reference to published comprehensive studies and guidelines would have been useful.
- 2 No literature is cited that critically considers FNA biopsy. Tee et al. published an analysis of 12 studies including more than 54 000 patients and concluded that FNA biopsy potentially misses one-third of malignancies (2). McCartney et al. conducted a model analysis to compare four strategies for investigating thyroid nodules. Routine FNA biopsy came out worst (3). The effect of the low prevalence of malignant thyroid tumors in Germany on the results of FNA biopsy was not shown adequately. We fail to understand why the perspective of the American Thyroid Association (ATA) is being uncritically applied to the German thyroid scenario. The ATA guideline has thus far not been accepted by European thyroid specialists.
- 3 It is true that guidelines so far have not included tumor scintigraphy using Tc-99m-MIBI (methoxyisobutylisonitrile) for the purpose of further investigation of hypofunctioning thyroid nodules, so called “cold” thyroid nodules. However, it would have been appropriate to point out the high negative predictive value of >97% of a normal MIBI finding to rule out malignancy, and

the positive predictive value of about 15–20% for a malignant thyroid tumor (18 publications, >1000 patients with a histopathological gold standard [4]).

DOI: 10.3238/arztebl.2017.0025b

REFERENCES

1. Feldkamp J, Führer D, Luster M, Musholt TJ, Spitzweg C, Schott M: Fine needle aspiration in the investigation of thyroid nodules—indications, procedures and interpretation. *Dtsch Arztebl Int* 2016; 113: 353–9.
2. Tee YY, Lowe AJ, Brand CA, Judson RT: Fine-needle aspiration may miss a third of all malignancy in palpable thyroid nodules: a comprehensive literature review. *Ann Surg* 2007; 246: 714–20.
3. McCartney CR, Stukenborg GJ: Decision analysis of discordant thyroid nodule biopsy guideline criteria. *J Clin Endocrinol Metab* 2008; 93: 3037–44.
4. Schmidt M: Tc-99m-MIBI for thyroid imaging. In: Bucerius J, Ahmadzadehfar H, Biersack HJ: ^{99m}Tc-sestamibi—clinical applications, Springer-Verlag 2012: 133–58.

Prof. Dr. med. Matthias Schmidt
Universitätsklinikum Köln
matthias.schmidt@uk-koeln.de

Prof. Dr. med. Dr. Dipl.-Phys. Wolfgang Schäfer
Kliniken Maria Hilf GmbH, Mönchengladbach

Unanswered Questions

I am a general practitioner who does thyroid ultrasonography himself. My experience with two large nuclear medical practices has been that in nodules of unclear dignity, MIBI scintigraphy is undertaken almost as a matter of routine. If the results of such an investigation do not indicate a malignant nodule, then no further diagnostic evaluation is required. If malignancy is still suspected, then the advice is to have surgery.

Fine needle aspiration biopsy instead of MIBI scintigraphy seems to me to be the exception rather than the rule. In my experience, there are few centers where patients might be referred to for FNA biopsy.

If one were to follow the assessment provided by Feldkamp et al., the diagnostic pathway for the investigation of thyroid nodules would have to be changed (1). However, it should not be forgotten that patients view a second scintigram as much more innocuous than fine needle aspiration. I would therefore invite the authors to comment again on this issue.

DOI: 10.3238/arztebl.2017.0026a

REFERENCES

1. Feldkamp J, Führer D, Luster M, Musholt TJ, Spitzweg C, Schott M: Fine needle aspiration in the investigation of thyroid nodules—indications, procedures and interpretation. *Dtsch Arztebl Int* 2016; 113: 353–9.

Dr. med. Eberhard Backus
Frechen
info@hausarzt-backus-frechen.de

In Reply:

Schäfer and Schmidt in their response discuss thyroid nodules with a diameter of >1 cm and normal findings. In order to keep the *Figure* showing our diagnostic algorithm as simple as possible we did not discuss this aspect. Similarly, because of restrictions on word count and cited literature, we did not include any further studies on the validity of fine needle aspiration (FNA). We agree with our correspondents that the current recommendations of the American Thyroid Association (ATA) cannot unconditionally be generalized to the European setting. Independently of these issues, however, we wish to reiterate that FNA biopsy of the thyroid if undertaken by an experienced examiner and an experienced cytologist has a high degree of accuracy in the diagnostic evaluation of thyroid cancer and can also help prevent surgical procedures without a clear indication.

All three letters mention MIBI scintigraphy as a method for the differential diagnostic evaluation of thyroid nodules. Our review article aimed to focus on FNA (1). To date, the number of studies on MIBI scintigraphy that have been conducted is too small, and the procedure has not been approved for the diagnostic evaluation of thyroid nodules. For these reasons, MIBI scintigraphy has not been included in current guidelines. Our colleagues point out that in the studies published to date, a negative MIBI result is highly likely to indicate a benign thyroid nodule. The future role of MIBI scintigraphy as a complementary approach will need to be investigated in prospective studies.

Scholz explains the primary diagnostic evaluation of medullary thyroid nodules (MTC). We agree with him that cytology is of secondary importance in this setting. Calcitonin measurement is the most sensitive method and should be done at the very least before surgery for a thyroid nodule/(multi)nodular goiter.

DOI: 10.3238/arztebl.2017.0026b

REFERENCES

1. Feldkamp J, Führer D, Luster M, Musholt TJ, Spitzweg C, Schott M: Fine needle aspiration in the investigation of thyroid nodules—indications, procedures and interpretation. *Dtsch Arztebl Int* 2016; 113: 353–9.

On behalf of the authors:
Prof. Dr. Matthias Schott
Universitätsklinikum Düsseldorf
matthias.schott@med.uni-duesseldorf.de

Conflict of interest statement

The authors of all contributions declare that no conflict of interest exists.