Published in final edited form as:

Cytokine Growth Factor Rev. 2017 April ; 34: 67–72. doi:10.1016/j.cytogfr.2017.01.003.

Crosstalk between insulin-like growth factor (IGF) receptor and integrins through direct integrin binding to IGF1

Yoshikazu Takada^{1,2}, Yoko K. Takada^{1,2}, and Masaaki Fujita³

¹Departments of Dermatology, Biochemistry and Molecular Medicine, School of Medicine, University of California, Davis, Sacramento, California 95817

²Graduate Institute of Translational Medicine, College of Medical Science and Technology, Taipei Medical University, 250 Wu-Hsing Street, Taipei 11031, Taiwan, R.O.C

³Department of Clinical Immunology and Rheumatology, The Tazuke-Kofukai Medical Research Institute, Kitano Hospital, Osaka, Japan

Abstract

It has been generally accepted that integrin cell adhesion receptors are involved in growth factor signaling (integrin-growth factor crosstalk), since antagonists to integrins often suppress growth factor signaling. Partly because integrins have been originally identified as cell adhesion receptors to extracellular matrix (ECM) proteins, current models of the crosstalk between IGF1 and integrins propose that ECM ligands (e.g., vitronectin) bind to integrins and IGF1 binds to IGF receptor type 1 (IGF1R), and two separate signals merge inside the cells. Our research proves otherwise. We discovered that IGF1 interacts directly with integrins, and induces integrin-IGF-IGF1R complex formation on the cell surface. IGF1 signaling can be detected in the absence of ECM (anchorage-independent conditions). Integrin antagonists block both ECM-integrin interaction and IGF-integrin interaction, and do not distinguish the two. This is one possible reason why integrin-IGF1 interaction has not been detected. With these new discoveries, we believe that the direct IGF-integrin interaction should be incorporated into models of IGF1 signaling. The integrin-binding defective mutant of IGF1 is defective in inducing IGF signaling, although the mutant still binds to IGF1R. Notably, the IGF1 mutant is dominant-negative and suppresses cell proliferation induced by wt IGF1, and suppresses tumorigenesis in vivo, and thus the IGF1 mutant has potential as a therapeutic.

Keywords

IGF1; integrin; crosstalk; dominant-negative antagonist; signaling

Address correspondence to: Yoshikazu Takada, MD, PhD, Department of Dermatology, and Biochemistry and Molecular Medicine, University of California Davis School of Medicine, Research III, Suite 3300, 4645 Second Avenue, Sacramento, CA 95817, Tel 916-734-7443, Fax 916-734-7505, ytakada@ucdavis.edu and Graduate Institute of Translational Medicine, College of Medical Science and Technology, Taipei Medical University, 250 Wu-Hsing Street, Taipei 11031, ytakada@tmu.edu.tw.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

IGF-1/IGF1R axis is a therapeutic target of cancer

Insulin-like growth factor-1 (IGF1) is a polypeptide hormone that is homologous to proinsulin. Most of (80%) of the IGF1 in serum is synthesized by the liver and secreted, and it functions as an endocrine hormone. The remaining 20% of the IGF1 is synthesized in the periphery. Usually, connective tissue cell types, such as stromal cells, and IGF1 that is synthesized in the periphery can function to regulate cell survival by autocrine and paracrine mechanisms [1]. IGF1 is also secreted by many cancer cells at abnormally high levels. Once released by cancer cells, IGF1 binds and activates the type 1 IGF receptor (IGF1R) on their surface. IGF1R is ubiquitously present on multiple cell types. Ligand binding induces phosphorylation of specific tyrosine residues of IGF1R. These phosphotyrosines then bind to adapter molecules such as Shc and insulin receptor substrate-1. Phosphorylation of these proteins leads to activation of the phosphoinositide 3-kinase and mitogen activated protein kinase signaling pathways [2]. IGF1, thereby, confers cancer cells resistance to chemotherapy and radiation therapy. Thus, IGF1 is a major therapeutic target for cancer. Several strategies to target IGF1 signaling have been developed, including, siRNA and monoclonal antibodies for IGF-IR, and kinase inhibitors to inhibit the enzymatic activity of the receptor. The elucidation of the IGF1 signaling pathway (e.g., role of integrins, see below) should have a major impact in designing new therapeutic strategies [1].

IGF1R is a marker of cancer stem cells

Several reports suggest that IGF1R is a marker of cancer stem cells (CSCs). IGF1R signaling is critical for maintaining “stemness” of CSCs, which is detected by the expression of Oct-4 and Nanog transcription factors. IGF1R is activated by IGF2, which is secreted by cancer-associated fibroblasts in a paracrine manner. IGF2 enhances expression of Oct-4 and Nanog in lung CSCs through IGF1R signaling and Akt phosphorylation (paracrine model) [3]. Development of new strategies to suppress IGF signaling should have a major impact in cancer treatment. IGF1 decoy is expected to inhibit IGF1R signaling in CSCs, and suppress stemness or the number of CSCs.

Previous studies on IGF1 signaling and integrin $\alpha v \beta 3$

Integrins are a family of cell adhesion receptors that recognize extracellular matrix ligands and cell surface ligands [4]. They are transmembrane α - β heterodimers, and at least 18 α and 8 β subunits are known [5]. They are involved in signal transduction upon ligand binding [4]. Ligation of integrins also triggers a large variety of signal transduction events that serve to modulate cell behavior, including proliferation, survival/apoptosis, shape, polarity, motility, gene expression, and differentiation.

It has been well established that integrin $\alpha v \beta 3$ plays a critical role in IGF signaling in addition to IGF1R [6–8]. “Ligand occupancy” of $\alpha v \beta 3$ (*i.e.* the binding of extracellular matrix (ECM) proteins such as vitronectin to $\alpha v \beta 3$) enhances signaling induced by IGF1 binding to IGF1R [1]. It has been proposed that ECM ligands bind to integrins and IGF1 binds to IGF1R, and two separate signals merge inside the cells. Blocking of vitronectin- $\alpha v \beta 3$ integrin interaction using echistatin, a snake venom disintegrin that specifically

inhibits $\alpha v\beta 3$, inhibits IGF1-induced IGF1R phosphorylation, DNA synthesis [9] and cell migration [7]. The disulfide-linked loop of integrin $\beta 3$ (The 177–184) in the ligand-binding site of this integrin is involved in vitronectin binding, and is necessary for IGF1 stimulated cell migration and proliferation [10]. The antibody against the disulfide-linked loop of $\beta 3$ inhibits IGF1 signaling (IGF1-stimulated Shc phosphorylation and ERK1/2 activation) [10]. Anti- $\alpha v\beta 3$ mAb and echistatin block IGF1-induced cell migration [11]. Also, echistatin blocked IGF1-stimulated DNA synthesis and IRS-1 phosphorylation, and attenuated IGF1R-linked down stream signaling events, such as activation of PI-3K and MAP kinase ERK1/2 [9]. We recently discovered that IGF1 interacts directly with integrins. Integrin antagonists block both ECM-integrin interaction and IGF-integrin interaction, and do not distinguish the two. This is one possible reason why integrin-IGF1 interaction has not been detected. *With these new discoveries*, we believe that the IGF-integrin interaction should be incorporated into models of IGF1 signaling.

Direct binding of IGF1 to integrin $\alpha v\beta 3$ is critical for IGF signaling

The first indication that IGF1 binds to integrins is docking simulation, which we have been using to identify potential new integrin ligands. Docking simulation predicted that IGF1 binds to integrin $\alpha v\beta 3$ well. We confirmed that this is really the case using human IGF1 we generated in our laboratory, and commercial IGF1 [12]. In surface plasmon resonance (SPR) analysis of IGF1- $\alpha v\beta 3$ interaction using immobilized recombinant soluble $\alpha v\beta 3$ to a sensor chip, we obtained $K_D 5 \times 10^{-7}$ M, which is a reasonable affinity for integrin-ligand interaction. The simulation predicted the potential integrin-binding sites in IGF1 [12]. Mutating the Arg residues at positions 36/37 in the predicted integrin-binding site of IGF1 into Glu (designated R36E/R37E mutation) markedly reduced integrin binding of IGF1 [12]. The R36E/R37E mutant is defective in inducing IGF signaling (e.g., IGF1R phosphorylation, ERK1/2 activation, and cell proliferation), although the mutant still binds to IGF1R [12]. This suggests that the direct binding of integrins to IGF1 is critical for IGF signaling in addition to binding to IGF1R.

IGF1 induces integrin-IGF1-IGF1R ternary complex

Furthermore, WT IGF1 induces integrin-IGF1-IGF1R ternary complex formation, while R36E/R37E does not (Fig. 1a), suggesting that the ternary complex formation plays a role in IGF1 signaling [12]. Our studies show phosphorylation of IGF1R is not required for IGF1-induced integrin-IGF1R association. Inhibitors of IGF1R (PPP) and Src (PP2) does not suppress $\alpha v\beta 3$ -IGF-IGF1R ternary complex formation [13], suggesting that phosphorylation of IGF1R is not required for ternary complex formation. We propose a model, in which IGF1 binding to IGF1R induces recruitment of integrin $\alpha v\beta 3$ to the IGF-IGF1R complex, and then $\beta 3$ and IGF1R are phosphorylated [13] (Fig. 1b). It is likely that $\alpha v\beta 3$ is together with the IGF1-IGF1R complex for triggering IGF signaling.

IGF1-induced physical association of integrin $\beta 3$ and IGF1R is detected by BiFC and FRET studies

The association of $\alpha v\beta 3$ and IGF1R has recently been studied in bimolecular fluorescence complementation (BiFC) studies using $\beta 3$ tagged with YFP1–158 and IGF1R tagged with YFP159–299 [14]. $\alpha v\beta 3$ and IGF1R did not interact on the cell surface, but WT IGF1 potentiated the association. Essentially the same results were obtained in FRET studies (CFP-tagged integrin $\beta 3$ and YFP tagged IGF1R) [14]. These findings are consistent with our biochemical findings that IGF1 binding to $\alpha v\beta 3$ induces ternary complex formation with IGF1R.

IGF1 signaling, integrin $\alpha 6\beta 4$, and cancer

The $\alpha 6\beta 4$ integrin is a receptor for the laminin family of extracellular matrix proteins, and its expression is associated with poor patient prognosis and reduced survival in a variety of human cancers [15]. $\alpha 6\beta 4$ combines with multiple oncogenic receptor tyrosine kinases, such as ErbB2 [16], EGFR [17], and Met [18] and enhances the signaling function. However, how $\alpha 6\beta 4$ and RTKs interact is not well understood. $\alpha 6\beta 4$ in normal epithelial cells is located in basal surface of the cells, and $\alpha 6\beta 4$ is relocated to the apical region of the cells when cells were transformed or stimulated with EGF [19]. The change in $\alpha 6\beta 4$ localization is accompanied by the loss of polarity and shape of the epithelial cells. It is expected that $\alpha 6\beta 4$ is exposed to growth factors when it moves from the basal surface of the cells, in which $\alpha 6\beta 4$ is shielded from growth factors, to the apical region.

The $\beta 4$ subunit was originally identified as a tumor-related antigen expressed in metastatic cancer [20]. In contrast with its function in regulating stable adhesion through the formation of hemidesmosomes in normal epithelial cells, $\alpha 6\beta 4$ promotes motility and invasion in carcinoma cells [21]. Moreover, suppression of $\alpha 6\beta 4$ expression by siRNA diminishes invasive potential [22]. $\alpha 6\beta 4$ expression has been correlated with the progression and metastatic potential of several different tumors including breast cancer and pancreatic cancer [23,24,25]. $\alpha 6\beta 4$ is up regulated in various types of cancer (squamous, breast, thyroid, bladder, colorectal, and gastric)[19]. In addition, $\alpha 6\beta 4$ is expressed in tumor blood vessels and promotes the invasion phase of tumor angiogenesis [26].

We demonstrated that integrin $\alpha 6\beta 4$ directly binds to IGF1. WT IGF1 induced ternary complex formation ($\alpha 6\beta 4$ -IGF1-IGF1R) in MCF-7 breast cancer cells, but R36E/R37E IGF1 did not (see below), suggesting that $\alpha 6\beta 4$ -IGF1 interaction plays a role in IGF1 signaling possibly through integrin-IGF1R crosstalk.

Overexpression of $\alpha 6\beta 4$ enhances, and down regulation of $\alpha 6\beta 4$ suppresses, proliferation of MCF-7 cells [27,15], suggesting that $\alpha 6\beta 4$ plays a role in proliferation of this cell type. We found that $\alpha 6\beta 4$ binds to IGF1 at a level comparable to that of $\alpha v\beta 3$. Interestingly, $\alpha 6\beta 4$ does not adhere to R36E/R37E, suggesting that $\alpha 6\beta 4$ binds to IGF1 in a manner similar to that of $\alpha v\beta 3$. Also, $\alpha 6\beta 4$ co-precipitated with wt IGF1 but not well with R36E/R37E, suggesting that $\alpha 6\beta 4$ specifically recognize IGF1. We found that WT IGF1 induced co-precipitation of IGF1R and integrin $\beta 4$, but R36E/R37E was defective in this function. This

suggests $\alpha 6\beta 4$ -IGF1-IGF1R ternary complex formation is involved in IGF signaling in cancer cells.

The R36E/R37E IGF1 mutant suppresses cell proliferation, anchorage-independent growth (in soft agar), and tumorigenesis

We stably expressed WT and mutant IGF1 in mouse and human breast cancer cells using secretion vector. WT IGF1 enhances proliferation of stably transfected cells, while R36E/R37E suppresses it. We tested if WT and mutant IGF1 affects anchorage-independent growth, an important hallmark of cell transformation, of cancer cells. Interestingly, WT IGF1 enhances the number of colonies in soft agar, while R36E/R37E slightly reduced it. We studied if IGF1 can induce signaling in anchorage-independent conditions in transformed cells (Chinese hamster ovary, CHO) that express $\alpha v\beta 3$ ($\beta 3$ -CHO) cells. We found that IGF1 signals were more clearly detectable in anchorage-independent conditions (polyHEMA-coated plates) than in regular tissue culture conditions. This suggests that IGF signaling is masked by signals from cell-matrix interaction in regular tissue culture. IGF signaling required $\alpha v\beta 3$ expression, and R36E/R37E was defective in inducing signals in polyHEMA-coated plates. These results suggest that $\alpha v\beta 3$ -IGF1 interaction, not $\alpha v\beta 3$ -extracellular matrix interaction, is essential for IGF signaling.

We tested whether R36E/R37E IGF1 affects tumorigenesis in vivo using cancer cells that constitutively secrete WT IGF1 or R36E/R37E. R36E/R37E suppresses tumorigenesis in vivo, while WT IGF1 enhances it. This is consistent with the results in vitro that R36E/R37E is dominant-negative.

We found that excess R36E/R37E suppresses the binding of labeled-WT IGF1 to the cell surface [28], and suppresses ternary complex formation induced by WT IGF1 [28]. This demonstrates that R36E/R37E competes with WT IGF1 for the binding to IGF1R on the cell surface.

IGF signaling occurs in anchorage-independent conditions

It has been proposed that integrin-ECM interaction is required for signaling from growth factors [29,30]. In IGF signaling, “ligand occupancy” of $\alpha v\beta 3$ through vitronectin binding to $\alpha v\beta 3$ enhances signaling induced by IGF1 binding to IGF1R [1]. We found, however, that exogenous WT IGF1 did not induce detectable signaling in regular tissue culture (2-dimensional culture) [13]. We hypothesized that massive proliferative signals from cell-ECM interaction masks the signals from exogenous IGF1. We thus suppressed cell-ECM interaction by coating the plastic surface with hydrogel (polyHEMA), which generate anchorage-independent conditions [31]. We also used soft agar plates, a well-established anchorage-independent condition. Under these conditions we detected the effect of exogenous WT IGF1, the dominant-negative effect of R36E/R37E, and the effect of $\alpha v\beta 3$ or $\alpha 6\beta 4$ expression [32]. (Fig. 2)

These findings are consistent with a previous report that cell-matrix adhesion masks the heparin-binding EGF signaling because cell-matrix adhesion provides cells sufficient

proliferative signals through cell-matrix adhesion, but that it is possible to detect the proliferative effect of heparin binding EGF on cancer cells in vitro in three- or two-dimensional culture in which cell-matrix interaction is reduced [33]. Thus it is unclear if cell-ECM interaction really enhances IGF1 signaling because it masks the effect of exogenous IGF1 signaling. It is likely that exogenous IGF may not be required for proliferation or survival when cells interact with ECM. If tumor growth in anchorage-independent conditions, in which cell-ECM interaction is reduced, mimics tumorigenesis in vivo, it is highly likely that IGF1-integrin interaction plays a major role in tumorigenesis rather than integrin-ECM interaction. Thus we will need to study how ECM and IGF (and perhaps other cytokines) contribute to proliferation and cell survival in future studies.

Can mutants of human proteins be used as therapeutic agents?

There is a precedent that a mutant of human protein was used for human diseases. A mutant of human growth hormone (hGH) has been used as an antagonist of GH receptor in the treatment of acromegaly (Pegmicovent) [34]. The Gly-120 of h GH was mutated to Arg (G120R) and this mutant was further modified by polyethyleneglycol (PEG)-5000 to elongate half-life. Pegvisomant prevents functional dimerization of hGH receptor by sterically inhibiting conformational changes within the GHR dimers [34]. Pegvisomant is generally well tolerated with a safety profile similar to that reported in clinical trials and can effectively reduce IGFI in patients with acromegaly refractory to conventional therapy [35].

Integrin-growth factor crosstalk through direct integrin binding to growth factor is not limited to IGF1

Our previous studies found that FGF1 directly bind to integrins using docking simulation [36]. Based on the docking model, we generate integrin-binding-defective FGF1 by introducing mutations in the predicted integrin-binding site. The integrin-binding-defective mutant (the R50E mutant) is not only defective in inducing signaling, it also suppresses FGF signaling induced by wild-type FGF1 (dominant-negative effect by definition) [37]. Wild-type FGF1 induces ternary complex formation (integrin-FGF-FGF receptor), but R50E does not. WT FGF1 induces ternary complex formation (integrin, FGF1, and FGF receptor-1 (FGFR1), when cells are treated with FGF1. The integrin-binding-defective mutant of FGF1 does not induce ternary complex formation, suggesting that the ability of FGF1 to interact with integrins is required for FGF signaling. Notably, we found that the integrin-binding defective mutant of FGF1 acts as an antagonist to FGF signaling (dominant-negative mutants) [37,38], suggesting that the FGF mutant has potential as a therapeutic. We described very similar findings in neuregulin-1 [39], and fractalkine [40]. Thus integrin-growth factor crosstalk through direct integrin binding to growth factor is not limited to IGF1.

Conclusion

Our studies suggest that integrin binding to IGF1 and subsequent ternary complex formation is required for IGF1 signaling (Fig. 1). The integrin-binding defective IGF1 mutant is an interesting tool for studying the mechanism of integrin-IGF1 crosstalk. The dominant-

negative IGF1 mutant has potential as a therapeutic in cancer. Thus we need to fully evaluate the potential of the IGF1 mutant as a therapeutic agent. How can we deliver the dominant-negative IGF1 mutant to diseased tissues? Small devices that deliver insulin to diabetic patients have been developed, in which timing and amount of delivery are fully programmable. One possibility is that we will be able to exploit these devices for delivering IGF1 instead of insulin.

Acknowledgments

Funding: This work was supported by The Kanoe Foundation for the Promotion of Medical Science (MF), The Tazuke-Kofukai Medical Research Institute (MF), and Mitsubishi Tanabe Pharma Corporation (MF).

Abbreviations

CHO	Chinese hamster ovary
CSC	cancer stem cells
ECM	extracellular matrix
IGF1R	insulin-like growth factor type I receptor
IGF1	insulin-like growth factor-1
polyHEMA	poly-2-hydroxyethyl methacrylate

References

1. Clemmons DR. Modifying IGF1 activity: an approach to treat endocrine disorders, atherosclerosis and cancer. *Nat Rev Drug Discov.* 2007; 6(10):821–33. [PubMed: 17906644]
2. Clemmons DR, Maile LA. Interaction between insulin-like growth factor-I receptor and alphaVbeta3 integrin linked signaling pathways: cellular responses to changes in multiple signaling inputs. *Mol Endocrinol.* 2005; 19(1):1–11. [PubMed: 15528274]
3. Chen WJ, Ho CC, Chang YL, Chen HY, Lin CA, Ling TY, Yu SL, Yuan SS, Chen YJ, Lin CY, Pan SH, Chou HY, Chen YJ, Chang GC, Chu WC, Lee YM, Lee JY, Lee PJ, Li KC, Chen HW, Yang PC. Cancer-associated fibroblasts regulate the plasticity of lung cancer stemness via paracrine signalling. *Nature communications.* 2014; 5:3472.
4. Hynes RO. Integrins: bidirectional, allosteric signaling machines. *Cell.* 2002; 110(6):673–87. [PubMed: 12297042]
5. Takada Y, Ye X, Simon S. The integrins. *Genome Biol.* 2007; 8(5):215. [PubMed: 17543136]
6. Soung YH, Clifford JL, Chung J. Crosstalk between integrin and receptor tyrosine kinase signaling in breast carcinoma progression. *BMB Rep.* 2010; 43(5):311–8. [PubMed: 20510013]
7. Jones JI, Prevette T, Gockerman A, Clemmons DR. Ligand occupancy of the alpha-V-beta3 integrin is necessary for smooth muscle cells to migrate in response to insulin-like growth factor. *Proc Natl Acad Sci U S A.* 1996; 93(6):2482–7. [PubMed: 8637900]
8. Zheng B, Duan C, Clemmons DR. The effect of extracellular matrix proteins on porcine smooth muscle cell insulin-like growth factor (IGF) binding protein-5 synthesis and responsiveness to IGF-I. *The Journal of biological chemistry.* 1998; 273(15):8994–9000. [PubMed: 9535886]
9. Zheng B, Clemmons DR. Blocking ligand occupancy of the alphaVbeta3 integrin inhibits insulin-like growth factor I signaling in vascular smooth muscle cells. *Proc Natl Acad Sci U S A.* 1998; 95(19):11217–22. [PubMed: 9736716]
10. Maile LA, Busby WH, Sitko K, Capps BE, Sergeant T, Badley-Clarke J, Clemmons DR. Insulin-like growth factor-I signaling in smooth muscle cells is regulated by ligand binding to the

- 177CYDMKTTTC184 sequence of the beta3-subunit of alphaVbeta3. *Mol Endocrinol.* 2006; 20(2): 405–13. [PubMed: 16195248]
11. Jones JI, Doerr ME, Clemmons DR. Cell migration: interactions among integrins, IGFs and IGFs. *Prog Growth Factor Res.* 1995; 6(2–4):319–27. [PubMed: 8817675]
 12. Saegusa J, Yamaji S, Ieguchi K, Wu CY, Lam KS, Liu FT, Takada YK, Takada Y. The direct binding of insulin-like growth factor-1 (IGF-1) to integrin alphavbeta3 is involved in IGF-1 signaling. *The Journal of biological chemistry.* 2009; 284(36):24106–14. [PubMed: 19578119]
 13. Fujita M, Takada YK, Takada Y. Insulin-like Growth Factor (IGF) Signaling Requires alphavbeta3-IGF1-IGF Type 1 Receptor (IGF1R) Ternary Complex Formation in Anchorage Independence, and the Complex Formation Does Not Require IGF1R and Src Activation. *The Journal of biological chemistry.* 2013; 288(5):3059–69. [PubMed: 23243309]
 14. Tahimic CG, Long RK, Kubota T, Sun MY, Elalieh H, Fong C, Menendez AT, Wang Y, Vilardaga JP, Bikle DD. Regulation of Ligand and Shear Stress-induced Insulin-like Growth Factor 1 (IGF1) Signaling by the Integrin Pathway. *The Journal of biological chemistry.* 2016; 291(15):8140–9. [PubMed: 26865633]
 15. Lu S, Simin K, Khan A, Mercurio AM. Analysis of integrin beta4 expression in human breast cancer: association with basal-like tumors and prognostic significance. *Clinical cancer research : an official journal of the American Association for Cancer Research.* 2008; 14(4):1050–8. [PubMed: 18281537]
 16. Guo W, Pylyayeva Y, Pepe A, Yoshioka T, Muller WJ, Inghirami G, Giancotti FG. Beta 4 integrin amplifies ErbB2 signaling to promote mammary tumorigenesis. *Cell.* 2006; 126(3):489–502. [PubMed: 16901783]
 17. Mariotti A, Kedeshian PA, Dans M, Curatola AM, Gagnoux-Palacios L, Giancotti FG. EGF-R signaling through Fyn kinase disrupts the function of integrin alpha6beta4 at hemidesmosomes: role in epithelial cell migration and carcinoma invasion. *The Journal of cell biology.* 2001; 155(3): 447–58. [PubMed: 11684709]
 18. Trusolino L, Bertotti A, Comoglio PM. A signaling adapter function for alpha6beta4 integrin in the control of HGF-dependent invasive growth. *Cell.* 2001; 107(5):643–54. [PubMed: 11733063]
 19. Mercurio AM, Rabinovitz I. Towards a mechanistic understanding of tumor invasion--lessons from the alpha6beta 4 integrin. *Semin Cancer Biol.* 2001; 11(2):129–41. [PubMed: 11322832]
 20. Falcioni R, Kennel SJ, Giacomini P, Zupi G, Sacchi A. Expression of tumor antigen correlated with metastatic potential of Lewis lung carcinoma and B16 melanoma clones in mice. *Cancer research.* 1986; 46(11):5772–8. [PubMed: 3756921]
 21. Borradori L, Sonnenberg A. Structure and function of hemidesmosomes: more than simple adhesion complexes. *J Invest Dermatol.* 1999; 112(4):411–8. [PubMed: 10201522]
 22. Lipscomb EA, Dugan AS, Rabinovitz I, Mercurio AM. Use of RNA interference to inhibit integrin (alpha6beta4)-mediated invasion and migration of breast carcinoma cells. *Clin Exp Metastasis.* 2003; 20(6):569–76. [PubMed: 14598892]
 23. Streuli CH, Akhtar N. Signal co-operation between integrins and other receptor systems. *The Biochemical journal.* 2009; 418(3):491–506. [PubMed: 19228122]
 24. Bon G, Folgiero V, Di Carlo S, Sacchi A, Falcioni R. Involvement of alpha6beta4 integrin in the mechanisms that regulate breast cancer progression. *Breast Cancer Res.* 2007; 9(1):203. [PubMed: 17319974]
 25. Wilhelmssen K, Litjens SH, Sonnenberg A. Multiple functions of the integrin alpha6beta4 in epidermal homeostasis and tumorigenesis. *Molecular and cellular biology.* 2006; 26(8):2877–86. [PubMed: 16581764]
 26. Nikolopoulos SN, Blaikie P, Yoshioka T, Guo W, Giancotti FG. Integrin beta4 signaling promotes tumor angiogenesis. *Cancer Cell.* 2004; 6(5):471–83. [PubMed: 15542431]
 27. Jones JL, Royall JE, Critchley DR, Walker RA. Modulation of myoepithelial-associated alpha6beta4 integrin in a breast cancer cell line alters invasive potential. *Exp Cell Res.* 1997; 235(2):325–33. [PubMed: 9299156]
 28. Fujita M, Ieguchi K, Cedano-Prieto DM, Fong A, Wilkerson C, Chen JQ, Wu M, Lo SH, Cheung AT, Wilson MD, Cardiff RD, Borowsky AD, Takada YK, Takada Y. An Integrin Binding-defective Mutant of Insulin-like Growth Factor-1 (R36E/R37E IGF1) Acts as a Dominant-negative

- Antagonist of the IGF1 Receptor (IGF1R) and Suppresses Tumorigenesis but Still Binds to IGF1R. *The Journal of biological chemistry*. 2013; 288(27):19593–603. [PubMed: 23696648]
29. Eliceiri BP. Integrin and growth factor receptor crosstalk. *Circ Res*. 2001; 89(12):1104–10. [PubMed: 11739274]
 30. Schwartz MA, Ginsberg MH. Networks and crosstalk: integrin signalling spreads. *Nat Cell Biol*. 2002; 4(4):E65–8. [PubMed: 11944032]
 31. Folkman J, Moscona A. Role of cell shape in growth control. *Nature*. 1978; 273(5661):345–9. [PubMed: 661946]
 32. Fujita M, Ieguchi K, Davari P, Yamaji S, Taniguchi Y, Sekiguchi K, Takada YK, Takada Y. Cross-talk between integrin $\alpha 6 \beta 4$ and insulin-like growth factor-1 receptor (IGF1R) through direct $\alpha 6 \beta 4$ binding to IGF1 and subsequent $\alpha 6 \beta 4$ -IGF1-IGF1R ternary complex formation in anchorage-independent conditions. *The Journal of biological chemistry*. 2012; 287(15):12491–500. [PubMed: 22351760]
 33. Mizushima H, Wang X, Miyamoto S, Mekada E. Integrin signal masks growth-promotion activity of HB-EGF in monolayer cell cultures. *Journal of cell science*. 2009; 122(Pt 23):4277–86. [PubMed: 19887590]
 34. Kopchick JJ, Parkinson C, Stevens EC, Trainer PJ. Growth hormone receptor antagonists: discovery, development, and use in patients with acromegaly. *Endocr Rev*. 2002; 23(5):623–46. [PubMed: 12372843]
 35. Schreiber I, Buchfelder M, Droste M, Forssmann K, Mann K, Saller B, Strasburger CJ. Treatment of acromegaly with the GH receptor antagonist pegvisomant in clinical practice: safety and efficacy evaluation from the German Pegvisomant Observational Study. *Eur J Endocrinol*. 2007; 156(1):75–82. [PubMed: 17218728]
 36. Mori S, Wu CY, Yamaji S, Saegusa J, Shi B, Ma Z, Kuwabara Y, Lam KS, Isseroff RR, Takada YK, Takada Y. Direct Binding of Integrin $\{\alpha\}v\{\beta\}3$ to FGF1 Plays a Role in FGF1 Signaling. *The Journal of biological chemistry*. 2008; 283(26):18066–75. [PubMed: 18441324]
 37. Yamaji S, Saegusa J, Ieguchi K, Fujita M, Takada YK, Takada Y. A novel fibroblast growth factor-1 (FGF1) mutant that acts as an FGF antagonist. *PloS one*. 2010; 5(4):e10273. [PubMed: 20422052]
 38. Mori S, Tran V, Nishikawa K, Kaneda T, Hamada Y, Kawaguchi N, Fujita M, Takada YK, Matsuura N, Zhao M, Takada Y. A Dominant-Negative FGF1 Mutant (the R50E Mutant) Suppresses Tumorigenesis and Angiogenesis. *PloS one*. 2013; 8(2):e57927. [PubMed: 23469107]
 39. Ieguchi K, Fujita M, Ma Z, Davari P, Taniguchi Y, Sekiguchi K, Wang B, Takada YK, Takada Y. Direct binding of the EGF-like domain of neuregulin-1 to integrins ($\{\alpha\}v\{\beta\}3$ and $\{\alpha\}6\{\beta\}4$) is involved in neuregulin-1/ErbB signaling. *The Journal of biological chemistry*. 2010; 285(41):31388–98. [PubMed: 20682778]
 40. Fujita M, Takada YK, Takada Y. Integrins $\alpha v \beta 3$ and $\alpha 4 \beta 1$ Act as Coreceptors for Fractalkine, and the Integrin-Binding Defective Mutant of Fractalkine Is an Antagonist of CX3CR1. *J Immunol*. 2012; 189(12):5809–5819. [PubMed: 23125415]

Biographies

Yoshikazu Takada:

I obtained MD and PhD degrees in Japan, and studied liver peroxisomes in animals for 10 years, I moved to Dana-Farber cancer institute in Boston in 1986 and was involved in identification of $\beta 1$ integrins. I moved to the Scripps Research Institute in La Jolla in 1990,

and then UC Davis School of Medicine (2003-present) and at the PhD program for translational medicine, Taipei Medical University. Current interest is integrin-growth factor crosstalk and integrin signaling. Using docking simulation, we identified several new integrin ligands including FGF, IGF, neuregulin, fractalkine, secreted phospholipase A2 type IIA, and C-reactive protein. Our studies aim to identify new integrin functions that are not related cell adhesion to ECM.

Masaaki Fujita:

I studied chemistry, structural biology, and protein engineering at Tokyo Institute of Technology in Tokyo, Japan. I then completed a MD/PhD program at Kobe University School of Medicine, Kobe Japan. I conducted clinical and basic research in the field of rheumatology and immunology during residency. I spent several years as a postdoctoral scholar and then assistant researcher at UC Davis, School of Medicine, Sacramento, USA. I was involved in the project on integrin-growth factor crosstalk during this period. Currently, I am a physician scientist (vice chair of Department of Clinical Immunology and Rheumatology) at The Tazuke-Kofukai Medical Research Institute, Kitano Hospital in Osaka, Japan. I intend to connect basic science and medicine and advance bench to bedside research.

Highlights

- IGF1 directly binds to integrins and induces ternary complex formation (integrin, IGF1, and IGF1R) on the cell surface. The integrin-binding defective IGF1 mutant is defective in inducing IGF signaling, although the mutant still binds to IGF1R.
- The integrin-binding defective IGF1 mutant is dominant-negative and suppresses cell proliferation induced by wt IGF1, and suppresses tumorigenesis in vivo.
- IGF1 signaling is detected in anchorage-independent conditions (in the absence of integrin-ECM interaction), suggesting that integrin-IGF1 interaction, rather than integrin-ECM interaction, is important for IGF signaling.
- In addition to IGF1, several other cytokines (FGF, neuregulin-1, fractalkine) directly bind to integrins and induce ternary complex formation, suggesting that integrins act as co-receptors common to these cytokines.

Fig. 1.

A model of IGF signaling. IGF1 binds to IGF1R, and then integrin $\alpha\beta3$ (and $\alpha6\beta4$ and perhaps other integrins as well) is recruited to the IGF1- IGF1R complex. a. Our studies found that WT IGF1 directly binds to integrins and induces ternary complex formation on the cell surface. The integrin-binding defective mutant (R36E/R37E) of IGF1 is defective in inducing signals and ternary complex formation while the mutant still binds to IGF1R. b. IGF1 binds first its cognate high-affinity receptor IGF1R, and then integrins are recruited to the IGF1/IGF1R complex, resulting the formation of integrin-IGF1-IGF1R ternary complex. The integrin-binding defective IGF1 mutant (R36E/R37E, IGF1 decoy) generates an inactive IGF1R/IGF1 decoy complex upon binding to IGF1R, and integrins cannot be recruited to the IGF1R/IGF1 decoy complex. c. Does ternary complex formation require IGF1R activation? Our studies found that PPP, a specific IGF1R inhibitor, did not block the ternary complex formation, indicating that integrin-IGF1-IGF1R ternary complex formation does not require IGF1R phosphorylation. We speculate that ternary complex formation happens before IGF1R is phosphorylated. P= phosphate.

Fig. 2.

R36E/R37E suppresses cell viability and tumorigenesis, whereas WT IGF1 enhances cell viability and tumorigenesis. Met-1 cells that secrete WT or IGF1 decoy (R36E/R37E) were injected into the mammary fat pads of FVB mice. WT IGF1 markedly enhanced tumor growth, but IGF1 decoy delayed tumor growth. b. A potential mechanism of the antagonistic action of IGF1 decoy. IGF1 decoy competes with WT IGF1 for binding to IGF1R and suppresses the ternary complex formation on the cell surface, resulting in the inactive IGF1R/IGF1 decoy complex.