Hydration-reduced lattice thermal conductivity of olivine in Earth's upper mantle

Yun-Yuan Chang^a, Wen-Pin Hsieh^{a,1}, Eh Tan^a, and Jiuhua Chen^{b,c}

^aInstitute of Earth Sciences, Academia Sinica, Taipei 11529, Taiwan; ^bCenter for High Pressure Science and Technology Advanced Research, Changchun 130012, China; and ^cCenter for the Study of Matter at Extreme Conditions, Florida International University, Miami, FL 33191

Edited by Ho-kwang Mao, Carnegie Institution of Washington, Washington, DC, and approved March 16, 2017 (received for review September 28, 2016)

Earth's water cycle enables the incorporation of water (hydration) in mantle minerals that can influence the physical properties of the mantle. Lattice thermal conductivity of mantle minerals is critical for controlling the temperature profile and dynamics of the mantle and subducting slabs. However, the effect of hydration on lattice thermal conductivity remains poorly understood and has often been assumed to be negligible. Here we have precisely measured the lattice thermal conductivity of hydrous San Carlos olivine ($\text{Mg}_{0.9}\text{Fe}_{0.1}\text{SiO}_4$ (Fo90) up to 15 gigapascals using an ultrafast optical pump–probe technique. The thermal conductivity of hydrous Fo90 with $\sim 7,000$ wt ppm water is significantly suppressed at pressures above ~ 5 gigapascals, and is approximately 2 times smaller than the nominally anhydrous Fo90 at mantle transition zone pressures, demonstrating the critical influence of hydration on the lattice thermal conductivity of olivine in this region. Modeling the thermal structure of a subducting slab with our results shows that the hydration-reduced thermal conductivity in hydrated oceanic crust further decreases the temperature at the cold, dry center of the subducting slab. Therefore, the olivine–wadsleyite transformation rate in the slab with hydrated oceanic crust is much slower than that with dry oceanic crust after the slab sinks into the transition zone, extending the metastable olivine to a greater depth. The hydration-reduced thermal conductivity could enable hydrous minerals to survive in deeper mantle and enhance water transportation to the transition zone.

hydration | thermal conductivity | geodynamics | metastable olivine | subducting slab

H₂O plays a critical role in driving and affecting many geophysical phenomena and dynamic processes in Earth's interior (1–3). It has been suggested that olivine, a primary mineral in the upper mantle, and its high-pressure polymorphs (wadsleyite and ringwoodite) could store a large amount of water (hydrogen ions) in their crystalline defects and act as a major water reservoir within Earth (2–4). Incorporation of water in these mantle minerals has been shown to influence their physical properties (5–12) and the dynamics of the mantle and subducting slabs (13). In particular, water enrichment could influence the minerals' thermal transport properties, which, in turn, alters the temperature gradient in the mantle and subducting slabs. At the center of subducting slabs, the cold temperature inhibits the olivine–wadsleyite/ringwoodite phase transformation (14). A thin wedge of olivine could therefore persist in a metastable state far below the 410-km depth. Previous experiments on transformation kinetics have shown that the presence of water inside the slabs greatly enhances the olivine–wadsleyite/ringwoodite transformation rate under the same pressure and temperature conditions (15–18). However, the hydration effect on the lattice thermal conductivity of mantle minerals and its consequential influence on the temperature inside the slabs that also controls the transformation rate remain largely unknown. Precise determination of the thermal conductivity of hydrous upper mantle minerals at relevant conditions would provide critical insight into the temperature evolution and dynamics of subducting slabs, and therefore the fate of metastable olivine in the mantle

transition zone, with potential implications for the origin of deep earthquakes (19).

In the past few decades, the lattice and radiative thermal conductivities of upper mantle minerals have been extensively studied, as these two mechanisms transfer heat through Earth's upper mantle in different ways. Previous optical absorption measurements have indicated that, under relevant pressure–temperature (*P*–*T*) conditions in the upper mantle, small amounts of incorporated water may strongly influence the radiative thermal conductivities of olivine (20), wadsleyite (21), and ringwoodite (21). By contrast, precise measurements of the lattice thermal conductivity of upper mantle minerals under extreme *P*–*T* conditions are very challenging. Therefore, little experimental data are available and the measurement accuracy is insufficient due to the limitations of previous experimental methods. Although the lattice thermal conductivity and diffusivity of nominally anhydrous olivine have been investigated for decades (22–25), the reported results at ambient and high pressures remain inconclusive among different studies. Moreover, in contrast to radiative thermal conductivity, which is sensitive to the hydration content, the role that hydration plays in the lattice thermal conductivity of olivine has often been assumed to be negligible (20, 25) due to its limited water storage capacity, typically of the order of ~ 0.1 wt% to 1 wt%. The validity of this hypothesis, however, has not been experimentally tested, and whether the influence of hydration changes as pressure increases remains unknown. Our study of the hydration effect on the olivine lattice thermal conductivity under relevant mantle conditions, in combination with our joint thermomechanical modeling of a subducting slab, provides insight into the extent of metastable olivine in the mantle transition zone.

Significance

Thermal conductivity of mantle minerals is critical for controlling the temperature profile and dynamics of the mantle and subducting slabs. However, the effect of hydration on lattice thermal conductivity remains poorly understood. We studied lattice thermal conductivity of olivine ($\text{Mg}_{0.9}\text{Fe}_{0.1}\text{SiO}_4$ (Fo90) to 15 GPa using ultrafast optics. The thermal conductivity of hydrous Fo90 with $\sim 7,000$ wt ppm water is 2 times smaller than its anhydrous counterpart at transition zone pressures. Modeling thermal structure of a subducting slab shows that the hydration-reduced thermal conductivity in the oceanic crust further decreases the temperature within the subducting slab, which substantially lowers the olivine–wadsleyite transformation rate and extends the metastable olivine to greater depths. Such an effect could enhance water transportation to the transition zone.

Author contributions: W.-P.H. designed research; Y.-Y.C., W.-P.H., and E.T. performed research; Y.-Y.C. and W.-P.H. analyzed data; J.C. synthesized samples; and Y.-Y.C., W.-P.H., E.T., and J.C. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

¹To whom correspondence should be addressed. Email: wphsieh@earth.sinica.edu.tw.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1616216114/-DCSupplemental.

Results and Discussion

We combined ultrafast time-domain thermoreflectance (TDTR) with high-pressure diamond anvil cell (DAC) techniques to precisely measure the lattice thermal conductivity (Λ) of anhydrous (polycrystalline) and hydrous (single crystal) San Carlos olivine ($\text{Mg}_{0.9}\text{Fe}_{0.1}\text{SiO}_4$ (Fo90)) at room temperature and high pressures up to ~ 15 GPa. The results are shown in Fig. 1. Given the anisotropic thermal conductivity along different crystal orientations (23), the pressure-dependent thermal conductivity of our anhydrous polycrystalline Fo90 (Λ_{anhy} ; black symbols in Fig. 1) shows a slight variation comparable to the measurement uncertainty of about 10 to 15% (see [Supporting Information](#) for the details). At ambient pressure, $\Lambda_{\text{anhy}} \approx 3.6 \text{ W}\cdot\text{m}^{-1}\cdot\text{K}^{-1}$, in reasonable agreement with previous measurements of polycrystalline Fo90 (22) and slightly lower than single-crystal Fo92 (25) and Fo93 (23), suggesting that, due to the iron mass disorder effect (cf. ref. 26), the more iron ions in the olivine, the stronger the phonon-defect scattering, leading to a lower lattice thermal conductivity. Upon compression, Λ_{anhy} increases rapidly with increasing pressure to $\sim 11.6 (\pm 2.3) \text{ W}\cdot\text{m}^{-1}\cdot\text{K}^{-1}$ at 13.1 GPa, a pressure near the 410-km discontinuity.

Compared with the anhydrous Fo90, the lattice thermal conductivity of the hydrous single crystal Fo90 (Λ_{hy}) shows a weaker pressure dependence (see the red symbols in Fig. 1). Note that a different set of red symbols represents the measurement along a different orientation of the single-crystal sample. As expected, the incorporation of $\sim 7,000$ wt ppm (0.7 wt%) water in Fo90 does not significantly change the lattice thermal conductivity at ambient pressure or even below $P \approx 5$ GPa (the characterization of water content is described in [Calibration of Water Content](#) and [Figs. S2 and S3](#)). However, the Λ_{hy} remains low upon further compression and reaches only ~ 5.7 (± 0.6) $\text{W}\cdot\text{m}^{-1}\cdot\text{K}^{-1}$ at 13.6 GPa, a factor of

Fig. 1. Lattice thermal conductivity of anhydrous (black symbols) and hydrous (red symbols) San Carlos olivine Fo90 as a function of pressure at room temperature. Each set of symbols represents an individual measurement run, with solid symbols for compression and open symbols for decompression cycles. The effects of hydration and pressure on the thermal conductivity of Fo90 incorporated with 7,000 wt ppm water are insignificant until ~ 5 GPa, after which its increase with pressure is suppressed and becomes much smaller than its anhydrous counterpart by a factor of 2 at the pressures near the mantle transition zone, which is marked by the blue dashed line. The orange cross symbols represent the conductivity of Fo90 with a water content lower than 200 wt ppm, which is similar to the anhydrous Fo90 until ~ 10 GPa. The measurement uncertainties are $\sim 10\%$ in the low-pressure regime and $\sim 15\%$ at 15 GPa. Literature results for polycrystalline anhydrous Fo90 (green symbols) at 298 K from ref. 22 are plotted for comparison.

2 smaller than the Λ_{anhy} at a similar pressure. The concentrated distribution of thermal conductivity as a function of pressure based on four sets of measurements along different orientations suggests that the anisotropy of the hydrous Fo90 thermal conductivity may have been suppressed by the hydration effect under high pressure. The orange cross symbols in Fig. 1 represent the Λ_{hy} of Fo90 incorporated with a water content lower than 200 wt ppm (see [Calibration of Water Content](#) and [Fig. S2](#)), which is essentially the same as the Λ_{anhy} in the low-pressure regime but becomes slightly smaller by ~ 20 to 25% at $P > 10$ GPa. Note that, because the variation in the thermal conductivity during 8 GPa to 11 GPa is within our measurement uncertainty of about 15%, there isn't a thermal conductivity anomaly within such a pressure range. Compared with the effect of proton weakening on the sound velocities and thermodynamic properties (e.g., heat capacity) of olivine and its high-pressure polymorphs [typically of the order of only a few percent (5, 11, 12)], the proton-induced lattice thermal conductivity reduction at high pressures, by a factor as large as 2, is dramatic and much more significant than previously expected. Such a large pressure-induced suppression of hydrous Fo90 lattice thermal conductivity is likely due to higher-density ionic defects when compressed to higher pressures. Such weaker pressure dependence of thermal conductivity was also reported in MgO doped with ionic defects of iron at high pressures (26). At low-pressure regime, the replacement of cations (Si^{4+} , Mg^{2+} , and Fe^{2+}) in Fo90 with a small amount of H^+ does not substantially affect the heat capacity, phonon group velocities, and phonon scattering rates. As pressure increases, these ionic defects may aggregate and form a higher-density network that serves as an additional phonon-defect scattering mechanism to substantially reduce the phonon mean free path for thermal energy transport.

In Fig. 2, we plot the ratio of our lattice thermal conductivity of hydrous Fo90 against its anhydrous counterpart ($\Lambda_{hy}/\Lambda_{anhy}$) as a function of pressure to characterize the influences of hydration and pressure on thermal conductivity. The decrease of this ratio with pressure indicates an enhancement of the hydration effect with increasing pressure, which could be general for other hydrous mantle minerals.

In addition to the lattice thermal conductivity, which increases with pressure but decreases with temperature, the radiative thermal conductivity (Λ_{rad}) of olivine is important as well for heat transfer in Earth's upper mantle, particularly at high-temperature conditions. Hofmeister (20) showed that the Λ_{rad} of Fo90 is sensitive to the hydration content and grain size of a crystal. At 1,500 K, the Λ_{rad} of subcentimeter-sized Fo90 incorporated with 10 wt ppm to 100 wt ppm water is smaller than its anhydrous counterpart and the lattice thermal conductivity by a factor of ~ 4 and ~ 8 , respectively, if the hydration effect on the lattice thermal conductivity was assumed to be negligible (cf. figures 5 and 6 in ref. 20). If we assume that the temperature dependence of hydrous Fo90 lattice thermal conductivity (Λ_{hy}), which is incorporated with proton defects follows a typically assumed $T^{-1/2}$ dependence (see our high-temperature measurements at ambient pressure in [High-Temperature Thermal Conductivity Measurements](#), Fig. S6, and Table S1 as well as refs. 22 and 26), the Λ_{hy} of Fo90 with $\sim 7,000$ wt ppm water would then be reduced to $\sim 2.5 \text{ W m}^{-1} \text{ K}^{-1}$ at 13.6 GPa and 1,500 K. This estimation suggests that the lattice thermal conductivity of hydrous olivine at relevant P - T conditions near the mantle transition zone remains significantly larger than the radiative component, indicating that thermal conduction is a major contributor to heat transfer in the hydrous mantle, compared with the thermal radiation mechanism. Therefore, it is reasonable to use the lattice thermal conductivity to evaluate the effect of hydration on the thermal profile of subducting metastable olivine slabs in Earth's transition zone.

We further investigated the resultant temperature anomaly of subducting slabs using 2D thermomechanical models (27) (see [Supporting Information](#) for the modeling details). The degree of hydration below the oceanic Moho before subduction remains

Fig. 2. The ratio of the lattice thermal conductivity of hydrous Fo90 to its anhydrous counterpart as a function of pressure. The red dashed line represents the 7,000 wt ppm water case, and the black dashed line represents the low water content case. Here, $\Lambda_{anh}/(\Lambda_{hy})$ was obtained by averaging the data in three (four) different measurement runs shown as the black (red) symbols in Fig. 1. The uncertainties of the ratio are typically $\sim 15\%$ in the low-pressure regime and $\sim 15\%$ to 20% at $P \approx 13.6$ GPa. This behavior features the effect of hydration, which enhances with increasing pressure.

poorly constrained, but the bending of oceanic plates during subduction could create extensional stress and favorable conditions for pumping water into the mantle (28). Extensional earthquakes in the Western Aleutians and Southern Bonin subduction zones can occur at 20 km below the crust, suggesting partial serpentinization at this depth (29). Seismic exploration has found a 12-km-thick serpentinized mantle beneath the Cocos plate offshore Costa Rica (30), although it is still unclear whether this amount of hydration layer is universal or not. We, therefore, tested three cases with different degrees of hydration. Case A is the reference case with anhydrous oceanic crust and mantle, which have constant thermal conductivity. Case B contains a hydrous oceanic crust and an anhydrous mantle. Case C contains the hydrous oceanic crust as in case B, and an additional 20-km-thick hydrous mantle right below oceanic Moho. The hydrous crust in case B as well as the hydrous crust and mantle in case C are assumed to contain 7,000 wt ppm of water and have reduced thermal conductivity as we experimentally found at high pressure and room temperature.

The slabs in our models have an age of 80 My and a plate velocity of $5 \text{ cm}\cdot\text{y}^{-1}$ at the trench. After the slabs, defined as the region 150°C colder than normal mantle, reach the transition zone, their thicknesses are $\sim 150 \text{ km}$. The temperature in the center of the slab is colder than the surrounding mantle by 560°C to 590°C at a 600-km depth (Fig. 3A). This temperature anomaly is $\sim 200^\circ\text{C}$ smaller than that used in other studies (16, 18). In those studies, the slab temperature was assumed to be 873 K at 660 km and had a constant thermal gradient of $0.6 \text{ K}\cdot\text{km}^{-1}$. Using faster velocity and an older age increases the temperature anomaly in our models. However, the temperature gradient in our models evolves dynamically and changes with depth, making a direct comparison with those studies difficult. The hydrous oceanic crust acts as a thermal insulator in case B and lowers the slab temperature by 10°C in all depth ranges, compared with case A. Compared with case B, an additional 20-km layer of thermally insulating mantle in case C has a less profound effect at shallow depths but still lowers the slab temperature by another 10°C at a 600-km depth. The distance from Moho to the coldest part of the slab at this 600-km depth is about

22 km for case A, 19 km for case B, and 17 km for case C, indicating an increased effect of thermal insulation from hydrous rocks.

The persistence of the metastable olivine wedge is sensitive to temperature and hydration and was used to constrain the water content in the slab. We further computed the olivine–wadsleyite phase growth rate and transformed volume fraction (16, 19) inside the cold slab (Fig. 3B). The growth rate in case B is $\sim 2/3$ of the rate in case A throughout the transition zone, and the depth of 50% transformed fraction in case B is 17 km deeper than in case A. The coldest region in case C is inside the hydrous zone. Because the growth rate is greatly enhanced by the high hydration content locally and is mildly inhibited by the lower temperature, the growth rate in case C is several orders of magnitude higher, so that the phase transformation completes at the equilibrium depth. Although the exact amount of transformed fraction might be affected by other unaccounted factors, importantly, we found that even though case B contains hydrous oceanic crust because the cold center of the slab is anhydrous, a significant fraction of metastable olivine still exists below a 500-km depth. The hydration-reduced thermal conductivity deepens the stability field of metastable olivine in case B compared with the case of anhydrous oceanic crust in case A. We also performed thermo-mechanical modeling with the $T^{-1/2}$ dependence for the thermal conductivity of mantle minerals at relevant P - T conditions and obtained consistent results of lower temperature inside the slab due to hydration (Fig. S7 and [Supporting Information](#)).

In contrast to water's effect that increases the olivine-wadsleyite transformation rate via a transformation kinetics mechanism (15–18), our findings here suggest that a hydrous oceanic crust, which is located near the edge of the slab and has lower thermal conductivity, will lower the temperature inside the slab. The colder temperature slows down the transformation rate, allowing metastable olivine at the center of the slab to exist at greater depths than previously thought. The metastable olivine is compatible with, and even benefits from, a hydrous oceanic crust, and its existence is not

Fig. 3. Temperature anomaly and the olivine–wadsleyite transformed fraction of the slabs. (A) The mantle and 7-km-thick oceanic crust in case A are anhydrous. The mantle in case B is anhydrous, but the oceanic crust is hydrated with reduced thermal conductivity as shown in Fig. 2. Case C is similar to case B but with an additional 20-km hydrated layer of mantle beneath the oceanic crust. The simulation ran for 60 My to reach a thermally steady state statistically. The temperature anomaly in the coldest part of the slab is averaged for the last 10 My, and the shaded region is the min/max variation in the temperature anomaly over the time span. (B) The olivine–wadsleyite transformed fraction in the coldest part of the slab. The coldest part of the slab in cases A and B is anhydrous and can sustain metastable olivine. In case C, the coldest part of the slab is within the hydrated mantle layer. With enhanced growth rate, olivine completely transforms to wadsleyite at equilibrium depth.

necessarily linked to a completely dry condition in all parts of the slab. In addition, the hydrated slab will slow down the heat dissipation during frictional sliding of earthquakes, and therefore promotes transformational deep earthquakes.

Although the water content in the entire slab had been suggested to vary widely, depending on the slab's age and spreading rate (cf. ref. 31), our data for the thermal conductivity of hydrous Fo90 with <200 wt ppm H₂O (orange cross symbols in Fig. 1) already showed an about 25% lower value than the anhydrous Fo90 at about 14 GPa. We therefore can expect that, even if the hydrous olivine in the oceanic crust contains water content of only a few hundred to 1,000 wt ppm, it will still play an effective role in the thermal insulating effect. In other words, the possible low water content in olivine in the oceanic crust still lowers the temperature at the center of the subducting slab and slows down the olivine–wadsleyite/ringwoodite transformation rate around the 410-km depth.

Conclusions

The hydration-induced significant reduction of lattice thermal conductivity at higher pressures could be a general feature in other mantle minerals, such as garnet, wadsleyite, and ringwoodite, which can contain comparable or even larger amounts of hydration than

olivine (5). Therefore, the effective lattice thermal conductivities of the hydrous subducted slab, upper mantle, and transition zone aggregates are likely much smaller than previously expected. The stability of hydrous minerals is strongly affected by temperature, and lower thermal conductivity can help hydrous minerals in subducted slabs survive to greater depths, enhancing water transportation to the transition zone, which will lower the thermal conductivity further. There is potentially positive feedback between the water concentration and reduced thermal conductivity in the transition zone. Future experimental studies of the hydration effect on the lattice thermal conductivity of other mantle minerals at relevant P - T conditions is essential to further constrain the heat transfer and to model the dynamics in Earth's upper mantle and transition zone.

ACKNOWLEDGMENTS. J.C. acknowledges Dr. Bin Yang and Dr. Chunli Ma for their assistance during sample preparation. The work by W.-P.H. was supported by the Academia Sinica and the Ministry of Science and Technology (MOST) of Taiwan, Republic of China, under Contract MOST 103-2112-M-001-001-MY3 and 105-2116-M-001-024. Y.-Y.C. was supported under MOST 103-2811-M-001-182 and 104-2811-M-001-179. E.T. was supported under Academia Sinica AS-104-TP-A04. J.C. acknowledges the National Science Foundation Award EAR-1015509, and the Key State Laboratory for Inorganic Synthesis & Preparative Chemistry. This work was also partially supported by the National Natural Science Foundation of China (Grant U1530402).

1. Nestola F, Smyth JR (2016) Diamonds and water in the deep Earth: A new scenario. *Int Geol Rev* 58:263–276.
2. Hirschmann MM, Kohlstedt D (2012) Water in Earth's mantle. *Phys Today* 65:40–45.
3. Hirschmann MM (2006) Water, melting, and the deep Earth H₂O cycle. *Annu Rev Earth Planet Sci* 34:629–653.
4. Pearson DG, et al. (2014) Hydrous mantle transition zone indicated by ringwoodite included within diamond. *Nature* 507:221–224.
5. Jacobsen SD (2006) Effect of water on the equation of state of nominally anhydrous minerals. *Rev Mineral Geochem* 62:321–342.
6. Hirth G, Kohlstedt DL (1996) Water in the oceanic upper mantle: implications for rheology, melt extraction and the evolution of the lithosphere. *Earth Planet Sci Lett* 144:93–108.
7. Chen J, Inoue T, Yurimoto H, Weidner D (2002) Effect of water on olivine-wadsleyite phase boundary in the (Mg, Fe)₂SiO₄ system. *Geophys Res Lett* 29:1875.
8. Kelbert A, Schultz A, Egbert G (2009) Global electromagnetic induction constraints on transition-zone water content variations. *Nature* 460:1003–1006.
9. Karato S (1990) The role of hydrogen in the electrical conductivity of the upper mantle. *Nature* 347:183–187.
10. Gardes E, Gaillard F, Tarits P (2014) Toward a unified hydrous olivine electrical conductivity law. *Geochem Geophys Geosyst* 15:4984–5000.
11. Jacobsen SD, et al. (2008) Effects of hydration on the elastic properties of olivine. *Geophys Res Lett* 35:6–11.
12. Jahn S, Rahner R, Dachs E, Mrosko M, Koch-Müller M (2013) Thermodynamic properties of anhydrous and hydrous wadsleyite, β -Mg₂SiO₄. *High Pressure Res* 33:584–594.
13. Crowley JW, Gérault M, O'Connell RJ (2011) On the relative influence of heat and water transport on planetary dynamics. *Earth Planet Sci Lett* 310:380–388.
14. Mosenfelder JL, Marton FC, Ross CR, Kerschhofer L, Rubie DC (2001) Experimental constraints on the depth of olivine metastability in subducting lithosphere. *Phys Earth Planet Inter* 127:165–180.
15. DuFrane WL, Sharp TG, Mosenfelder JL, Leinenweber K (2013) Ringwoodite growth rates from olivine with ~75ppmw H₂O: Metastable olivine must be nearly anhydrous to exist in the mantle transition zone. *Phys Earth Planet Inter* 219:1–10.
16. Hosoya T, Kubo T, Ohtani E, Sano A, Funakoshi KI (2005) Water controls the fields of metastable olivine in cold subducting slabs. *Geophys Res Lett* 32:L17305.
17. Kubo T, Ohtani E, Kato T, Shinmei T, Fujino K (1998) Effects of water on the alpha-beta transformation kinetics in San Carlos olivine. *Science* 281:85–87.
18. Perrillat JP, et al. (2013) Mechanism and kinetics of the α - β transition in San Carlos olivine Mg_{1.8}Fe_{0.2}SiO₄. *J Geophys Res Earth* 118:110–119.
19. Kirby SH, Stein S, Okal EA, Rubie DC (1996) Metastable mantle phase transformations and deep earthquakes in subducting oceanic lithosphere. *Rev Geophys* 34:261–306.
20. Hofmeister AM (2004) Enhancement of radiative transfer in the upper mantle by OH⁻ in minerals. *Phys Earth Planet Inter* 146:483–495.
21. Thomas S-M, Bina CR, Jacobsen SD, Goncharov AF (2012) Radiative heat transfer in a hydrous mantle transition zone. *Earth Planet Sci Lett* 357:358:130–136.
22. Xu Y, et al. (2004) Thermal diffusivity and conductivity of olivine, wadsleyite and ringwoodite to 20 GPa and 1373 K. *Phys Earth Planet Inter* 143:321–336.
23. Osako M, Ito E, Yoneda A (2004) Simultaneous measurements of thermal conductivity and thermal diffusivity for garnet and olivine under high pressure. *Phys Earth Planet Inter* 143:311–320.
24. Gibert B (2003) Thermal diffusivity of olivine single-crystals and polycrystalline aggregates at ambient conditions—A comparison. *Geophys Res Lett* 30:2172.
25. Pertermann M, Hofmeister AM (2006) Thermal diffusivity of olivine-group minerals at high temperature. *Am Mineral* 91:1747–1760.
26. Dalton DA, Hsieh W-P, Hohensee GT, Cahill DG, Goncharov AF (2013) Effect of mass disorder on the lattice thermal conductivity of MgO periclase under pressure. *Sci Rep* 3:2400.
27. Tan E, Lavier LL, VanAvendonk HJA, Heurt A (2012) The role of frictional strength on plate coupling at the subduction interface. *Geochem Geophys Geosyst* 13:Q10006.
28. Faccenda M, Gerya TV, Burlini L (2009) Deep slab hydration induced by bending-related variations in tectonic pressure. *Nat Geosci* 2:790–793.
29. Emry EL, Wiens DA (2015) Incoming plate faulting in the Northern and Western Pacific and implications for subduction zone water budgets. *Earth Planet Sci Lett* 414:176–186.
30. VanAvendonk HJA, Holbrook WS, Lizarralde D, Denyer P (2011) Structure and serpentinization of the subducting Cocos plate offshore Nicaragua and Costa Rica. *Geochem Geophys Geosyst* 12:Q06009.
31. Faccenda M (2014) Water in the slab: A trilogy. *Tectonophysics* 614:1–30.
32. Mao HK, Bell PM, Shaner JW, Steinberg DJ (1978) Specific volume measurements of Cu, Mo, Pd, and Ag and calibration of the ruby R1 fluorescence pressure gauge from 0.06 to 1 Mbar. *J Appl Phys* 49:3276–3283.
33. Paterson M (1982) The determination of hydroxyl by infrared absorption in quartz, silicate glasses and similar materials. *Bull Mineral* 105:20–29.
34. Bolfan-casanova N, Keppler H, Rubie DC (2000) Water partitioning between nominally anhydrous minerals in the MgO–SiO₂–H₂O system up to 24 GPa: Implications for the distribution of water in the Earth's mantle. *Earth Planet Sci Lett* 182:209–221.
35. Libowitzky E, Rossman GR (1997) An IR absorption calibration for water in minerals. *Am Mineral* 82:1111–1115.
36. Bell DR, Rossman GR, Maldener J, Endisch D, Rauch F (2003) Hydroxide in olivine: A quantitative determination of the absolute amount and calibration of the IR spectrum. *J Geophys Res* 108:2105.
37. Demouchy S, Bolfan-Casanova N (2016) Distribution and transport of hydrogen in the lithospheric mantle: A review. *Lithos* 240–243:402–425.
38. Cahill DG (2004) Analysis of heat flow in layered structures for time-domain thermoreflectance. *Rev Sci Instrum* 75:5119–5122.
39. Kang K, Koh YK, Chiristescu C, Zheng X, Cahill DG (2008) Two-tint pump-probe measurements using a femtosecond laser oscillator and sharp-edged optical filters. *Rev Sci Instrum* 79:114901.
40. Ge Z, Cahill DG, Braun PV (2006) Thermal conductance of hydrophilic and hydrophobic interfaces. *Phys Rev Lett* 96:186101.
41. Schmidt A, Chiesa M, Chen X, Chen G (2008) An optical pump-probe technique for measuring the thermal conductivity of liquids. *Rev Sci Instrum* 79:064902.
42. Zheng X, Cahill DG, Krasnochtchekov P, Averbach RS, Zhao JC (2007) High-throughput thermal conductivity measurements of nickel solid solutions and the applicability of the Wiedemann-Franz law. *Acta Mater* 55:5177–5185.
43. Chen B, Hsieh W-P, Cahill DG, Trinkle DR, Li J (2011) Thermal conductivity of compressed H₂O to 22 GPa: A test of the Leibfried-Schlömann equation. *Phys Rev B* 83:132301.
44. Hsieh W-P, Chen B, Li J, Kelsblins P, Cahill DG (2009) Pressure tuning of the thermal conductivity of the layered muscovite crystal. *Phys Rev B* 80:180302.
45. Hsieh W-P (2015) Thermal conductivity of methanol-ethanol mixture and silicone oil at high pressures. *J Appl Phys* 117:235901.
46. Li L, Wentzcovitch RM, Weidner DJ, DaSilva CRS (2007) Vibrational and thermodynamic properties of forsterite at mantle conditions. *J Geophys Res* 112:B05206.
47. Cahill DG, Watanabe F (2004) Thermal conductivity of isotopically pure and Ge-doped Si epitaxial layers from 300 to 550 K. *Phys Rev B* 70:235322.
48. Cundall PA (1989) Numerical experiments on localization in frictional materials. *Ingenieur-Archiv* 59:148–159.
49. Chen Y, Morgan W (1990) A nonlinear rheology model for mid-ocean ridge axis topography. *J Geophys Res Solid* 95:17,583–17,604.
50. Kubo T, Kaneshima S, Torii Y, Yoshioka S (2009) Seismological and experimental constraints on metastable phase transformations and rheology of the Mariana slab. *Earth Planet Sci Lett* 287:12–23.
51. Turnbull D (1956) Phase changes. *Solid State Phys* 3:225–306.