Published in final edited form as:

Ann Intern Med. 2008 August 19; 149(4): 264–269.

Navigating Language Barriers under Difficult Circumstances

Yael Schenker, MD, Bernard Lo, MD, Katharine M. Ettinger, JD, and Alicia Fernandez, MD

University of California, San Francisco, and California Pacific Medical Center, San Francisco, California

Abstract

The proportion of the U.S. population with limited English proficiency is growing. Physicians often find themselves caring for patients with limited English proficiency in settings with limited language services. There has been little exploration of the decisions physicians face when providing care across language barriers. The authors offer a conceptual framework to aid physicians in thinking through difficult choices about language services and provide responses to common questions encountered in the care of patients with limited English proficiency. Specifically, they describe 4 factors that should inform the decision to call an interpreter (the clinical situation, degree of language gap, available resources, and patient preference), discuss who may be an appropriate interpreter, and offer strategies for when a professional interpreter is not available. The authors use a hypothetical case to illustrate how decisions about language services may evolve over the course of an interaction. This conceptual and practical approach can help clinicians to improve the quality of care provided to patients with limited English proficiency.

Mrs. A., a 60-year-old diabetic woman who speaks Spanish, arrives in the emergency department with a chief symptom of burning with urination. Dr. B. examines her vital signs and finds that Mrs. A. has a fever. Dr. B. speaks very limited Spanish. A hospital interpreter is available, but not for 1 hour. Dr. B. wonders whether he should wait for a professional interpreter or proceed with the care of Mrs. A. using his own limited Spanish-speaking skills.

This scenario is increasingly common. The proportion of the U.S. population with limited English proficiency nearly doubled from 1980 to 2000 and is projected to increase further (1). More than 20 million people in the United States speak little or no English. Federal laws mandate access to language assistance (2, 3), and hospital guidelines recommend routine use of professional interpreters (4–6). Interpreter services, however, are often limited and poorly integrated into clinical practice (6, 7). Clinician second-language skills and responses to language barriers vary widely (8–11). Many physicians feel uncertain about how best to care for patients with limited English proficiency and uneasy about the quality of care they provide (12–16).

Requests for Single Reprints: Alicia Fernandez, MD, University of California, San Francisco, Box 1364, San Francisco, CA 94143; afernandez@medsfgh.ucsf.edu.

Current Author Addresses: Drs. Schenker, Lo, and Fernandez: University of California, San Francisco, Box 1364, San Francisco, CA 94143. Dr. Ettinger: California Pacific Medical Center, 2395 Sacramento Street, San Francisco, CA 94115.

Potential Financial Conflicts of Interest: None disclosed.

Although suggestions have been made about the appropriate use of interpreters (17, 18), there has been little exploration of how to approach decisions about language services, particularly when such services are limited. We analyze common questions and provide a conceptual framework. Specifically, we describe factors that should inform the decision to call an interpreter, discuss who may be an appropriate interpreter, and offer strategies for when a professional interpreter is not available. Although we recognize that cultural barriers accompany language barriers (19), we focus on language services. We use the hypothetical case of Mrs. A. to illustrate how decisions about language services may evolve over time.

What Factors Should Inform the Decision to Call an Interpreter?

In this case, Dr. B. must decide whether to wait for a hospital interpreter, seek some other form of language assistance, or proceed using his own limited Spanish-speaking skills. Four factors must be considered when deciding whether to involve an interpreter: the clinical situation, degree of language gap, available resources, and patient preference. We consider each factor separately, then illustrate how they influence each other and may change during a clinical encounter.

Clinical Situation

Physicians commonly decide whether to use an interpreter on the basis of what information they need to make a diagnosis and provide treatment. In emergent clinical situations, physicians must provide care, even if language assistance is unavailable. In complex clinical scenarios, physicians are more likely to request an interpreter than for seemingly straightforward encounters (15). Although this approach is often reasonable, physicians also must understand the patient's perspective to accurately assess the severity and complexity of the clinical situation and address the patient's concerns (20, 21). For example, in our hypothetical case, Mrs. A. may also be experiencing severe abdominal pain and nausea. Without language assistance, Dr. B. might have missed this key information. Alternatively, Mrs. A. might have feared that her urinary symptoms were a sign of a sexually transmitted disease.

Degree of Language Gap

The need for an interpreter is easily recognized if the patient and physician do not share a common language. However, the degree of language gap is more difficult to assess when the patient or physician is partially fluent in the other's language. In such situations, physicians frequently attempt to get by using limited second-language skills (14, 15). Language skills are often directional, meaning that spoken ability may exceed understanding, or vice versa. For example, Dr. B. may be able to ask simple screening questions (such as "Do you have abdominal pain?") in Spanish, but not understand complex responses. In addition, vocabulary and fluency may be adequate for certain clinical scenarios but not for others. For example, Dr. B.'s vocabulary may be limited to symptoms commonly encountered in the emergency department, such as abdominal pain. Accurate assessment of one's own language skills is difficult. At a minimum, physicians providing care in a second language should be able to follow-up on a patient's responses and assess patient comprehension to avoid underestimating the language gap.

Available Resources

Availability, accessibility, and quality of interpreter services vary widely among institutions (6). Most hospitals have some mechanism to address language barriers for the most common languages, partly in response to laws that forbid discrimination based on primary language (2). However, institutional constraints may affect the type of interpreter available (that is, on-site professional interpreter vs. telephone interpreter vs. bilingual staff member), the wait time for an interpreter, and the quality of the interpreter. In the hypothetical case, Dr. B. has access to an on-site professional interpreter, but the wait time is 1 hour. Decisions to wait for an interpreter must balance the availability of language assistance with the urgency of the clinical situation.

Patient Preference

Physicians frequently make decisions about interpreter use without taking patient preferences into account. Many patients are not aware that they have a legal right to a professional interpreter at no cost (22). It can be challenging to assess patient preference in the setting of language barriers. Even when a physician has some skills in the patient's language, the power differential between physicians and patients may make it difficult for a patient to express a preference for formal language assistance over a physician's partial language skills. However, failure to elicit patient preference about use of an interpreter is associated with decreased patient satisfaction and may lead to poor adherence to treatment plans and overuse of medical services (23–26). Patients should be offered a professional interpreter and be informed that interpreters are free and trained in confidentiality.

Pulling the Factors Together

How might Dr. B. take into account all 4 factors when deciding about language assistance for the patient?

In assessing the language gap, Dr. B. discovered that Mrs. A. spoke very little English, but his Spanish was adequate to ask a few key questions and understand Mrs. A.'s “yes” and “no” responses. After the pertinent review of systems yielded negative results, Dr. B. decided that obtaining laboratory data was the appropriate next step in Mrs. A.'s care. With the help of a nurse who spoke somewhat more Spanish, he explained to Mrs. A., “I'm concerned you may have an infection. The hospital will provide a professional interpreter free of charge, but with a wait of 1 hour. I think it is important to proceed with blood and urine tests in the meantime.” Mrs. A. expressed her understanding and agreement with this plan, a hospital interpreter was called, and a nurse was sent in immediately to draw Mrs. A.'s blood. The patient's blood and urine test results reveal that she has a high blood glucose level and a kidney infection.

Under these circumstances, the initial decision to proceed without a professional interpreter was appropriate. However, the 4 factors informing the decision to request an interpreter will change over the course of an encounter. Because the data suggest that Mrs. A. has pyelonephritis, Dr. B. must now ensure that he has not missed anything in his initial limited history and examination, gather information about drug allergies and recent antibiotic use, explain the diagnosis and treatment to Mrs. A., and address Mrs. A.'s questions and

concerns. In reassessing his own language skills, Dr. B. will probably find that they are no longer adequate. In reassessing Mrs. A.'s preference, he may discover that she would now prefer to speak through a professional interpreter. At this stage, Dr. B. should use the hospital interpreter he requested earlier, assuming that the interpreter is now available.

Mrs. A. is given intravenous fluids and antibiotics, and preparations are made for admission. Mrs. A.'s daughter arrives and offers to serve as an interpreter. The admitting physician, Dr. C., speaks no Spanish. Dr. C. wonders whether she should accept the daughter's offer or call for a professional interpreter.

Who Is an Appropriate Interpreter, and Who Decides?

There are several different kinds of interpreters, each with advantages and disadvantages (Table 1). Again, the clinical situation, degree of language gap, available resources, and patient preference must be considered. Professional interpreters are the most skilled in medical interpretation, the least likely to make errors, and the only type of interpreter associated with overall improvement of care for patients with limited English proficiency (27, 28). However, professional on-site interpreters may not be immediately available, and professional telephone interpreters may feel impersonal or be difficult to use for patients with hearing or speech impairments (29). Although bilingual staff may be convenient and available, their language skills are usually not tested and may be inadequate (30, 31). Having family members serve as interpreters may be congruent with cultural expectations (32). Relatives may also serve as patient advocates and participate in decisions about a patient's care. However, family members often have inadequate language skills, resulting in higher rates of clinically significant errors (28). Family members may also interpret selectively to fit their own beliefs.

Physicians should inform patients with limited English proficiency of the available resources for language assistance and offer a professional interpreter at each major stage of an encounter. Bilingual staff and family members can help convey this information to patients. Physicians should ensure that patients understand the advantages of professional interpreters and not assume that patients prefer to have family members interpret. Once informed, patients may decide what type of interpreter they prefer.

In some clinical situations, however, the patient's choice of an ad hoc interpreter (that is, a family member or acquaintance) may not be appropriate (Table 2). The chosen interpreter might not be competent in either English or the patient's primary language, or the medical situation may be too complex for the interpreter to understand (28). Alternatively, the patient may not feel comfortable discussing a sensitive topic in the presence of a relative or friend.

For example, Dr. C. may have initially respected Mrs. A.'s preference to have her adult daughter interpret, but then discovered that Mrs. A.'s daughter spoke very little English. It is then Dr. C.'s responsibility to interrupt the interaction, explain why a professional interpreter would be preferable, and offer to request one. She might say to Mrs. A.'s daughter, "I really appreciate your help with interpretation, but I fear I am not doing a good job of understanding your mother. I would like to ask a hospital interpreter to assist us, in *addition*

to you, so that I may provide better care for her. Could you please ask your mother if this would be okay?" By recognizing the ad hoc interpreter's efforts and framing the need for a professional interpreter in clinical terms, physicians may continue to respect the patient's preferences while strongly recommending a professional interpreter.

In rare cases, use of an ad hoc interpreter may place the patient at risk for harm, for example, if the ad hoc interpreter is suspected of abusing elderly persons or domestic violence. In such cases, the physician should request a professional interpreter and conduct the interaction without the suspected person present.

The use of young children as interpreters is common but problematic. Asking children to interpret distorts the child–parent relationship and places a substantial emotional strain on the child (33–35). Some states, including Massachusetts and Rhode Island, have passed laws barring minors from serving as interpreters for family members (36). Except in emergency situations, when no alternatives are available, the use of children as interpreters should be avoided.

Mrs. A. is hospitalized. The next morning, the medical team caring for the patient comes to rounds at her bedside at 8 a.m. No one on the team speaks Spanish. No one is available to translate: The Spanish interpreter arrives at 11 a.m., a bilingual nurse went off shift at 7 a.m., and Mrs. A.'s daughter has gone home. The team therefore checks Mrs. A.'s vital signs, does a perfunctory examination, and leaves without speaking to the patient. Mrs. A. cannot tell them that she has developed an itchy rash on her back.

What Is the Appropriate Response If an Interpreter Is Not Available?

Inpatients commonly have severe illnesses or need invasive procedures, and thus must make important treatment decisions. Failure to communicate with a patient throughout hospitalization and at discharge may compromise the patient's care. A daily conversation between an inpatient and the treating physician should be the standard of care. In this case, the language gap is nearly complete and the clinical situation requires language assistance. Yet, Mrs. A. has no opportunity to voice a preference for an interpreter, and consequently the team cannot address her concerns, explain the treatment plan, or evaluate her rash.

When a professional interpreter is not immediately available, clinicians must investigate all possible options for language assistance (Table 1). Limits on duty hours may make it difficult for resident physicians to wait for a professional interpreter to become available. Creativity and advance planning may be required to obtain language assistance. In this scenario, the team could have arranged for interpreter services in advance, used a remote interpreter over the telephone, enlisted the services of a staff member or ad hoc interpreter (for example, by calling a bilingual family member from Mrs. A.'s bedside phone), or changed the time of rounds to coincide with the hospital interpreter's schedule (while ensuring that Mrs. A. received timely attention). Physicians must be willing to accept some personal inconvenience during such cases in order to act in a patient's best interests.

What Can Be Done When Interpreter Services Are Inadequate?

Availability of interpreter services is not always adequate. Recent national surveys indicate that between 38% and 68% of hospitals use professional interpreters (6, 37). More than half of hospitals surveyed reported frequent use of untrained bilingual staff and family members or friends as interpreters. Professional interpreter services may be less available at small or rural institutions, at night or on the weekends, and for less common languages.

A systems approach is necessary to improve language services. Hospitals need minimum communication standards for the care of patients with limited English proficiency, accurate collection of language data, and language services integrated into the flow of patient care. To avoid depending on the initiative of individual physicians, the hospital admissions office could identify patients with limited English proficiency and automatically schedule interpreters for a time that is convenient for the daily rounds (12, 38). Insurers should provide financial incentives for hospitals to meet benchmarks for interpreter services.

Until such changes occur, physicians must also advocate for improved language services. The chief of service or quality improvement department should be informed when access to interpreters is consistently unacceptable. Physicians should notify the risk management department when inadequate access to interpreters compromises patient care. If such attempts to improve language services fail, physicians and patients should be aware that the federal Office for Civil Rights is charged with ensuring that access to health services does not reflect discrimination on the basis of language or national origin. Discrimination complaints filed with the Office for Civil Rights have led to systemic changes in language services at medical centers around the United States (2, 39).

Conclusion

Physicians are increasingly treating patients across language barriers and in settings with limited availability of language services. Clinicians and patients often find themselves navigating language barriers under difficult circumstances. Although improved quality and availability of interpreter services are needed, clinicians are ultimately responsible for ensuring safe and effective communication with patients. A conceptual and practical approach to navigating language barriers can help to improve the quality of care for patients with limited English proficiency.

Acknowledgments

The authors thank Leila Alpers, Robert Brody, Leah Karliner, and Jeff Kohlwes for their comments on an earlier draft of this manuscript.

Grant Support: By the National Institutes of Health (NIH) Career Development Award K23-RR018324-01 (Dr. Fernandez) and NIH Center Grant MH062246 (Dr. Lo), NIH Roadmap Clinical and Translational Sciences Award U01 AI46749 (Dr. Lo), and the Greenwall Foundation (Dr. Lo).

References

1. United States Census Bureau. United States Census 2000. Accessed at www.census.gov/main/www/cen2000.html on 16 June 2008

2. Chen AH, Youdelman MK, Brooks J. The legal framework for language access in healthcare settings: Title VI and beyond. *J Gen Intern Med*. 2007; 22(2):362–7.
3. Perkins, J. Ensuring Linguistic Access in Health Care Settings: An Overview of Current Legal Rights and Responsibilities. Kaiser Commission of Medicaid and the Uninsured. 2003. Accessed at www.kff.org/uninsured/kcmu4131report.cfm on 16 June 2008
4. U.S. Department of Health and Human Services. The Office of Minority Health. National Standards on Culturally and Linguistically Appropriate Services (CLAS). Accessed at www.omhrc.gov/templates/browse.aspx?lvl=2&lvlid=15 on 16 June 2008
5. The Joint Commission 2008 Requirements Related to the Provision of Culturally and Linguistically Appropriate Health Care. The Joint Commission. Apr. 2008 Accessed at www.JointCommission.org/PatientSafety/HLC/HLC_Joint_Commission_Standards.htm on 16 June 2008
6. Wilson-Stronks, A., Galvez, E. Hospitals, Language, and Culture: A Snapshot of the Nation Exploring Cultural and Linguistic Services in the Nations Hospitals: A Report of Findings. The Joint Commission and The California Endowment. 2007. Accessed at www.JointCommission.org/PatientSafety/HLC/ on 16 June 2008
7. Dower, C., Kaiser, J. UCSF Center for the Health Professions: 2007. Improving Language Access in California Hospitals. Accessed at www.futurehealth.ucsf.edu/hwtc/languageaccess.html on 16 June 2008
8. Lee KC, Winickoff JP, Kim MK, Campbell EG, Betancourt JR, Park ER, et al. Resident physicians' use of professional and nonprofessional interpreters: a national survey [Letter]. *JAMA*. 2006; 296:1050–3. [PubMed: 16954482]
9. Kuo DZ, O'Connor KG, Flores G, Minkovitz CS. Pediatricians' use of language services for families with limited English proficiency. *Pediatrics*. 2007; 119:e920–7. [PubMed: 17371933]
10. Schenker Y, Wang F, Selig SJ, Ng R, Fernandez A. The impact of language barriers on documentation of informed consent at a hospital with on-site interpreter services. *J Gen Intern Med*. 2007; 22(2):294–9. [PubMed: 17957414]
11. Karliner LS, Pérez-Stable EJ, Gildengorin G. The language divide. The importance of training in the use of interpreters for outpatient practice. *J Gen Intern Med*. 2004; 19:175–83. [PubMed: 15009797]
12. Regenstien M. Measuring and improving the quality of hospital language services: insights from the Speaking Together collaborative. *J Gen Intern Med*. 2007; 22(2):356–9. [PubMed: 17957425]
13. Gadon M, Balch GI, Jacobs EA. Caring for patients with limited English proficiency: the perspectives of small group practitioners. *J Gen Intern Med*. 2007; 22(2):341–6. [PubMed: 17957422]
14. Burbano O'Leary SC, Federico S, Hampers LC. The truth about language barriers: one residency program's experience. *Pediatrics*. 2003; 111(5 Pt 1):e569–73. [PubMed: 12728111]
15. Diamond, L., Schenker, Y., Curry, L., Bradley, EH., Fernandez, A. Getting by: Under-use of interpreters by resident physicians. Society of General Internal Medicine; Pittsburgh, Pennsylvania: Apr 9-12. 2008
16. Abbe M, Simon C, Angiolillo A, Ruccione K, Kodish ED. A survey of language barriers from the perspective of pediatric oncologists, interpreters, and parents. *Pediatr Blood Cancer*. 2006; 47:819–24. [PubMed: 16615062]
17. Chen, AH., Jacobs, EA. Providing Care to Patients Who Speak Limited English. In: King, Talmadge E., Wheeler, Margaret B., editors. *Medical management of Vulnerable and Underserved Patients: Principles, Practice and Populations*. New York: McGraw-Hill; 2007. p. 265-274.
18. San Francisco: California Academy of Family Physicians; 2007. Addressing Language and Culture: A Practice Assessment for Healthcare Professionals. Accessed at www.familydocs.org/practice-resources/cultural-proficiency.php on 16 June 2008
19. Fernandez A, Schillinger D, Grumbach K, Rosenthal A, Stewart AL, Wang F, et al. Physician language ability and cultural competence. An exploratory study of communication with Spanish-speaking patients. *J Gen Intern Med*. 2004; 19:167–74. [PubMed: 15009796]

20. Cohen AL, Rivara F, Marcuse EK, McPhillips H, Davis R. Are language barriers associated with serious medical events in hospitalized pediatric patients? *Pediatrics*. 2005; 116:575–9. [PubMed: 16140695]
21. Divi C, Koss RG, Schmaltz SP, Loeb JM. Language proficiency and adverse events in US hospitals: a pilot study. *Int J Qual Health Care*. 2007; 19:60–7. [PubMed: 17277013]
22. Grubbs V, Chen AH, Bindman AB, Vittinghoff E, Fernandez A. Effect of awareness of language law on language access in the health care setting. *J Gen Intern Med*. 2006; 21:683–8. [PubMed: 16808766]
23. Baker DW, Hayes R, Fortier JP. Interpreter use and satisfaction with interpersonal aspects of care for Spanish-speaking patients. *Med Care*. 1998; 36:1461–70. [PubMed: 9794340]
24. Apter AJ, Reisine ST, Affleck G, Barrows E, ZuWallack RL. Adherence with twice-daily dosing of inhaled steroids. Socioeconomic and health-belief differences. *Am J Respir Crit Care Med*. 1998; 157:1810–7. [PubMed: 9620910]
25. Carrasquillo O, Orav EJ, Brennan TA, Burstin HR. Impact of language barriers on patient satisfaction in an emergency department. *J Gen Intern Med*. 1999; 14:82–7. [PubMed: 10051778]
26. Wilson E, Chen AH, Grumbach K, Wang F, Fernandez A. Effects of limited English proficiency and physician language on health care comprehension. *J Gen Intern Med*. 2005; 20:800–6. [PubMed: 16117746]
27. Karliner LS, Jacobs EA, Chen AH, Mutha S. Do professional interpreters improve clinical care for patients with limited English proficiency? A systematic review of the literature. *Health Serv Res*. 2007; 42:727–54. [PubMed: 17362215]
28. Flores G, Laws MB, Mayo SJ, Zuckerman B, Abreu M, Medina L, et al. Errors in medical interpretation and their potential clinical consequences in pediatric encounters. *Pediatrics*. 2003; 111:6–14. [PubMed: 12509547]
29. Garcia EA, Roy LC, Okada PJ, Perkins SD, Wiebe RA. A comparison of the influence of hospital-trained, ad hoc, and telephone interpreters on perceived satisfaction of limited English-proficient parents presenting to a pediatric emergency department. *Pediatr Emerg Care*. 2004; 20:373–8. [PubMed: 15179145]
30. Elderkin-Thompson V, Silver RC, Waitzkin H. When nurses double as interpreters: a study of Spanish-speaking patients in a US primary care setting. *Soc Sci Med*. 2001; 52:1343–58. [PubMed: 11286360]
31. Moreno MR, Otero-Sabogal R, Newman J. Assessing dual-role staff-interpreter linguistic competency in an integrated healthcare system. *J Gen Intern Med*. 2007; 22(2):331–5. [PubMed: 17957420]
32. Kuo D, Fagan MJ. Satisfaction with methods of Spanish interpretation in an ambulatory care clinic. *J Gen Intern Med*. 1999; 14:547–50. [PubMed: 10491243]
33. Flores G. The impact of medical interpreter services on the quality of health care: a systematic review. *Med Care Res Rev*. 2005; 62:255–99. [PubMed: 15894705]
34. Flores G. Language barriers to health care in the United States. *N Engl J Med*. 2006; 355:229–31. [PubMed: 16855260]
35. Jacobs B, Kroll L, Green J, David TJ. The hazards of using a child as an interpreter. *J R Soc Med*. 1995; 88:474P–475P.
36. Youdelman MK. The medical tongue: U.S. laws and policies on language access. *Health Aff (Millwood)*. 2008; 27:424–33. [PubMed: 18332499]
37. Hasnain-Wynia, R., Yonek, J., Pierce, D., Kang, R., Hedges Greising, C. Hospital Language Services For Patients with Limited English Proficiency: Results from A National Survey. Health Research and Educational Trust; National Health Law Program. 2006. Accessed at www.hret.org/ on 6 June 2008
38. Schyve PM. Language differences as a barrier to quality and safety in health care: the Joint Commission perspective. *J Gen Intern Med*. 2007; 22(2):360–1. [PubMed: 17957426]
39. Office for Civil Rights. Accessed at <http://hhs.gov/ocr/> on 16 June 2008

Key Summary Points

Four factors must be considered when making decisions about language services: the clinical situation, the degree of language gap, available resources, and patient preference.

As the clinical situation evolves, patient preferences and needs may shift, requiring ongoing assessment of the need for professional language assistance.

Patients should be included in the decision about whether to call an interpreter. As a standard approach, patients should be informed that professional interpreters are available free of cost.

Physicians using their own second-language skills must continuously assess for patient understanding.

When an institution's interpreter services are inadequate, physicians should advocate for improvement.

Minimum communication standards are needed for hospitalized patients with limited English proficiency.

Physicians have a professional responsibility to ensure that patients with limited English proficiency receive similar care to that of patients who speak English.

Table 1
Types of Interpreters*

Type of Interpreter	Advantages	Disadvantages
Physician using his or her own foreign-language skills	Fast and convenient Protects patient confidentiality, as no third party is involved in the interaction Improves rapport and results in more patient-centered interactions	Physicians may not accurately assess their own language skills Risk for miscommunication if language skills are not sufficient May overlook patient preferences about language assistance
Ad hoc interpreter (e.g., adult family member, friend, or another patient)	Convenient Bilingual family members may serve as patient advocates May be congruent with cultural expectations	Inadequate bilingual language and interpreting skills are common and may result in high rates of errors and omissions Communication may be modified to suit the interpreter's agenda May compromise patient privacy and confidentiality Patients may withhold sensitive information Demands of interpreting may conflict with family role
Bilingual staff (e.g., physicians, nurses, medical assistants, clerks, and janitors)	Convenient No added salary costs Staff may be familiar to patients	Language skills are not assessed and may be inadequate Staff cannot perform other duties
Professional interpreter	Less likely to make errors Trained to protect patient privacy and confidentiality Patients may be more likely to share sensitive information	May have to wait for interpreter to be available Telephone interpreter services may be awkward to use Patients may prefer not to involve a stranger in their care May disrupt family relationships

* Adapted from reference 17.

Table 2
When a Patient's Choice of an Interpreter Is Not Appropriate*

The ad hoc interpreter is not competent in English or the patient's language.

The medical situation is too complex for the ad hoc interpreter to understand or translate.

The patient does not feel comfortable discussing sensitive information with the ad hoc interpreter.

The ad hoc interpreter experiences substantial role conflict (e.g., an adult son interpreting for an end-of-life discussion).

The interpreter is a child.

Use of an ad hoc interpreter places the patient at risk for harm (e.g., the interpreter is suspected of elder abuse or domestic violence).

* Including ad hoc interpreters, such as family members and friends.