Published in final edited form as:

Breast J. 2017 March ; 23(2): 200–205. doi:10.1111/tbj.12780.

Breast Cancer in Pregnancy: Avoiding Fetal Harm When Maternal Treatment Is Necessary

Christina N. Cordeiro, MD* and Mary L. Gemignani, MD, MPH†

*Department of Gynecology and Obstetrics, Johns Hopkins School of Medicine, Baltimore, MA

†Breast Service, Department of Surgery, Memorial Sloan Kettering Cancer Center, New York, NY

Abstract

The management of breast cancer during pregnancy poses unique challenges and requires a multidisciplinary approach. In this review, we discuss the treatment of breast cancer in pregnancy and recent updates regarding the safety of surgical and chemotherapeutic treatments, including both oncologic and fetal outcomes.

The treatment of breast cancer during pregnancy mirrors that outside of pregnancy, with a few important differences dictated by the balance of maternal versus fetal health. Overall, surgical treatment, neoadjuvant chemotherapy, and/or adjuvant chemotherapy are feasible in most women during pregnancy. Further research to determine the safety of these therapies in pregnancy-associated breast cancer is warranted.

Keywords

breast cancer; pregnancy; chemotherapy

INTRODUCTION

Pregnancy-associated breast cancer (PABC), also referred to as gestational breast cancer, is defined as breast cancer diagnosed during pregnancy or in the first year postpartum.¹ It has an incidence of 1 in 3000 pregnancies with a median age of diagnosis of 33 years. As women delay childbearing for personal and/or professional reasons, there may be an increased incidence in PABC during pregnancy or before completion of childbearing.² It has been reported that 80% of PABCs are infiltrating ductal carcinoma, 49–84% are estrogen receptor/progesterone receptor negative, 28–58% are HER2/neu overexpressed, and about 67% present with positive lymph nodes.^{2–5} Although PABC was initially thought to have a worse prognosis than non-PABC, this is not true when matched for age and stage.

Corresponding author: Mary L. Gemignani, MD, MPH, Breast Service, Department of Surgery, Memorial Sloan Kettering Cancer Center, 300 East 66th Street, New York, NY 10065, (T) 646 888 5359, (F) 646 888 4921, gemignam@mskcc.org.

CONFLICTS OF INTEREST

The authors have no conflicts of interest to disclose.

Financial Disclosure: Dr. Mary L. Gemignani was funded in part by NIH/NCI Cancer Center Support Grant No. P30 CA008748. The authors have no conflicts of interest to report.

Delay in diagnosis can occur because of a patient's confusion between cancer-related changes with pregnancy-related changes of the breast. Even for trained practitioners, hormonal influences of pregnancy on breast tissue affect the ability to detect a mass during pregnancy. Physical examination of the breast becomes more difficult as the pregnancy progresses, or postpartum if lactation is established.⁵ Patient denial and, potentially, physician reluctance to intervene during pregnancy, may also lead to delayed diagnosis.⁴ Prompt evaluation of any breast mass, with or without biopsy, is warranted. Here we review current recommendations for the diagnosis and treatment of breast cancer in pregnancy, focusing upon recent updates regarding the safety of surgical and chemotherapeutic treatments in pregnancy, including both oncologic and fetal outcomes.

Diagnosis

Ultrasound is the preferred imaging method for breast mass evaluation in pregnancy and is often the first step, allowing for ultrasound-guided biopsy if necessary.⁶ Sensitivity and negative predictive values for ultrasound in the detection of breast malignancy in pregnant patients have both been reported at 100%⁷, although in other studies, sensitivity has been reported as low as 70%.⁸ In this latter study, however, images were reviewed at various institutions, and the authors noted that eight breast cancer cases were incorrectly classified as "probably benign".⁸ Mammography, if needed for further evaluation, has a low risk of radiation (estimated at 4mGy), especially with abdominal shielding. However, mammography is often not as informative because these women are often young with dense breasts.^{2,5,9} Mammography sensitivity for diagnosis has been reported ranging from 25–75% in some reports, and as high as 81–90% in others.^{3,8}

Although some authors suggest MRI can be considered due to the increased density of breasts during pregnancy⁵, MRI with contrast potentially poses a risk to the developing fetus⁶, and MRI without contrast is not helpful for breast imaging. Gadolinium should be avoided because the effect of free gadolinium ions in amniotic fluid is unknown and its use has been associated with fetal malformations in rats.^{5,6} Moreover, the lactational changes in breasts during the peri- and postpartum periods result in rapid enhancement after administration of IV contrast, making lactational changes and malignancy difficult to differentiate.^{9,10} Thus, the clinical utility of MRI in PABC is questionable. In Europe, novel MRI contrast agents gadobenate dimeglumine and gadoterate meglumine are currently both approved for use in pregnant women⁶; further study regarding their efficacy/safety is necessary.

Treatment of breast cancer in pregnancy is similar to that outside of pregnancy, with modifications for fetal indications. There is no evidence therapeutic abortion improves maternal survival, and such management is only indicated when progressive development of malignant disease is expected or fetal harm due to continued intensive adjuvant therapy is likely.¹¹

Surgical Treatment

Mastectomy versus breast-conservation surgery—Mastectomy is generally recommended in the first and second trimesters.^{12,13} Breast and axillary surgery can be

performed during any trimester of pregnancy with minimal risk to the fetus, especially after 12 weeks when the risk of spontaneous miscarriage is minimal.^{2,6,13} However, there is increased risk of miscarriage associated with surgery in the first trimester of pregnancy; hence elective surgery should be deferred to the second or third trimester if possible.¹⁴ Lumpectomy can be performed in the third trimester with delay of radiation therapy (RT) to the postpartum period. Neoadjuvant chemotherapy or lumpectomy followed by adjuvant chemotherapy can be administered during pregnancy for oncologic indications, or patient choice, if remote from delivery. Lumpectomy alone in the first trimester with delay of radiation to the postpartum period may have a detrimental oncologic impact if chemotherapy is not planned.¹³ Overall, breast conservation versus mastectomy for stage I and II PABC has been shown to have similar survival in PABC.² Radiation exposure during pregnancy is most likely to be detrimental before 10 weeks gestation (pre-implantation) or during organogenesis (until the end of the eighth week).¹² Nakagawa¹⁵ reported no significant risk to the fetus for radiation if fetal exposure does not exceed 100 mGy.

Several studies have aimed at estimating the fetal dose during tangential breast irradiation, using adult anthropomorphic phantom models consisting of transverse sections to which rings could be added to simulate the changing geometry of the pregnant female body. Typically, dosing for radiation exposure in these studies is based upon data from humans, which have estimated the tangential breast irradiation field dimensions to which patients are exposed during breast cancer treatment.¹⁶ Using such an anthropomorphic phantom model, studies were able to estimate the conceptus dose resulting from various field dose sizes during the three pregnancy trimesters. They report that for women receiving tangential breast irradiation, with a treatment dose of 50,000 mGy, the dose to the fetus was 21–76mGy in the first trimester, a time period during which doses above 100mGy are associated with severe mental retardation or fetal malformations. In the second trimester, the estimated dose was 22–246mGy, during which doses above 10mGy may be associated with decreased IQ, and those above 500–600mGy associated with mental retardation, microcephaly, and growth retardation. In the third trimester, fetal dose ranged from 22–586mGy, a time when fetal growth restriction may occur above 500mGy, implying a possible detrimental effect on IQ in the second trimester, and fetal growth in the third trimester.^{16,17} Similarly, Martin-Rincón¹⁸, administering a 50,000mGy dose to the isocenter of their field target to the field parameters for over 300 patients, estimated a dose of 38.1+/-1.3mGy with wedges and 39.2+/-2.2mGy for open field.¹⁸ Overall, therefore, it appears the majority of radiotherapeutic regimens do not affect fetal IQ factors, but may affect fetal growth.

To date, although these studies seem to indicate the dose to the developing fetus is low, no reports on actually administering RT to the breast during pregnancy have been reported. Of note, the literature includes a case report of radiation given to a woman before pregnancy was detected. Pregnancy was diagnosed at three weeks gestational age, during week two of radiation therapy, and treatment was continued until six gestational weeks. Fetal dose estimations were made at 39mGy using phantom models, but no fetal outcomes were reported.¹⁹ Thus, whether the risk of radiation during pregnancy is worth the possible risk currently remains unknown. It is known that abdominal shielding may further decrease fetal dose by 50–75%. Some current reports seem to suggest radiation therapy should not be

considered contraindicated in pregnancy¹⁷, although some have argued that it may only be safe in the first and second trimesters.²⁰

Sentinel lymph node biopsy (SLNB)—The concerns of SLNB in pregnancy are the fetal effects of exposure to radiation and/or blue dye. Radiation dose estimates of technetium sulfur colloid are significantly less than the National Council on Radiation Protection and Measurement limits for a pregnant woman. Lymphazurin blue is generally avoided because of the risk of allergic reactions and anaphylaxis, while methylene blue is avoided because it is associated with jejunal atresia during the first trimester.²⁰ SLNB is the standard of care for axillary staging of patients with clinically node-negative breast cancer. It decreases operative time and risk of complications, including lymphedema. Thus, it would be of benefit to offer SLNB in PABC.²¹

Some data exist regarding radiation-related risks to the fetus with SLNB in PABC. Ellner²² reported that the abdomen absorbs less than 1.3×10^{-3} m Gy/MBq of radiation during the use of technetium radiotracers for lymph node mapping, with a total average dose of 39MBq. This is less than the average daily background radiation of 8.2mGy per day.²³ In accordance with this finding, a recent prospective trial on 12 patients demonstrated no adverse effects to the fetus associated with low-dose lymphoscintigraphy with 99m-Tc.¹³

With respect to the use of blue dye, Pruthi²⁴ estimated that the extrapolated estimated fetal exposure to blue dye, based upon the pharmacokinetics of methylene blue in 10 non-pregnant women and the organ distribution of methylene blue, was 0.25mg. This value is likely further reduced by physiologic changes in pregnancy. They concluded that there is likely minimal fetal risk associated with use of this technique.²⁴ Additionally, Gropper²¹ described a 47-patient cohort with node-negative breast cancer in pregnancy, of whom 16 had SLNB with 99-Tc sulfur colloid and seven with isosulfan blue dye alone, with 24 of 25 pregnancies resulting in healthy deliveries. The authors concluded SLNB is feasible and appears safe in PABC. However, blue dyes in general (lymphazurin and isosulfan blue) are generally avoided because of the risk of anaphylaxis.

Reconstruction—Reconstruction following mastectomy during PABC is usually delayed until after delivery because achieving symmetry is considered difficult due to pregnancy-associated breast engorgement², as well as to fetal and maternal concerns. Recently, Lohsirawat²⁵ described reconstruction in pregnancy in 78 patients: 22 underwent unilateral mastectomy; 13 of 22 had immediate reconstruction (12 with a tissue expander and one with an immediate implant). There was no infection, hematoma, capsular contraction, or flap necrosis, and 75% of patients completed expansion inpartum. Eleven of 12 patients continued their pregnancy; one had a termination at nine weeks. With median follow-up of 32 months postpartum, one patient had expander leakage after external radiation and one had a local recurrence 19 months postmastectomy. This study suggests inpartum reconstruction is a feasible option warranting further investigation. Notably, the authors reported some difficulty related to increased breast size and an unpredictable degree of breast stiffness during pregnancy, indicating the need for appropriate patient counseling and provider understanding of breast cancers in pregnancy.

Chemotherapy and Neonatal Outcomes

Generally, chemotherapeutic treatment of PABC should be the same as those prescribed for non-pregnant patients.⁶ Chemotherapy administered during the first trimester—specifically, during organogenesis (weeks 4–12)—poses the highest risk of fetal teratogenesis, with an increased risk associated with multi-agent therapy^{26,27} and thus, ideally, it would be administered after the first trimester. Several studies have confirmed the safety of the standard 5-fluorouracil, doxorubicin, and cyclophosphamide regimen.⁶ Notably, tamoxifen is associated with a 20% birth defect risk and contraindicated in pregnancy.¹³

Several studies report on the maternal oncogenic and fetal outcomes of various combinations of anthracyclines and taxanes given to treat PABC. Hahn²⁸ reports specifically on the use of the standard regimen of 5-fluorouracil, doxorubicin, and cyclophosphamide given to a 57-woman cohort in the second or third trimesters. A median number of four cycles (range 1–6) was given at an average gestational age of 23 weeks (range 11–34 weeks). After median duration of 38.5 months, the majority of patients were disease free and delivered before 34 weeks with no stillbirths or perinatal deaths. A parent survey of children born to these women, aged 2–157 months, reported birth complications similar to population norms. The most common birth complication was difficulty breathing, reported in 10%. Among 18 school-aged children, only two required special attention in school, one with a diagnosis of attention deficit disorder. Additionally, follow-up data on 104 women with primary or recurrent breast cancer during pregnancy, treated with either adriamycin and cytoxan or one of 11 taxanes, reported recurrence in 30 and death in 21.²⁹ The fetal malformations rate was no greater than the general population, and the mean gestational age at delivery was 36 weeks, with some iatrogenic preterm birth cases following induction of labor to allow for initiation of postpartum chemotherapy treatment. Interestingly, patients reported more nausea and paresthesias as a result of the same chemotherapeutic agent given during versus after pregnancy (for patients who had intra- and postpartum treatment).²⁹

Ring³⁰ reported on 28 women who received various neoadjuvant, adjuvant, or palliative chemotherapy regimens (e.g., doxorubicin and cyclophosphamide, epirubicin and cyclophosphamide, or cyclophosphamide, methotrexate, and fluorouracil) during the second or third trimesters. Seventeen patients received adjuvant chemotherapy, totaling 116 cycles for the cohort. Patients received a median of six cycles (range 4–8) of chemotherapy during and after pregnancy. The average gestational age at the start of chemotherapy was 20 weeks (range 15–33). The most significant complication reported was febrile neutropenia (three cases). Infants were delivered at a median gestational age of 37 weeks; only one patient had spontaneous preterm labor and no fetal abnormalities were reported. After follow-up of 7–159 months, 63% patients with stage I–IIIB cancer had disease-free survival.

With respect to taxol use in pregnancy, Cardonick³¹ reported on a 15-woman cohort—12 with breast cancer—prospectively followed after intrapartum exposure to either paclitaxel or docetaxel. Median gestational age at delivery was 36.9 weeks, with two patients induced for preeclampsia. Complications at time of delivery included apnea or prematurity, GERD, neutropenia, hyperbilirubinemia, and respiratory distress syndrome. One infant had hypertrophic pyloric stenosis. Follow-up for a median 46 months showed appropriate infant growth.

One modification of chemotherapeutic treatment, which is appealing in the case of pregnancy, is that of dose-dense regimens, allowing for a shorter time period of treatment. Cardonick³¹ described 10 women exposed to dose-dense doxorubicin and cyclophosphamide. A dose-dense schedule allows for completion farther from time of delivery, requiring only 12 weeks total. Pegfilgrastim (Neulasta) or filgrastim (granix) was used in 6 of 10 women in the dose-dense group as compared with 16 of 99 women in the conventional chemotherapy group. In this study, there were no statistically significant differences in gestational age at delivery, rates of preterm labor, intrauterine growth restriction, congenital anomalies, or neutropenia, indicating the safety of this protocol.³¹

A recent meta-analysis of 18 cases using trastuzumab recommends against its use in pregnancy, as increased risk of renal and pulmonary complications were noted in newborn infants.¹³ In this report, trastuzumab was administered in pregnancy for metastatic (55.6%) or adjuvant chemotherapy, predominately in the second or third trimester (83%). Seventy-three percent of women exposed in the second or third trimester had apparently reversible oligo- or an-hydramnios, as compared with none treated during the first trimester. One-quarter of patients died. Ten of 19 infants were healthy at birth and at nine months follow-up; most had mild to severe pulmonary and/or renal disease, and/or infectious outcomes at birth. These authors concluded that trastuzumab should not be given during pregnancy, particularly because it is just as effective six months after receiving chemotherapy intrapartum.¹³

CONCLUSIONS

In summary, the treatment of breast cancer in pregnancy mirrors that outside of pregnancy, with a few important differences dictated by the balance of maternal versus fetal health, and oncologic versus obstetric outcome. In part, treatment recommendations are trimester-dependent. Breast-conserving surgery or mastectomy can be considered (with SLNB appearing feasible and safe) after first treatment. Overall, surgical treatment, possibly with neoadjuvant and/or adjuvant chemotherapy, is often feasible. In the first trimester, the risks of possible treatment delays must be weighed against the risks to the fetus secondary to oncologic treatment. Intrapartum chemotherapy is an option to optimize oncologic outcomes if continuation of pregnancy is desired. Several authors have reported good oncologic and fetal outcomes using standard breast chemotherapeutic regimens given after the first trimester of pregnancy. Tamoxifen and trastuzumab should be avoided, and dose-dense regimens with granulocyte stimulating factors may be considered.

Additionally, there is a need for further research to determine the safety of diagnostic and therapeutic procedures that are routinely used in the non-pregnant woman as well as long-term data on the oncologic safety of these approaches. Furthermore, although existing studies on surgical and chemotherapeutic treatment of these malignancies in pregnancy report overall good fetal outcomes, long-term studies on children treated with these agents in utero are needed. In order to ensure the most timely and safe treatment to women with pregnancy-associated breast cancer, improved education of providers regarding the safety of various surgical and chemotherapeutic treatments in pregnancy is important. This, in

conjunction with a multidisciplinary approach, will provide patients with the best options for optimizing oncologic as well as fetal outcomes.

Acknowledgments

FUNDING

Dr. Mary L. Gemignani was funded in part by NIH/NCI Cancer Center Support Grant No. P30 CA008748.

References

1. Meisel JL, Economy KE, Calvillo KZ, Schapira L, Tung NM, Gelber S, et al. Contemporary multidisciplinary treatment of pregnancy-associated breast cancer. *Springerplus*. 2013; 2(1):297. [PubMed: 23888269]
2. Keleher AJ, Theriault RL, Gwyn KM, Hunt KK, Stelling CB, Singletary SE, et al. Multidisciplinary management of breast cancer concurrent with pregnancy. *J Am Coll Surg*. 2002; 194(1):54–64. [PubMed: 11800340]
3. Asgeirsson KS. Pregnancy-associated breast cancer. *Acta Obstet Gynecol Scand*. 2011; 90(2):158–66. [PubMed: 21241261]
4. Basaran D, Turgal M, Beksac K, Ozyuncu O, Aran O, Beksac MS. Pregnancy-associated breast cancer: clinicopathological characteristics of 20 cases with a focus on identifiable causes of diagnostic delay. *Breast Care (Basel)*. 2014; 9(5):355–9. [PubMed: 25759617]
5. Rovera F, Frattini F, Coglitore A, Marelli M, Rausei S, Dionigi G, et al. Breast cancer in pregnancy. *Breast J*. 2010; 16(Suppl 1):S22–5. [PubMed: 21050304]
6. Amant F, Van Calsteren K, Halaska MJ, Gziri MM, Hui W, Lagae L, et al. Long-term cognitive and cardiac outcomes after prenatal exposure to chemotherapy in children aged 18 months or older: an observational study. *Lancet Oncol*. 2012; 13(3):256–64. [PubMed: 22326925]
7. Vashi R, Hooley R, Butler R, Geisel J, Philpotts L. Breast imaging of the pregnant and lactating patient: imaging modalities and pregnancy-associated breast cancer. *AJR Am J Roentgenol*. 2013; 200(2):321–8. [PubMed: 23345353]
8. Langer A, Mohallem M, Stevens D, Rouzier R, Lerebours F, Cherel P. A single-institution study of 117 pregnancy-associated breast cancers (PABC): Presentation, imaging, clinicopathological data and outcome. *Diagn Interv Imaging*. 2014; 95(4):435–41. [PubMed: 24485752]
9. Talele AC, Slanetz PJ, Edmister WB, Yeh ED, Kopans DB. The lactating breast: MRI findings and literature review. *Breast J*. 2003; 9(3):237–40. [PubMed: 12752635]
10. Ayyappan AP, Kulkarni S, Crystal P. Pregnancy-associated breast cancer: spectrum of imaging appearances. *Br J Radiol*. 2010; 83(990):529–34. [PubMed: 20335428]
11. Amant F, Brepoels L, Halaska MJ, Gziri MM, Calsteren KV. Gynaecologic cancer complicating pregnancy: an overview. *Best Pract Res Clin Obstet Gynaecol*. 2010; 24(1):61–79. [PubMed: 19740709]
12. Fajdic J, Gotovac N, Hrgovic Z, Fassbender WJ. Diagnosis and therapy of gestational breast cancer: a review. *Adv Med Sci*. 2008; 53(2):167–71. [PubMed: 18952540]
13. Zagouri F, Sergentanis TN, Chrysikos D, Papadimitriou CA, Dimopoulos MA, Bartsch R. Trastuzumab administration during pregnancy: a systematic review and meta-analysis. *Breast Cancer Res Treat*. 2013; 137(2):349–57. [PubMed: 23242615]
14. Grimm D, Woelber L, Trillsch F, Keller-v Amsberg G, Mahner S. Clinical management of epithelial ovarian cancer during pregnancy. *Eur J Cancer*. 2014; 50(5):963–71. [PubMed: 24462638]
15. Nakagawa K, Aoki Y, Kusama T, Ban N, Nakagawa S, Sasaki Y. Radiotherapy during pregnancy: effects on fetuses and neonates. *Clin Ther*. 1997; 19(4):770–7. [PubMed: 9377620]
16. Mazonakis M, Varveris H, Damilakis J, Theoharopoulos N, Gourtsoyiannis N. Radiation dose to conceptus resulting from tangential breast irradiation. *Int J Radiat Oncol Biol Phys*. 2003; 55(2):386–91. [PubMed: 12527052]

17. Kal HB, Struikmans H. Radiotherapy during pregnancy: fact and fiction. *Lancet Oncol.* 2005; 6(5): 328–33. [PubMed: 15863381]
18. Martin Rincon C, Jerez Sainz I, Modolell Farre I, Espana Lopez ML, Lopez Franco P, Muniz JL, et al. Evaluation of the peripheral dose to uterus in breast carcinoma radiotherapy. *Radiat Prot Dosimetry.* 2002; 101(1–4):469–71. [PubMed: 12382793]
19. Antypas C, Sandilos P, Kouvaris J, Balafouta E, Karinou E, Kollaros N, et al. Fetal dose evaluation during breast cancer radiotherapy. *Int J Radiat Oncol Biol Phys.* 1998; 40(4):995–9. [PubMed: 9531386]
20. Toesca A, Gentilini O, Peccatori F, Azim HA Jr, Amant F. Locoregional treatment of breast cancer during pregnancy. *Gynecol Surg.* 2014; 11(4):279–84. [PubMed: 25419205]
21. Gropper AB, Calvillo KZ, Dominici L, Troyan S, Rhei E, Economy KE, et al. Sentinel lymph node biopsy in pregnant women with breast cancer. *Ann Surg Oncol.* 2014; 21(8):2506–11. [PubMed: 24756813]
22. Ellner SJ, Hoh CK, Vera DR, Darrah DD, Schulteis G, Wallace AM. Dose-dependent biodistribution of [(99m)Tc]DTPA-mannosyl-dextran for breast cancer sentinel lymph node mapping. *Nucl Med Biol.* 2003; 30(8):805–10. [PubMed: 14698783]
23. Spanheimer PM, Graham MM, Sugg SL, Scott-Conner CE, Weigel RJ. Measurement of uterine radiation exposure from lymphoscintigraphy indicates safety of sentinel lymph node biopsy during pregnancy. *Ann Surg Oncol.* 2009; 16(5):1143–7. [PubMed: 19267158]
24. Pruthi S, Haakenson C, Brost BC, Bryant K, Reid JM, Singh R, et al. Pharmacokinetics of methylene blue dye for lymphatic mapping in breast cancer-implications for use in pregnancy. *Am J Surg.* 2011; 201(1):70–5. [PubMed: 21167367]
25. Lohsiriwat V, Peccatori FA, Martella S, Azim HA Jr, Sarno MA, Galimberti V, et al. Immediate breast reconstruction with expander in pregnant breast cancer patients. *Breast.* 2013; 22(5):657–60. [PubMed: 23871328]
26. Minig L, Otano L, Diaz-Padilla I, Alvarez Gallego R, Patrono MG, Valero de Bernabe J. Therapeutic management of epithelial ovarian cancer during pregnancy. *Clin Transl Oncol.* 2013; 15(4):259–64. [PubMed: 23180347]
27. Salani R, Billingsley CC, Crafton SM. Cancer and pregnancy: an overview for obstetricians and gynecologists. *Am J Obstet Gynecol.* 2014; 211(1):7–14. [PubMed: 24316272]
28. Hahn KM, Johnson PH, Gordon N, Kuerer H, Middleton L, Ramirez M, et al. Treatment of pregnant breast cancer patients and outcomes of children exposed to chemotherapy in utero. *Cancer.* 2006; 107(6):1219–26. [PubMed: 16894524]
29. Cardonick E, Dougherty R, Grana G, Gilmandyar D, Ghaffar S, Usmani A. Breast cancer during pregnancy: maternal and fetal outcomes. *Cancer J.* 2010; 16(1):76–82. [PubMed: 20164696]
30. Ring AE, Smith IE, Jones A, Shannon C, Galani E, Ellis PA. Chemotherapy for breast cancer during pregnancy: an 18-year experience from five London teaching hospitals. *J Clin Oncol.* 2005; 23(18):4192–7. [PubMed: 15961766]
31. Cardonick E, Bhat A, Gilmandyar D, Somer R. Maternal and fetal outcomes of taxane chemotherapy in breast and ovarian cancer during pregnancy: case series and review of the literature. *Ann Oncol.* 2012; 23(12):3016–23. [PubMed: 22875836]

Synopsis

The management of breast cancer during pregnancy presents unique challenges. Here we review the treatment of breast cancer in pregnancy and recent updates regarding the safety of surgical and chemotherapeutic treatments.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript