Atopic dermatitis, cyclosporine, and nephrotoxicity

Naoko Otani · Tetsu Akimoto · Tomoyuki Yamazaki ·
Sumiko Honma · Daisuke Nagata

Received: 15 April 2014 / Accepted: 19 June 2014 / Published online: 9 July 2014
© Japanese Society of Nephrology 2014

Keywords Cyclosporine A · Infliximab · Drug-induced renal injuries · Renal biopsy · Atopic dermatitis

Dear Editor,

Cyclosporine has been covered for the treatment for atopic dermatitis (AD) by the health insurance program in Japan since October 2008 [1]. Considering the fact that AD is one of the most common skin disorders, the number of AD patients with cyclosporine nephrotoxicity may increase in the future despite the lack of embrace of cyclosporine use by a substantial percentage of the dermatology community [1–3], although the prompt recognition of the disease may not be straight forward. Here, we present our experience with such a case in a 26-year-old male Crohn's disease (CD) patient complicated with AD. He was diagnosed with CD in 2009, when he developed several perianal fistulas requiring surgical drainage and was treated with oral mesalazine (2250 mg/day). At the end of March 2010, infliximab (5 mg/kg) in alternate months was added to his therapeutic regimen due to incomplete control of the disease activity. Thereafter, his clinical remission was maintained; however, he was referred to our hospital at the beginning of August 2012 due to a 5-month history of a moderate decline in his renal function, with a serum creatinine (sCr) level ranging from 1.5 to 2.0 mg/dL. One

month before this referral, the administration of infliximab was discontinued. At the initial interview, he reported that he had also been diagnosed with AD in his adolescence, and had been treated by a dermatologist with 100 mg cyclosporine orally twice a day (3.6 mg/kg/day) for 22 months and sporadic oral fexofenadine hydrochloride. The laboratory analysis at the referral revealed a blood urea nitrogen level of 21.5 mg/dL; sCr of 1.4 mg/dL and potassium of 5.4 mmol/L. His urine was negative for protein, and no abnormalities were noted in the sediment. Both kidneys were of normal size and renal cortex echogenicity. There were marginal changes in the sCr level during the outpatient visit despite the infliximab discontinuation, and thus he was subjected to a histological analysis, which revealed arteriolar hyalinization with interstitial fibrosis (Fig. 1), which are characteristic findings of calcineurin inhibitor nephrotoxicity [4]. Finally, we instructed his dermatologist to stop the oral cyclosporine treatment, when the patient's 1 h post-preprandial dose cyclosporine concentration (C1) was 1000 ng/mL. Three months later, his sCr was finally settled around 1.0 mg/dL despite resuming the infliximab administration.

One may argue that this kind of diagnostic delay is simply the result of poor skills in the examination and history taking. Nevertheless, the known linkage between renal complications and the therapeutic agents used for inflammatory bowel disease, such as aminosalicylates and infliximab [5] initially encouraged not only the previous doctor, but also our staff, to attribute the illness of this patient to infliximab. Although cyclosporine-related renal injury is a well-described clinical entity [4], we should bear in mind that insufficient medical information sharing and the complex nature of comorbidities, which necessitate the use of various therapeutic agents with nephrotoxic potential prescribed by more than one medical specialist as

N. Otani · T. Akimoto · T. Yamazaki · S. Honma
Department of Internal Medicine, Koga Red Cross Hospital,
Koga, Japan

N. Otani · T. Akimoto (✉) · T. Yamazaki · D. Nagata
Division of Nephrology, Department of Internal Medicine,
Jichi Medical University, 3311-1 Yakushiji, Shimotsuke-Shi,
Shimotsuke, Tochigi 329-0498, Japan
e-mail: tetsu-a@jichi.ac.jp

Fig. 1 A renal biopsy was performed at the end of December 2012. Light micrographs showed a glomerulus with thickening of Bowman's capsule, which was associated with an arteriole demonstrating nodular hyaline deposits in the medial smooth muscle layers. *Inset* similar deposits were noted in some arterioles (**a** periodic acid–Schiff stain). The patchy and/or striped form of interstitial fibrosis with adjacent tubular atrophy was also noted (**b** Masson's trichrome stain). The *scale bar* is indicated in each panel

Conflict of interest None declared.

References

1. Saeki H, Furue M, Furukawa F, Hide M, Ohtsuki M, Katayama I, Sasaki R, Suto H, Takehara K, Committee for Guidelines for the Management of Atopic Dermatitis of Japanese Dermatological Association. Guidelines for management of atopic dermatitis. *J Dermatol.* 2009;36:563–77.
2. Furue M, Yamazaki S, Jimbow K, Tsuchida T, Amagai M, Tanaka T, Matsunaga K, Muto M, Morita E, Akiyama M, Soma Y, Terui T, Manabe M. Prevalence of dermatological disorders in Japan: a nationwide, cross-sectional, seasonal, multicenter, hospital-based study. *J Dermatol.* 2011;38:310–20.
3. Ryan C, Amor KT, Menter A. The use of cyclosporine in dermatology: part II. *J Am Acad Dermatol.* 2010;63:949–72.
4. Colvin RB, Nickleleit V. Renal transplant pathology. In: Jennete JC, Olson JL, Schwartz MM, Silvia FG, editors. *Heptinstall's pathology of the kidney*. 6th ed. Philadelphia: Lippincott Williams & Wilkins; 2007. p. 1346–490.
5. Oikonomou K, Kapsoritakis A, Eleftheriadis T, Stefanidis I, Potamianos S. Renal manifestations and complications of inflammatory bowel disease. *Inflamm Bowel Dis.* 2011;17:1034–45.

described in the current report, may preclude the prompt identification of the disease.