CORRESPONDENCE

Unnecessary Investigations in Environmental Medicine—A Retrospective Cohort Study

by Annette Greiner, and Hans Drexler
in issue 46/2016

Constructive Dialogue Necessary

The diagnostic strategies of clinical environmental medicine are outlined in the guidelines of the German Professional Association of Environmental Medicine (dbu; *Deutscher Berufsverbands der Umweltmediziner*) (1). These guidelines are based on the curriculum that was developed at that time together with the German Medical Association to obtain the additional title for this area of expertise. Although the additional titles have since been abolished, curricular training continues to follow these criteria. Indeed, a large part of the faulty diagnostics is not recommended, neither in the courses nor in the guideline.

It is not clear what inclusion criteria were used for this retrospective study, based on this article. The authors themselves state that the study included patients from occupational, environmental, and social medicine, so that the study can not be considered representative of patients of clinical environmental medicine.

The authors of a European Union (EU) study (2) concluded that 7% of all diseases are triggered by only nine environmental stressors in the six core EU countries. This corresponds to the extent of important national diseases.

We urgently need a constructive dialogue in this area if we want to reach solutions. This article follows the opposite strategy. It is particularly encouraging, for example, that the Robert Koch Institute hosts moderated discussions with colleagues from other disciplines, which are characterized by a factual and respectful approach. This allows doors to be opened rather than slammed shut.

DOI: 10.3238/arztebl.2017.0252a

REFERENCES

1. Bartram F, Bauer A, von Baehr V, et al.: Handlungsorientierte umweltmedizinische Praxisleitlinie – Leitlinienreport. Berlin: Deutscher Berufsverband der Umweltmediziner e. V. (eds.) 2010.
2. Hänninen O, Knol A: European perspectives on environmental burden of disease. Estimates for nine stressors in six European countries. Helsinki, Finland: University Printing 2011.
3. Greiner A, Drexler H: Unnecessary investigations in environmental medicine—a retrospective cohort study. *Dtsch Arztebl Int* 2016; 113: 773–80.

Dr. med. Kurt E. Müller

Mitglied beim Deutschen Berufsverband der Umweltmediziner e. V., Berlin, Germany
kurt.mueller@preventamed.de

Environmental Diseases Are Multifactorial

Clinical environmental medicine is therapy for patients with complex environmental-related multifactorial (chronic) diseases. Thus, diagnostics require more than just acute intoxications (blood and urine investigations) or dermal sensibilities in order to be successful.

In a study that aims to challenge unnecessary diagnostic tests (1), it is first necessary to explain why the respective diagnostic tests should be considered unnecessary. Likewise, no distinction is made between acute and chronic stress. Furthermore, with respect to the lymphocyte transformation test, the authors confuse immunology with toxicology. They also ignore the fact that, for toxic stresses and especially for immunological stresses, the reference values are less important than the individual vulnerability, which comprises specific susceptibilities to the detoxification mechanisms as well as the personal immune mechanisms that are genetically determined.

Scientifically supported diagnostic tests, such as lymphocyte transformation tests, blood or urine monitoring, and DMPS/DMSA mobilization (DMPS, dimercaptopropane sulfonic acid; DMSA, dimercaptosuccinic acid), are falsely thrown together with purely empirical or even esoteric methods (such as pendulum divination). In addition, false examination tests for chronic stress, such as simple blood and urine samples without the use of a chelating agent for mobilization, are listed. As the necessary substitution possibilities are not mentioned, these do not meet scientific criteria. The classical conventional medical method of hair analysis, which is known from the field of forensics for determining long-term exposure, is unambiguously described as an unsuitable test; the same applies to the saliva test.

Anyone who ultimately demands an improvement of symptoms once a known exposure is reduced is still bound to the linear cause-effect principle of Isaac Newton. However, complex scientific entities such as environmental-related multifactorial diseases are currently (in the 21st century) being cybernetically defined.

DOI: 10.3238/arztebl.2017.0252b

REFERENCES

1. Greiner A, Drexler H: Unnecessary investigations in environmental medicine—a retrospective cohort study. *Dtsch Arztebl Int* 2016; 113: 773–80.

Dr. medic-stom/RU Martin F. H. K. Klehmet

Praxis für Umweltzahnmedizin - Stomatologie, Bremen, Germany
123klehmet456@web.de

Diagnostic Use of Chelating Agents

The assessment of the DMPS/DMSA provocation test (DMPS, dimercaptopropane sulfonic acid; DMSA, dimercaptosuccinic acid) is not very balanced in the article by Greiner and Drexler (1). For example, the German Environment Agency (*Umweltbundesamt*, UBA) assumes that determining the chronic aluminium load is only possible with the use of a complexing agent (chelator). Simple urine and blood samples are unsuitable for the determination of chronic aluminium loads, since only the momentary exposure can be detected (2). However, Prof. Meissner from Dresden notes that chronic poisoning with metals is always the result of accumulation (3). Accumulation occurs over a longer period, by intake even of small quantities.

Mobilization tests with chelating agents lead to a release of metals from their depots and their increased excretion in the urine. These tests are important when the urine and blood samples do not provide clear results (3).

The so-called triple test was developed and standardized by the state medical laboratory in Luxembourg in 1995, in order to counteract the diagnostic uncertainty arising from simple blood and urine measurements and to obtain reliable information about the mercury exposure of individual patients. This procedure includes the determination of mercury in urine before and after DMPS administration (4).

These few examples show that there are other voices supporting the diagnostic use of DMPS and other chelating agents, which unfortunately were not included in the article by Greiner and Drexler.

DOI: 10.3238/arztebl.2017.0252c

REFERENCES

1. Greiner A, Drexler H: Unnecessary investigations in environmental medicine—a retrospective cohort study. *Dtsch Arztebl Int* 2016; 113: 773–80.
2. Umweltbundesamt: Stellungnahme der Kommission „Human-Biomonitoring“ des Umweltbundesamtes: Aluminium. *Bundesgesundheitsblatt – Gesundheitsforschung – Gesundheitsschutz* 1998; 41: 271.
3. Meißner D: Problematik, Klinik und Beispiele der Spurenelementvergiftung – Eine Einführung. *Toxichem Krimtech* 2011; 78: 447–52.
4. Hansen G, Victor R, Engeldinger E, Schweitzer C: Evaluation of the mercury exposure of dental amalgam patients by the Mercury Triple Test. *Occup Environ Med* 2004; 61: 535–40.

Peter Jennrich

Praxis für Allgemeinmedizin, Naturheilverfahren, Würzburg, Germany
peter-jennrich@gmx.de

In Reply:

The inclusion criteria, which Dr. Müller requested, are described in the Methods section of our article (1). In the prospective study, 653 patients were included consecutively, without exclusion criteria in order to prevent selection effects. Relating to the guidelines of the German Professional Association of Environmental Medicine (dbu), we would like to point out that the term “guideline” is not protected. Beside the scientifically accepted guidelines of the Association of Scientific Medical Societies in Germany (AWMF, *Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften e.V.*), there are guidelines in varying quality published by other societies. The diseases described in the study by Hänninen (2) are not specifically environmental diseases and can therefore be diagnosed and treated according to AWMF guidelines. We attempted to present our results objectively and without polemics, since a respectful relationship with colleagues is very important to us.

Dr. Klehmet first points out that a distinction must be made between acute and chronic exposure. However, this differentiation is only possible by using the history and the kinetics of the foreign substances or their metabolites, since quantification of burdens cannot distinguish between acute and chronic effects. It is correct that the lymphocyte transformation test is an immuno-

logical procedure. Nonetheless, only the type IV allergy from the Gell and Coombs classification can be detected, but not general immunological disturbances. The hypothesis of a clinical significance of genetic polymorphisms (susceptibilities) for detoxification has been controversially discussed in environmental medicine for 25 years. As long as there is no scientific evidence to show that these are also relevant in clinical environmental medicine, using genetic polymorphism determination for individual diagnosis based on an environmental medical issue cannot be recommended (3, 4).

Peter Jennrich comments in particular on the importance of the use of chelating agents. Without a doubt, chelating agents lead to increased excretion of metals in the urine. In particular, the first urine fraction obtained after administration of the chelating agent contains very high concentrations of metals. Thus, extrapolating of the concentration in the first urine fraction to one liter urine or even to a 24-hour excretion is therefore not possible. The metals are primarily mobilized from the blood and kidneys. Since the mobilization values are closely related to the respective initial values of the spontaneous Hg excretion, the test procedure does not yield any significant additional findings on Hg loading or storage (5). Thus, considering their potential side effects, using a chelating agent with of non-recognizable diagnostic value cannot be justified. Furthermore, there are no generally accepted reference values, so the metal concentrations can hardly be assessed after mobilization. The cited statement of the Umweltbundesamt (UBA) in this regard relates to the diagnosis of aluminum intoxication in renal insufficiency patients, but not to environmental medical questions, and thus is not related to our article.

DOI: 10.3238/arztebl.2017.0253

REFERENCES

1. Greiner A, Drexler H: Unnecessary investigations in environmental medicine—a retrospective cohort study. *Dtsch Arztebl Int* 2016; 113: 773–80.
2. Hänninen O, Knol A: European perspectives on environmental burden of disease. Estimates for nine stressors in six European countries. Helsinki, Finland: University Printing 2011.
3. Schulz TG, Hallier E: Die Bedeutung von genetischen Polymorphismen Fremdstoff-metabolisierender Enzyme in der Arbeitsmedizin. *Arbeitsmed Sozialmed Umweltmed* 1999; 34: 307–14.
4. RKI-Kommission „Methoden und Qualitätssicherung in der Umweltmedizin“: Genetische Polymorphismen (Sequenzvariationen) von Fremdstoff-metabolisierenden Enzymen und ihre Bedeutung in der Umweltmedizin. *Bundesgesundheitsblatt – Gesundheitsforschung – Gesundheitsschutz* 2004; 47: 1115–1123.
5. Schiele R: Amalgam fillings—tolerance. *Dtsch Zahnärztl Z* 1991; 46: 515–8.

Dr. med. Annette Greiner, Prof. Dr. med. Drexler

Institut und Poliklinik für Arbeits-, Sozial- und Umweltmedizin
Universität Erlangen-Nürnberg, Germany
annette.greiner@fau.de

Conflict of interest statement

All authors of all contributions declare that no conflict of interest exists.