CORRESPONDENCE

Antiviral Medications in Seasonal and Pandemic Influenza—A Systematic Review

by Regine Lehnert, Mathias Pletz, Annicka Reuss, and Tom Schaberg
in issue 47/2016

Unnecessary and Expensive Therapy

For several years, it has been known that the pharmaceutical company Roche had held back a large amount of study data on oseltamivir (Tamiflu). Further, after publication of these data, significant corrections were required for both the effectiveness as well as the number needed to treat (NNT). With its currently acknowledged shortening of disease duration by 0.5 days, Tamiflu has achieved the efficacy of Echinacea herbal products—which, according to studies, also shorten disease duration of patients with mild to moderate influenza by 0.5 days. Likewise, antibiotics reduce moderate bronchitis duration by 0.5 days and are therefore not indicated. Unfortunately, the Robert Koch Institute (RKI) and the Institute for Quality and Efficiency in Health Care (IQWiG) have yet to issue new, sensible statements, which has led to the fatal, expensive, and nonsensical practice of using neuraminidase inhibitors. Especially in times when politicians as well as health insurance companies propagate various prevention strategies, while patients with osteoporosis receive prophylactic therapy paid by a health insurance only after the occurrence of a pathological fracture, an unnecessary and expensive therapy should not be recommended to the detriment of the statutory health insurance funds. It is also peculiar that this article (1) appeared just at the beginning of the flu season. I trust that neither the authors of the review nor the various colleagues that they acknowledge had conflicts of interest.

DOI: 10.3238/arztebl.2017.0269a

REFERENCE

1. Lehnert R, Pletz M, Reuss A, Schaberg T: Antiviral medications in seasonal and pandemic influenza—a systematic review. *Dtsch Arztebl Int* 2016; 113: 799–807.

Michael Topp

Praxis für Allgemeinmedizin, Halstenbek, Germany
topp.michael@gmx.de

Open Questions

The article by Lehnert et al. (1) has left me, as a primary care provider for the treatment, quite perplexed:

Despite the excessive mortality due to influenza viruses, of 20 000/year in Germany (www.rki.de/SharedDocs/FAQ/Influenza/FAQ_Liste.html), it has still somehow not been possible to carry out studies with the required number of participants.

Thus, I am supposed to use neuraminidase inhibitors for patients for whom efficacy has not been demonstrated (for instance, pregnant women and immunosuppressed patients).

In fact, eTable 2 even shows that there is no reduction of disease duration for elderly patients, yet I am expected to still treat them.

The risk–benefit ratio of the antiviral therapy is believed to be good, based on a reduction of bronchitis and “patient-perceived pneumonia” in a patient group that I do not treat (healthy adults).

Finally, even though the studies show that neuraminidase inhibitors hardly are effective, I should still use them because there are no other therapeutic alternatives. This contradicts the requirements of our health insurance system for diseases where no therapeutic alternatives exist. Further, the study by Muthuri et al. (2), which states that mortality decreases when using neuraminidase inhibitors, was strongly criticized for its statistical methods (3, 4).

From 2005–2009, approximately 330 million euros of public funds were spent only in Germany on neuraminidase inhibitors, which were first stored and then destroyed.

Faced with these facts, what advice should I give to my patients? Unfortunately, I remain at a loss on this subject. I would like to protect my patients, but I do not feel that I can recommend these drugs to them or to our society in general.

DOI: 10.3238/arztebl.2017.0269b

REFERENCES

1. Lehnert R, Pletz M, Reuss A, Schaberg T: Antiviral medications in seasonal and pandemic influenza—a systematic review. *Dtsch Arztebl Int* 2016; 113: 799–807.
2. Muthuri SG, Venkatesan S, Myles PR, et al.: Effectiveness of neuraminidase inhibitors in reducing mortality in patients admitted to hospital with influenza A H1N1pdm09 virus infection: a meta-analysis of individual participant data. *Lancet Respir Med* 2014; 2: 395–404.
3. Kmiotowicz Z: Study claiming Tamiflu saved lives was based on „flawed“ analysis. *BMJ* 2014; 348: g2228.
4. Jones M: Mark Jones's reply to Myles and Leonardi-Bee's response to his critique of their paper reported in *The BMJ*. *BMJ* 2014; 348: g3001.

Dr. med. Thomas Maibaum

Praxis für Allgemeinmedizin, Rostock, Germany
thomas.maibaum@yahoo.de

In Reply:

As stated in the review (1), our aim was to summarize the available data on influenza antivirals in an objective way, and not to provide our own clinical recommendations. Instead, we complemented our results with treatment recommendations from professional societies and international public health institutes.

The primary question addressed by the clinical trials for drug approval was whether disease

duration is shortened by neuraminidase inhibitors (NIs). For this purpose, the patient groups were kept as homogeneous as possible—as is commonly done, also in other drug registration studies. Thus, particularly vulnerable patients, such as pregnant women or immunosuppressed or seriously ill patients, were excluded. This leads to the concern of missing data from randomized controlled trials (RCTs) for some relevant patient populations, such as hospitalized patients, as also highlighted in the correspondences.

In the absence of active control therapies, RCTs should have been placebo-controlled, which would have been difficult to justify on ethical grounds after approval of the NIs. In addition, the acuity of influenza disease (time window), the varying severity of influenza seasons, and regional differences in influenza activity (pre-selection of study centers) can render the implementation of RCTs difficult, particularly for clinically relevant patient populations. The latter two factors are especially unpredictable.

It should be kept in mind that the RCTs were performed during seasonal influenza waves leading to

generally mild disease. Hence, for these studies, it might not have been possible to show efficacy with regard to more clinically relevant endpoints, such as reduction of influenza-related complications. Observational studies cannot replace the evidence from RCTs (the methodological problems of observational studies are also discussed in the article).

However, since data from RCTs are lacking, the results of some very large observational studies should not be overlooked. In conclusion, we would also welcome better data on influenza antivirals, especially for the known risk groups.

DOI: 10.3238/arztebl.2017.0269c

REFERENCE

1. Lehnert R, Pletz M, Reuss A, Schaberg T: Antiviral medications in seasonal and pandemic influenza—a systematic review. *Dtsch Arztebl Int* 2016; 113: 799–807.

On behalf of the authors:

Regine Lehnert

Bundesinstitut für Arzneimittel und Medizinprodukte, Bonn, Germany
Regine.Lehnert@bfarm.de

Conflict of interest statement

The authors of all contributions declare that no conflict of interest exists.

CLINICAL SNAPSHOT

Differential Diagnosis: Child Abuse

The differential diagnosis of skin discoloration in children includes child abuse as well as accidental injury, hematologic diseases, and congenital skin lesions. A seven-month-old boy from a Turkish family was brought to the hospital because of the suspicion that he had been beaten by his father. The forensic medical examination revealed poorly demarcated, homogeneous, bluish discolorations of the skin, of varying sizes, on the back and buttocks. On re-examination five days later, the appearance of the lesions was unchanged and a definitive diagnosis of Mongolian spots was made.

Mongolian spots are benign congenital pigmented lesions that are common in some ethnic groups. They are usually located on the buttocks, and they disappear by puberty. Although it is clearly important that a diagnosis of child abuse should never be missed, the formulation of a differential diagnosis is important as well, so that misinterpretations with unfortunate legal consequences can be avoided. In case of uncertainty, a follow-up examination after an adequate period of time is recommended so that the evolution in color associated with true traumatic hematomata (bruises) can be observed or ruled out.

Dr. med. Benjamin Ondruschka, Dr. med. Christian König

Institut für Rechtsmedizin, Universität Leipzig, benjamin.ondruschka@medizin.uni-leipzig.de

Conflict of interest statement: The authors state that they have no conflict of interest.

Translated from the original German by Ethan Taub, M.D.

Cite this as:

Ondruschka B, König C: Differential diagnosis: child abuse. *Dtsch Arztebl Int* 2017; 114: 270. DOI: 10.3238/arztebl.2017.0270