CORRESPONDENCE

The Differential Diagnosis of Dyspnea

by Dominik Berliner, Nils Schneider, Tobias Welte,
and Johann Bauersachs in issue 49/2016

Assure That Diagnosis Is Useful to Therapy Goal

I thank the authors for an overview of the differential diagnosis of dyspnea that is responsive to curative treatment (1). That, however, was not the subject of the article's title. The symptom of shortness of breath is a predictor of increased mortality and often manifests itself in people with multimorbidity near the end of life. For instance, cardiac causes of dyspnea is a worse prognostic factor of long-term survival than colorectal cancer (2).

Even with symptoms such as deep unconsciousness or inspiratory crackles, the obvious differential diagnosis of "dying person" is not mentioned in the article. Yet in these cases, it is a matter of the code of medical ethics to refrain from burdensome examinations. For lung infections, it is often claimed that elderly people should always receive stationary treatment—yet this is often against their will and is even contraindicated for dying persons.

Nevertheless, the article explains the various invasive diagnostic possibilities, although only the markers FEV1, NT-proBNP, and the emphysematous change are predictors. Coronary angiography, for example, is not an indication for this case, even though Germany is a world leader in it—yet we lag behind in reducing hospital mortality rates after heart attack (3). This alone points to overdiagnostics and overtreatment, which is a major problem for end-of-life care (4).

CME articles have a responsibility to promote knowledge to young colleagues as well. In a country of invasive overdiagnostics, the list of possible examination procedures does not appear to be helpful.

DOI: 10.3238/arztebl.2017.0271a

REFERENCES

1. Berliner D, Schneider N, Welte T, Bauersachs J: The differential diagnosis of dyspnoea. *Dtsch Arztebl Int* 2016; 113: 834–45.
2. Stewart S, MacIntyre K, Hole DA, Capewell S, McMurray JJV: More 'malignant' than cancer? Five-year survival following a first admission for heart failure in Scotland. *Eur J Heart Fail* 2001; 3: 315–22.
3. OECD: Health at a glance 2015: OECD indicators. Paris: OECD publishing 2015. http://dx.doi.org/10.1787/health_glance-2015-en. (last accessed on 23 March 2017).
4. Thöns M: Patient ohne Verfügung. Das Geschäft mit dem Lebensende. 7th edition. München: Piper 2016

Dr. Matthias Thöns
Palliativnetz Witten e. V., Germany
thoens@sapv.de

Don't Forget Ketoacidosis

That the article (1) focuses on the differentiation of pulmonary and cardiac causes is understandable, due to their frequency. However, it thereby neglects metabolic causes, although these are by no means rare and are of high practical relevance.

I have repeatedly seen type 1 diabetes in children and youth lead to ketoacidosis that is misdiagnosed as pulmonary dyspnea. The snag for two young patients was that both had known pulmonary problems: a small child with recurrent obstructive bronchitis, and a youth with bronchial asthma. Neither child was known to have diabetes, which manifested itself acutely with severe metabolic imbalance. During the very full office visitation hours ("infection season"), each was treated with inhalation or with cortisone after a short consultation with the attending physician (pneumologist or pediatrician). The subsequent medical examination likewise did not lead to the correct diagnosis, giving them a rocky start.

Both came to the clinic with severe acidosis in extremis (with pH values of 6.8 and 6.9!), but thankfully, both survived without complications. A diabetic ketoacidosis must be taken into consideration for dyspnea at every age, especially in light of the ever increasing incidence of type 1 diabetes mellitus. Also, the respiratory sounds should be differentiated—breathing during an asthma attack or a psychogenic hyperventilation differs markedly from the deep breathing in metabolic acidosis ("kussmaul").

In general, it is important to keep in mind that any anamnesis should be adequately addressed with open questions before attempting to differentiate into more precise specifications. It is often the case that a broader anamnesis would have given decisive indications to avoid a misdiagnosis, such as the symptoms mentioned above that are typical for diabetes.

DOI: 10.3238/arztebl.2017.0271b

REFERENCES

1. Berliner D, Schneider N, Welte T, Bauersachs J: The differential diagnosis of dyspnoea. *Dtsch Arztebl Int* 2016; 113: 834–45.

Prof. Dr. med. Walter Burger
Berlin, Germany
walterburger@gmx.de

Indications for Kidney Disease Not Adequately Discussed

The continuing education contribution (1) gives a comprehensive overview of the symptom "shortness of breath." The pulmonary and cardiac causes of dyspnea and the possibilities of differential diagnoses are presented in detail. The frequent comorbidity of both organ systems is pointed out.

Causes of dyspnea outside the respiratory and cardiovascular systems are treated only briefly: anemia, ENT, neuromuscular disorders, and psychogenic and medicinal causes. However, I find a specific mention of renal diseases, and in particular, references to acute renal failure in chronic kidney disease, to be missing.

The authors only indirectly draw attention to the importance of kidney function (in Figure 1: basic evaluation, laboratory tests, renal function tests).

Why are the main nephrogenic causes of dyspnea, which are very easy to misinterpret, especially in geriatrics, not mentioned in detail?

DOI: 10.3238/arztebl.2017.0271c

REFERENCE

1. Berliner D, Schneider N, Welte T, Bauersachs J: The differential diagnosis of dyspnoea. Dtsch Arztebl Int 2016; 113: 834–45.

Dipl.-Med. Ulrike Thäle
Dresden, Germany

In Reply:

We are grateful for the helpful and important comments. Undoubtedly, continued diagnostics should only be performed upon considering the patient. In this respect, we thank Dr. Thöns for emphasizing that, especially at the end of life, overdiagnoses can be a burden rather than a meaningful explanation of cause that leads to therapy. Assessing whether diagnostics and the corresponding therapy, including inpatient care, are meaningful is one of the primary tasks of medical practice. This ethical question must be, and should be, asked on a daily basis and precedes any diagnostic activity. Our CME article aims to transmit the various diagnostic possibilities within a framework of acute or chronic dyspnea symptoms, with the assumption that a diagnosis is either desired by the patient or appears to be medically necessary under the circumstances.

Dr. Thöns also points out that CME articles are responsible for conveying the current state of knowledge to younger colleagues. Indeed, the use of the CRB-65

score, an internationally recognized and applied tool, can also be viewed in this context. Notably, younger colleagues frequently work in emergency departments, where they have to make rapid decisions about further diagnoses and therapies. Valid scores have the task of supporting such decision-making processes and providing assistance; they do not, of course, release the physicians from medical responsibilities that should be based on ethical considerations.

The other two authors likewise highlight an important point: the causes of dyspnea symptoms are manifold and require a collective observation. Thus, chronic dyspnea in pediatric patients is not likely to be caused by chronic obstructive pulmonary disease (COPD). Unfortunately, not all differential diagnoses can be broadly discussed in the limited scope of a journal article. In order to take this point into account, we provided a comprehensive description of the causes of dyspnea in *eTable 2*. Ketoacidosis and other metabolic causes are also listed here. A nephrogenic cause of dyspnea is not one of the most common causes according to the currently available literature—but it should always be taken into account for geriatric patients. However, renal failure is often only a secondary expression of an underlying disease, such as cardiac decompensation. Unfortunately, this detailed table could not be included in the printed version of the article. Finally, due to the expertise of the authors, the article focuses on the daily routine of patients in the adult age group. A similar article on dyspnea in children and adolescents could certainly be a useful supplement.

DOI: 10.3238/arztebl.2017.0272

REFERENCE

1. Berliner D, Schneider N, Welte T, Bauersachs J: The differential diagnosis of dyspnoea. Dtsch Arztebl Int 2016; 113: 834–45.

On behalf of the authors:
Dr. med. Dominik Berliner
Prof. Dr. med. Johann Bauersachs
Klinik für Kardiologie und Angiologie
Medizinische Hochschule Hannover, Germany
berliner.dominik@mh-hannover.de

Conflict of interest statement

The authors of all contributions declare that no conflict of interest exists.