Published in final edited form as:

J Bioanal Biomed. 2015 December ; 7(6): . doi:10.4172/1948-593X.1000e133.

The Biochemical Cascades of the Human Pancreatic β -Cells: The Role of MicroRNAs

Joseph W Kim¹, John Z Luo^{1,2}, and Luguang Luo^{1,*}

¹Department of Internal Medicine, Roger Williams Hospital, Boston University School of Medicine, Providence, RI, USA

²Doctor's Choice LLC, Warwick, RI, USA

Abstract

Diabetes mellitus is a disease that poses a burden to the health care system due to its prevalence and chronic nature. Understanding β cell pathophysiology may lead to future therapeutic options for diabetes mellitus type 1 and 2. MicroRNAs (MiR) fine-tune β cell biochemical cascades through specific protein targets. This review argues that miRs may play a critical role in human islet β cell biology and are potential candidates for a new pharmacological strategy. We have reviewed and presented how miRs fine tune four biochemical cascades in islet β cells: glucose stimulated insulin secretion, β cell replication, apoptosis, and development. Only studies that examine human pancreatic islets either *in vitro* or *in vivo* are included. The unveiling role of miR pathways in regulating human islet β cell biology could open the door for diagnostic and therapeutic methods for diabetes mellitus prevention and therapy.

Keywords

miRNAs (miR); Human islet; β cells; Diabetes; Metabolism

Introduction

Diabetes mellitus is a prevalent disease in the USA and worldwide. It has a significant burden on the health care system due to its chronic nature and its long-term complications [1–4]. Novel therapeutic and diagnostic strategies could potentially lower the cost to the health care system and ultimately lower the morbidity and mortality of diabetic patients. A better understanding of the physiological mechanisms of the pancreatic β cell could help develop new therapies. In spite of their different pathophysiologies, both diabetes mellitus type 1 and type 2 could benefit from therapies that focus on the β cells. One approach to understanding β cell physiology is the microRNA (miR). MiRs were discovered in the 1990s in *Caenorhabditis elegans*, and they epigenetically inhibit mRNA translation [5–8]. By inhibiting specific protein or enzyme targets, miRs can regulate a biochemical cascade

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

*Corresponding author: LuGuang Luo, Center for Stem Cell Research, Department of Medicine/Research, Prior Building Floor 2, 825 Chalkstone Avenue, Providence, Rhode Island, 02908, USA, Tel: +1 401 456 5344; Fax: +1 401 456 5759; Lluo@rwmc.org.

within the β cell. Depending on which protein is targeted, the overall biochemical cascade can be inhibited or stimulated [6] miRs, by targeting specific enzymes or proteins, can uniquely “fine tune” the biochemical cascades of the β cells. miRs change subtle aspects of a cascade, affecting when, why, or how it occurs. Studies that examine the effect of miRs on biochemical cascades focus on four pathways: glucose stimulated insulin secretion (GSIS) [9–19], replication [20–24], apoptosis [25–32], and development [33–35].

Multiple miRs regulate each cascade and have unique ways of fine tuning them (Figure 1). This review argues that miRs could be the mechanism of a new form of regulating human islet β cell biology. Using pharmacologic agents that target miR pathways, human islet β cells may be fine-tuned. First, the article will demonstrate how β cell biochemical cascades are fine-tuned by miR; second, potential agents or special vectors carrying miRs can be used to target miR pathways modifying human islet β cell biological processes.

miRs Fine-Tune β Cell Biochemical Pathways

miRs affect subtle aspects of GSIS or the response of GSIS to a stimulus

GSIS is the special release of insulin from β cells in response to glucose [36,37]. It includes β cell glucose detection, insulin released by exocytosis after insulin translation, and insulin storage in vesicles. Multiple enzymes regulate these multiple biochemical cascades in GSIS, and miRs regulate these enzymes to fine tune the GSIS process in human islet β cell in response to glucose [38,39]. MiR-124 affects β cell responsiveness to glucose. miR-124 targets Foxa2, which is involved in glucose detection. By inhibiting Foxa2, miR-124 can change how a β cell responds to dynamic levels of serum glucose [12]. Note that the total insulin secreted will remain unchanged.

MiR-124 also affects insulin exocytosis by targeting the protein Mtpn. By doing so, miR-124 can changes the rate by which insulin is secreted. The action of miR-124 will not affect the total amount of insulin released [12]. Hence, “how” GSIS occurs is altered. Unlike the regulation of miR-124, miR-204 affects the total insulin available for release. miR-204 inhibits MafA, an insulin transcription factor. When MafA is inhibited, the total insulin available in vesicles is reduced [16]. Note that the rate of insulin secretion is not affected. The responsiveness of the β cell to serum glucose levels may not change. Some miRs do not change how GSIS occurs, but rather they affect “when” it occurred. miR-133 suppresses GSIS in the presence of glucose, forming glucotoxicity. miR-133 inhibits polypyrimidine tract binding protein expression (PTB). This inhibits insulin synthesis, decreasing insulin available for exocytosis. Another miR, miR-146a, also inhibits GSIS in response to IL-1 β . Hence, it may form the basis by which IL-1 β and inflammation inhibit insulin secretion [32].

Finally, the example of miR-29 illustrates how delicately a miR can change a biochemical cascade. In the presence of miR-29, GSIS is inhibited. Though GSIS is inhibited, increased glucose concentrations can still increase insulin secretion. miR-29 does not completely terminate insulin secretion, but rather changes its “responsiveness” to serum glucose [28,40].

miRs affect apoptosis in response to a stimulus

Apoptosis is programmed cell death (a chronic cell loss process) differing from necrosis (an acute cell loss process). Initiating apoptosis usually requires a special trigger or stimulus. Unlike GSIS, apoptosis cannot be subtly altered but rather its direct response to a stimulus is. A group of miRs initiates β cell apoptosis through β cell detection of a cytotoxic agent. These include either lipids or cytokines. miR-24 is responsible for the mechanism of lipotoxicity in β cell. miR-24 is overexpressed in response to palmitate, a lipid [29]. Then it reduces bcl-2 levels, which induces apoptosis [41,42]. In another instance, the stimuli for the miR are adjacent cells. miR-146a inhibits apoptosis in response to the presence of bone marrow stem cells. miR-146a inhibits FAS, which is crucial to the canonical FAS-mediated apoptotic pathway. This improves β cell longevity and increases β cell mass [43].

miRs regulate β cell replication as a compensatory mechanism for diabetes mellitus type 2 or insulin resistance

When β cells undergo replication, the β cell mass increases, increasing the total insulin secreted from the pancreas. miR-7 regulates replication in response to insulin resistance. miR-7 inhibits the mTOR pathway to increase β cell replication through mTORC1. miR-7 is lowered during insulin resistance, demonstrated in cadaveric studies. [10,21]. As a result, insulin secretion increases in response to insulin resistance. Another miR, miR-184, increases β cell replication in response to diabetes mellitus type 2 through argonaute2. During states of insulin resistance, miR-184 is decreased. miR-184 no longer inhibits argonaute2. Once argonaute2 levels rise, cell proliferation increases [22]. This increases total insulin secretion in the setting of increased diabetes mellitus. In an abstract by Wu, bone marrow stem cells influence adjacent human pancreatic islets via miR-24-2. miR-24-2 inhibits P16INK4a, which inhibits PDX1. The net effect is increased β cell replication. This partly explains the observation that bone marrow co-cultured with human pancreatic islets improves the longevity of the islets and miR-24-2 may play a critical role in human β cell replication [43].

miR-7 affects β Cell development through PDX-1 and ISL-1

MiR-7 is involved in development by “fine tuning” the timing of this developmental process. The number of studies looking at miRs during human pancreatic β cell development is limited and less definitive [44]. But several studies suggest the role for miR-7 during development. First, knockout studies have demonstrated that miRs are necessary for β cell development [45]. Second, the expression of miR-7 correlates with β cell development in human fetal pancreatic tissue [35]. Third, expression of miR-7 is associated with human embryonic stem cells that convert into insulin producing cells (IPCs) [34]. It is possible that miR-7 is involved in converting human embryonic stem cells into insulin producing cells (IPCs). Finally, miR-7 is associated with PDX-1 and ISL-1. PDX-1 is found in differentiated β cells. ISL-1 is a marker for differentiated endocrine cells [46]. Another study suggests that miR-7 also controls the timing of β cell differentiation. Starting week 9 until to 14~18 weeks of gestation, miR-7 is expressed at a higher level, which coincides with β cell development process [32,33]. Thus, miR-7 may play a role in fine-tuning the timing of the necessary β cell developmental biochemical cascades.

miR Pathways can be Inhibited or Stimulated

Exposing pancreatic β cells to oligonucleotides with the same sequences as a miR could activate that pathway [47]. Alternatively, oligonucleotides with the complementary sequence could bind and inhibit these miRs [48]. A number of the studies looking at the effect of a miR on a biochemical cascade used oligonucleotides that either mimicked a target miR or had the complementary sequence [28,48,49]. However, *in vivo*, miRs are vulnerable to degradation by serum ribonucleases [6]. The naked miR requires a vector to deliver it to the pancreatic β cells. Previous studies have looked at different vectors to protect and deliver miRs to the target cells. These include exosomes [50,51], micro-vesicles [52], and modified viruses [53].

Conclusion

The studies examining miRs in the human pancreatic islet model are limited. But they offer a new potential therapeutic strategy for diabetes mellitus. miRs fine tune existing biochemical pathways in human pancreatic islets. By inhibiting or stimulating these miR pathways with oligonucleotides, a patient's β cells and glycemic status can clinically be fine-tuned.

Acknowledgments

This work was funded by grants from the US National Institutes of Health (NIH) 1R01DK097380-01A1 (L.L.) and National Center for Research Resources (COBRE, 5P20RR018757-10), Juvenile Diabetes Research Foundation International (JDRF) foundation 1-2007-180 (L.L.) and Roger Williams Hospital Research fund (L.L.).

References

1. CDC. Annual Number (in Thousands) of New Cases of Diagnosed Diabetes among Adults Aged 18–79 Years. United States: 2014. 1980–2014
2. WHO. Diabetes. 2015
3. Nathan DM. Long-term complications of diabetes mellitus. *N Engl J Med*. 1993; 328:1676–1685. [PubMed: 8487827]
4. Menke A, Casagrande S, Geiss L, Cowie CC. Prevalence of and trends in diabetes among adults in the united states, 1988–2012. *JAMA*. 2015; 314:1021–1029. [PubMed: 26348752]
5. Ozcan S. Mini review: microRNA function in pancreatic β cells. *Mol Endocrinol*. 2014; 28:1922–1933. [PubMed: 25396300]
6. Nair VS, Pritchard CC, Tewari M, Ioannidis JP. Design and Analysis for Studying microRNAs in Human Disease: A Primer on -Omic Technologies. *Am J Epidemiol*. 2014; 180:140–152. [PubMed: 24966218]
7. Alvarez-Garcia I, Miska EA. MicroRNA functions in animal development and human disease. *Development*. 2005; 132:4653–4662. [PubMed: 16224045]
8. Walker MD. Role of MicroRNA in pancreatic beta-cells: where more is less. *Diabetes*. 2008; 57:2567–2568. [PubMed: 18820213]
9. Karolina DS, Armugam A, Tavintharan S, Wong MT, Lim SC, et al. MicroRNA 144 impairs insulin signaling by inhibiting the expression of insulin receptor substrate 1 in type 2 diabetes mellitus. *PLoS One*. 2011; 6:e22839. [PubMed: 21829658]
10. Latreille M, Hausser J, Stützer I, Zhang Q, Hastoy B, et al. MicroRNA-7a regulates pancreatic β cell function. *J Clin Invest*. 2014; 124:2722–2735. [PubMed: 24789908]

11. Wijesekara N, Zhang LH, Kang MH, Abraham T, Bhattacharjee A, et al. MiR-33a modulates ABCA1 expression, cholesterol accumulation, and insulin secretion in pancreatic islets. *Diabetes*. 2012; 61:653–658. [PubMed: 22315319]
12. Sebastiani G, Po A, Miele E, Ventriglia G, Ceccarelli E, et al. MicroRNA-124a is hyperexpressed in type 2 diabetic human pancreatic islets and negatively regulates insulin secretion. *Acta Diabetol*. 2015; 52:523–530. [PubMed: 25408296]
13. Kang MH, Zhang LH, Wijesekara N, de Haan W, Butland S, et al. Regulation of ABCA1 protein expression and function in hepatic and pancreatic islet cells by miR-145. *Arterioscler Thromb Vasc Biol*. 2013; 33:2724–2732. [PubMed: 24135019]
14. Locke JM, da Silva Xavier G, Dawe HR, Rutter GA, Harries LW. Increased expression of miR-187 in human islets from individuals with type 2 diabetes is associated with reduced glucose-stimulated insulin secretion. *Diabetologia*. 2014; 57:122–128. [PubMed: 24149837]
15. Bolmeson C, Esguerra JL, Salehi A, Speidel D, Eliasson L, et al. Difference in islet-enriched miRNAs in healthy and glucose intolerant human subjects. *Biochem and Biophys Res Commun*. 2011; 404:16–22. [PubMed: 21094635]
16. Xu G, Chen J, Jing G, Shalev A. Thioredoxin-interacting protein regulates insulin transcription through microRNA-204. *Nat Med*. 2013; 19:1141–1146. [PubMed: 23975026]
17. Rafiq I, da Silva Xavier G, Hooper S, Rutter GA. Glucose-stimulated preproinsulin gene expression and nuclear trans-location of pancreatic duodenum homeobox-1 require activation of phosphatidylinositol 3-kinase but not p38 MAPK/SAPK2. *J Biol Chem*. 2000; 275:15977–15984. [PubMed: 10821851]
18. Li X. MiR-37, a microRNA related to diabetes. *Gene*. 2014; 533:1–4. [PubMed: 24120394]
19. Melkman-Zehavi T, Oren R, Kredo-Russo S, Shapira T, Mandelbaum AD, et al. miRNAs control insulin content in pancreatic β -cells via down regulation of transcriptional repressors. *EMBO J*. 2011; 30:835–845. [PubMed: 21285947]
20. Jiang X, Shan A, Su Y, Cheng Y, Gu W, et al. miR-144/451 Promote Cell Proliferation via Targeting PTEN/AKT Pathway in Insulinomas. *Endocrinology*. 2015; 156:2429–2439. [PubMed: 25919186]
21. Wang Y, Liu J, Liu C, Naji A, Stoffers DA. MicroRNA-7 regulates the mTOR pathway and proliferation in adult pancreatic β -cells. *Diabetes*. 2013; 62:887–895. [PubMed: 23223022]
22. Tattikota SG, Rathjen T, McAnulty SJ, Wessels HH, Akerman I, et al. Argonaute2 mediates compensatory expansion of the pancreatic β cell. *Cell Metab*. 2014; 19:122–134. [PubMed: 24361012]
23. Vijayaraghavan J, Maggi EC, Crabtree JS. MiR-24 regulates menin in the endocrine pancreas. *Am J Physiol Endocrinol Metab*. 2014; 307:E84–E92. [PubMed: 24824656]
24. Crabtree JS, Scacheri PC, Ward JM, Garrett-Beal L, Emmert-Buck MR, et al. A mouse model of multiple endocrine neoplasia, type, develops multiple endocrine tumors. *Proc Natl Acad Sci U S A*. 2001; 98:1118–1123. [PubMed: 11158604]
25. Eizirik DL, Grieco FA. On the immense variety and complexity of circumstances conditioning pancreatic β -cell apoptosis in type 1 diabetes. *Diabetes*. 2012; 61:1661–1663. [PubMed: 22723273]
26. Roggli E, Gattesco S, Caille D, Briet C, Boitard C, et al. Changes in microRNA expression contribute to pancreatic β -cell dysfunction in pre-diabetic NOD mice. *Diabetes*. 2012; 61:1742–1751. [PubMed: 22537941]
27. Melnik BC. The pathogenic role of persistent milk signaling in mTORC1- and milk-microRNA-driven type 2 diabetes mellitus. *Curr Diabetes Rev*. 2015; 11:46–62. [PubMed: 25587719]
28. Roggli E, Britan A, Gattesco S, Lin-Marq N, Abderrahmani A, et al. Involvement of microRNAs in the cytotoxic effects exerted by pro-inflammatory cytokines on pancreatic beta-cells. *Diabetes*. 2010; 59:978–986. [PubMed: 20086228]
29. Zhu Y, You W, Wang H, Li Y, Qiao N, et al. MicroRNA-24/MODY gene regulatory pathway mediates pancreatic β -cell dysfunction. *Diabetes*. 2013; 62:3194–3206. [PubMed: 23761103]
30. Hogh KL, Uy CE, Asadi A, Baker RK, Riedel MJ, et al. Overexpression of peroxisome proliferator-activated receptor α in pancreatic β -cells improves glucose tolerance in diet-induced obese mice. *Exp Physiol*. 2013; 98:564–575. [PubMed: 23042378]

31. Hellemans K, Kerckhofs K, Hannaert JC, Martens G, Van Veldhoven P, et al. Peroxisome proliferator-activated receptor alpha-retinoid X receptor agonists induce beta-cell protection against palmitate toxicity. *FEBS J.* 2007; 274:6094–6105. [PubMed: 17970749]
32. Fred RG, Bang-Berthelsen CH, Mandrup-Poulsen T, Grunnet LG, Welsh N. High glucose suppresses human islet insulin biosynthesis by inducing miR-133a leading to decreased polypyrimidine tract binding protein-expression. *PLoS One.* 2010; 5:e10843. [PubMed: 20520763]
33. Rosero S, Bravo-Egana V, Jiang Z, Khuri S, Tsinoremas N, et al. MicroRNA signature of the human developing pancreas. *BMC Genomics.* 2010; 11:509. [PubMed: 20860821]
34. Wei R, Yang J, Liu GQ, Gao MJ, Hou WF, et al. Dynamic expression of microRNAs during the differentiation of human embryonic stem cells into insulin-producing cells. *Gene.* 2013; 518:246–255. [PubMed: 23370336]
35. Joglekar MV, Joglekar VM, Hardikar AA. Expression of islet-specific microRNAs during human pancreatic development. *Gene Expr Patterns.* 2009; 9:109–113. [PubMed: 18977315]
36. Poy MN, Eliasson L, Krutzfeldt J, Kuwajima S, Ma X, et al. A pancreatic islet-specific microRNA regulates insulin secretion. *Nature.* 2004; 432:226–230. [PubMed: 15538371]
37. Komatsu M, Takei M, Ishii H, Sato Y. Glucose-stimulated insulin secretion: a newer perspective. *J Diabetes Investig.* 2013; 4:511–516.
38. Straub SG, Sharp GW. Glucose-stimulated signaling pathways in biphasic insulin secretion. *Diabetes Metab Res Rev.* 2002; 18:451–463. [PubMed: 12469359]
39. Bratanova-Tochkova TK, Cheng H, Daniel S, Gunawardana S, Liu YJ, et al. Triggering and augmentation mechanisms, granule pools, and biphasic insulin secretion. *Diabetes.* 2002; 51:S83–S90. [PubMed: 11815463]
40. Åslusarz A, Pulakat L. The two faces of miR-29. *J Cardiovasc Med (Hagerstown).* 2015; 16:480–490. [PubMed: 25689084]
41. Lal A, Pan Y, Navarro F, Dykxhoorn DM, Moreau L, et al. miR-24-mediated down regulation of H2AX suppresses DNA repair in terminally differentiated blood cells. *Nat Struct Mol Biol.* 2009; 16:492–498. [PubMed: 19377482]
42. Lal A, Navarro F, Maher CA, Maliszewski LE, Yan N, et al. miR-24 inhibits cell proliferation by targeting E2F, MYC, and other cell-cycle genes via binding to “seedless” 3′ UTR microRNA recognition elements. *Mol Cell.* 2009; 35:610–625. [PubMed: 19748357]
43. Luo JZ, Xiong F, Al-Homsi AS, Ricordi C, Luo L. Allogeneic bone marrow cocultured with human islets significantly improves islet survival and function in vivo. *Transplantation.* 2013; 95:801–809. [PubMed: 23416682]
44. Joglekar MV, Parekh VS, Hardikar AA. Islet-specific microRNAs in pancreas development, regeneration and diabetes. *Indian J Exp Biol.* 2011; 49:401–408. [PubMed: 21702218]
45. Lynn FC, Skewes-Cox P, Kosaka Y, McManus MT, Harfe BD, et al. MicroRNA expression is required for pancreatic islet cell genesis in the mouse. *Diabetes.* 2007; 56:2938–2945. [PubMed: 17804764]
46. Correa-Medina M, Bravo-Egana V, Rosero S, Ricordi C, Edlund H, et al. MicroRNA miR-7 is preferentially expressed in endocrine cells of the developing and adult human pancreas. *Gene Expr Patterns.* 2009; 9:193–199. [PubMed: 19135553]
47. Chen H, Lan HY, Roukos DH, Cho WC. Application of microRNAs in diabetes mellitus. *J Endocrinol.* 2014; 222:R1–R10. [PubMed: 24781254]
48. Weiler J, Hunziker J, Hall J. Anti-miRNA oligonucleotides (AMOs): ammunition to target miRNAs implicated in human disease? *Gene Ther.* 2006; 13:496–502. [PubMed: 16195701]
49. Czech MP, Aouadi M, Tesz GJ. RNAi-based therapeutic strategies for metabolic disease. *Nat Rev Endocrinol.* 2011; 7:473–484. [PubMed: 21502982]
50. Johnsen KB, Gudbergsson JM, Skov MN, Pilgaard L, Moos T, et al. A comprehensive overview of exosome as drug delivery vehicle - endogenous nano-carrier for targeted cancer therapy. *Biochim Biophys Acta.* 2014; 1846:75–87. [PubMed: 24747178]
51. Melnik BC, John SM, Schmitz G. Milk is not just food but most likely a genetic transfection system activating mTORC1 signaling for postnatal growth. *Nutr J.* 2013; 12:103. [PubMed: 23883112]

52. Cantaluppi V, Biancone L, Figliolini F, Beltramo S, Medica D, et al. Microvesicles derived from endothelial progenitor cells enhances neo-angiogenesis of human pancreatic islets. *Cell Transplant*. 2012; 21:1305–1320. [PubMed: 22455973]
53. Trajkovski M, Hausser J, Soutschek J, Bhat B, Akin A, et al. MicroRNAs 103 and 107 regulate insulin sensitivity. 2011; 474:649–653.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 1.
 β -cell biochemical cascades. MicroRNAs “fine tune” the biochemical pathways found in beta cells. These four pathways are examined in this review.