Published in final edited form as:

J Acquir Immune Defic Syndr. 2017 June 01; 75(2): 203–210. doi:10.1097/QAI.0000000000001371.

Predictors and impact of self-reported suboptimal effort on estimates of prevalence of HIV-associated neurocognitive disorders

Andrew J. Levine*,

Department of Neurology, David Geffen School of Medicine at the University of California Los Angeles, Los Angeles 90095, USA

Eileen Martin,

Department of Psychiatry, Rush University Medical Center, Chicago, 60612, USA

Ned Sacktor,

Department of Neurology, Johns Hopkins University School of Medicine, Baltimore, 21205, USA

Cynthia Munro,

Departments of Psychiatry and Neurology, Johns Hopkins University School of Medicine, Baltimore, 21205, USA

James Becker, and

Departments of Psychiatry and Neurology, University of Pittsburgh, Pittsburgh, 15260, USA

for the Multicenter AIDS Cohort Study- Neuropsychology Working Group

Abstract

Background—Prevalence estimates of HIV-associated neurocognitive disorders (HAND) may be inflated. Estimates are determined via cohort studies in which participants may apply suboptimal effort on neurocognitive testing, thereby inflating estimates. Additionally, fluctuating HAND severity over time may be related to inconsistent effort. To address these hypotheses, we characterized effort in the Multicenter AIDS Cohort Study.

Methods—After neurocognitive testing, 935 participants (525 HIV-, 410 HIV+) completed the Visual Analogue Effort Scale (VAES), rating their effort from 0-100%. Those with <100% then indicated the reason(s) for suboptimal effort. K-means cluster analysis established 3 groups: high (mean=97%), moderate (79%), and low effort (51%). Rates of HAND and other characteristics were compared between groups. Linear regression examined predictors of VAES score. Data from 57 participants who completed the VAES at two visits was analyzed to characterize the longitudinal relationship between effort and HAND severity.

Results—Fifty-two percent of participants reported suboptimal effort (<100%) effort, with no difference between serostatus groups. Common reasons included “tired” (43%) and “distracted” (36%). The lowest effort group had greater ANI and MND diagnosis (25 and 33%) as compared to the moderate (23% and 15%) and the high effort groups (12% and 9%). Predictors of suboptimal

*Corresponding Author: ajlevine@mednet.ucla.edu, 562-477-5963 (phone); 510-280-2993 (fax).

effort were self-reported memory impairment, African-American race, and cocaine use. Change in effort between baseline and follow-up correlated with change in HAND severity.

Conclusion—Suboptimal effort appears to inflate estimated HAND prevalence and explain fluctuation of severity over time. A simple modification of study protocols to optimize effort is indicated by the results.

Keywords

Suboptimal effort; HIV-associated neurocognitive disorders; prevalence; visual analogue scale; neuropsychology of HIV

Introduction

Estimates of the prevalence of HIV-associated neurocognitive disorders (HAND) are considerably varied. Between 22-84% of infected individuals meet criteria at any one time, with an average of about 50% based on larger consortium studies¹⁻⁵. The majority of HAND diagnoses are mild, termed Asymptomatic Neurocognitive Impairment (ANI) according to current research criteria⁶. However, the inclusion of ANI in current diagnostic schema may have had the unintended consequence of high rates of false positive diagnoses, thereby inflating HAND prevalence estimates^{7,8}. This is primarily due to the low threshold required to be considered neurocognitively impaired. Indeed, a significant percentage of healthy HIV-uninfected individuals with no known neurologic or psychiatric illness would meet criteria for ANI, save for the fact that they are not HIV-infected⁸⁻¹².

Such findings do not invalidate ANI as a useful diagnosis. Indeed, a recent study found that individuals diagnosed with ANI at baseline progressed to symptomatic impairment faster than neurocognitively normal participants, as determined via self-report or performance-based measures¹³. While such findings underscore that ANI may be prodromal for more severe HAND, it does not quell the likelihood that many cases are misdiagnosed. Of note, that study did not report how many of the neurocognitively normal also progressed to ANI, or how many of those with ANI recovered to a neurocognitively normal status. This is important, as changes in HAND status is common in both directions (recovery and worsening)^{14,15,6}. The explanation for this variability in HAND severity across visits remains incomplete.

One largely unexplored factor in fluctuating HAND severity and inflation of HAND prevalence is the influence of inadequate effort on psychometric testing outcomes. Such *suboptimal effort*, as defined here, occurs when research participants perform below their potential on neurocognitive tests. That is, due to a one or a combination of several potential factors (e.g., fatigue, apathy, distraction, boredom, intoxication), participants' scores do not reflect their true ability. This is different from *feigned effort*, in which performance is intentionally deficient with the goal of appearing cognitively impaired, although *suboptimal effort* has previously been used synonymously with *feigned effort*¹⁶⁻¹⁸. Feigned effort is a well-documented phenomenon in several patient populations, generally in the context of forensic or disability evaluations in which there is potential for secondary gain¹⁹⁻²². A variety of instruments, collectively termed performance validity tests (PVTs), have been

developed for the purpose of identifying feigned effort. The sole effort study in the context of HAND used a PVT²³ among a research cohort. Not surprisingly, the participants almost all performed above the established cutoff for suspect effort. However, such tests are not valid for assessing suboptimal effort (as defined here), as their scoring criteria were developed for the detection of intentional response bias rather than insufficient implementation of cognitive abilities.

Anecdotal evidence suggests that a substantial portion of individuals enrolled in epidemiological studies may not put forth adequate and/or consistent effort on neurocognitive testing, an observation supported by empirical evidence in other research populations (e.g., college students seeking course credit)²⁴⁻²⁶. Such suboptimal effort may have substantial downstream effects on HAND prevalence estimates and our understanding of variability of HAND severity over time. To investigate the role of suboptimal effort in HIV neurocognitive studies, we developed a novel visual analogue measure that was integrated into the Multicenter AIDS Cohort Study (MACS) neurocognitive battery. Combined with the wide range of other information collected from MACS participants, this information was used to explore reason for, and predictors and consequences of, suboptimal effort. Our hypothesis was that suboptimal effort would contribute to over-estimation of HAND rates and help to explain variability in test performance over time.

Methods

This study was conducted within the MACS, which has followed gay and bisexual men since the early 1980s. The Visual Analogue Effort Scale (VAES) was developed by the authors specifically for investigating suboptimal effort. Visual analogue scales are easy to administer and allow measurement of mental states in a continuous fashion. They demonstrate equivalent or better psychometric properties compared to ordinal scales, particularly with regards to assessing subjective states^{27,28}. The VAES was administered to 935 participants at the conclusion of neurocognitive testing. Participants rated their effort on a line, with a range of 0-100%. Those who reported <100% effort (i.e., suboptimal effort) were directed to indicate the reason(s) for suboptimal effort. Fifty-seven participants completed the VAES at two consecutive visits, allowing for longitudinal analysis of effort and HAND severity in a subset of cases.

The following variables were included in our analysis of predictors and outcomes of suboptimal effort:

Neurocognitive functioning—Participants complete a battery of neuropsychological tests as part of the standard study protocol²⁹. This includes measures of working memory, learning, memory, executive functioning, motor functioning, and processing speed. T-scores were calculated using normative data with demographic corrections for age, education, and ethnicity. We examined both domain-specific T-scores and overall global neurocognitive functioning based on the average of the six domain T-scores.

HAND Severity—HAND status is determined for both HIV+ and HIV-uninfected participants via an algorithm developed by MACS investigators that adheres to the 2007

“Frascati” research criteria^{6,30}. HAND status is based on neurocognitive test performance and self-reported deficits in activities of daily living³¹. Participants are rated as neurocognitively normal, mildly impaired, moderately impaired, or severely impaired. The latter three correspond to Asymptomatic Neurocognitive Impairment, Minor Neurocognitive Disorder, and HIV-Associated Dementia.

Substance Use—MACS Participants complete a substance use questionnaire that assesses frequency of use during the six months prior to the visit, including none, monthly, weekly, and daily. Participants report frequency of use for alcohol, stimulants, marijuana/hashish, cocaine, opiates, or other recreational drugs (e.g., MDMA).

Depression—Depression severity is determined with the Center for Epidemiologic Studies Depression Scale (CES-D)³² as part of the standard MACS protocol. Scores on the CES-D are used as a continuous variable, with higher scores indicating greater degree of depression.

Memory self-rating—Participants are asked, “On a scale from 1 to 10, with 10 being normal for you, how would you rate your own memory ability now?”

Employment/Student status—Participants indicate whether they are employed full time or part time, in school, or retired.

Statistical analyses

Effort, as determined with the VAES, was used as a continuous variable. K-means cluster analysis categorized participants into one of three groups: High, Moderate, or Low effort, as described below. Differences in frequencies of reasons for suboptimal effort was compared between HIV status group and other categorical factors using chi-square analysis, whereas continuous variables (e.g., effort) were compared using ANOVA. Linear regression used employed to determine predictors of suboptimal effort. Outliers (>3 SD above or below mean) with regards to neurocognitive domain scores were removed.

Results

Suboptimal effort characteristics in MACS

The average VAES score was 91.4% (sd=13), with a range of 20-100. There was no difference between HIV- (mean = 91.9, sd = 12.6) and HIV+ (mean = 90.9, sd = 13.6; $F = 1.33$, $p = .249$). Just over half of the sample (51.7%) indicated suboptimal effort. While there was not a statistically significant difference in effort across the four study sites ($F = 1.69$, $p = .167$), UCLA was observed to have the lowest effort (90.5%), whereas the site with the highest effort was Johns Hopkins (94.3%).

Table 1 displays reasons provided by the MACS participants who acknowledged suboptimal effort. For the 483 individuals who indicated suboptimal (<100%) effort, the most common reasons were “Tired/Fatigued” (43.4%) and “Distracted/Poor concentration” (36%). Less than 8% of those reporting suboptimal effort indicated “unmotivated” as a cause. This may seem contradictory, as effort and motivation appear synonymous. However, motivation is considered a different construct than effort³³. Consider that one can feel unmotivated yet put

forth full effort. As such, motivation is a feeling, while effort has the added dimension of action. Over 31% responded “Other”. Common reasons provided among this subset were physical discomfort or illness (17%), memory difficulties (17%), boredom (15%), and anxiety (7%). Less common reasons (<5%) included hunger, learning disability, depression, testing environment, dislike for tests, and being in a hurry. There were no differences in reasons provided between serostatus groups.

Effort groups

K-means cluster analysis was used to create distinct Effort groups. We settled on a 3-cluster solution, with groups defined as High Effort (N=707, mean effort = 97%, Initial cluster center = 100%), Moderate Effort (N=175, mean effort = 79%, Initial cluster center = 60%), and Poor Effort (N=53, mean effort = 51%, Initial cluster center = 20%). We then examined how these effort groups differed. After correcting for multiple comparisons, groups differed in the cognitive domains of Executive, Speed, Learning, and Memory. Post hoc (Tukey's B) analysis showed that all domains, the Moderate and Poor Effort groups differed from the High Effort group, but not each other. Groups also differed with regards to education, with Poor effort group having significantly lower years of formal education compared to the Moderate and High effort groups. Groups did not differ with regards to age or depression (CES-D). African Americans and Hispanics were over-represented in the Low and Moderate effort groups as compared to Caucasian and “other” ethnicities.

Effect of suboptimal effort on HAND diagnoses

To begin to understand the effect of suboptimal effort on HAND prevalence, we examined the correlations between effort level and neurocognitive functioning. Pearson correlation analysis, following adjustment for multiple comparisons, revealed that effort was weakly correlated with all neurocognitive domain T-scores ($R = .23$, $p < .00001$), with the exception of working memory and motor functioning (Table 3). Correlations between all neurocognitive domains was significant ($p < .00001$).

We then examined how the effort groups differed in frequencies of HAND diagnoses. Note that overall, 14.9% of the sample met criteria for ANI (15.3% of HIV+ and 14.5% HIV-), 11.2% for MND (13% of HIV+ and 8.9% of HIV-), and 2.7% for HAD (3.7% of HIV+ and 1.7% of HIV-). The Poor Effort group had greatest percent of ANI and MND diagnosis (25% and 33%, respectively) as compared to the High Effort group (12% and 9%, respectively) and the Moderate Effort group (23% and 15%, respectively) ($\chi^2 = 58.12$, $p < .001$). Unexpectedly, the Poor Effort group had no cases of HAD, whereas the Moderate and High Effort groups had 4.1% and 2.6% with HAD, respectively.

Next, we looked at the relationship between change in effort and change in HAND severity between two study visits among 57 participants. This group had a mean age of 51 years ($sd = 16$), mean education of 14.8 years ($sd = 2.7$), and mean effort of 92.2% ($sd = 10\%$). Two-thirds (66.7%) were HIV+, 58% were reported suboptimal effort at baseline, and 65% were Caucasian. The mean change in effort was +1.1% ($sd = 7.7$), with a range of between -20 to +20%. Twenty-two (39%) of participants indicated no change in effort, whereas 16 indicated a decrease and 19 indicated an increase. For HAND severity, 39 (68%) were stable between

the two visits, 9 progressed to a more severe stage, and 7 improved to a less severe stage. The Spearman rank correlation between change in effort and HAND severity was significant ($r = -.335$, $p = .013$). In a linear regression model, with change in HAND severity as the criterion variable and change in effort, race, CES-D (at 2nd visit), cocaine use (at 2nd visit), and self-reported memory ability (at 2nd visit) as predictors, only change in effort remained in the final model (Adjusted $R^2 = .091$, R^2 change = .110, F change = 5.579, significance of F change = .023). Note that none of the participants involved in the longitudinal analysis reported cocaine use during the previous 6 months.

Predictors of effort level

We then sought to determine predictors of effort level based on the continuous score from the VAES. Based on initial correlations (not shown), the following predictor variables were included: age, education, employment status (employed or retired vs. unemployed and not retired), race (Caucasian vs. African American, regardless of Hispanic ethnicity), HIV status, depression (CES-D), alcohol use, cocaine use, and self-reported memory ability. In the final statistically significant model (Adjusted $R^2 = .079$, R^2 change = .006, $p = .028$), decreasing effort was predicted by lower self-reported memory ability ($\beta = .236$, $p < .001$), African American race ($\beta = -.092$, $p = .012$), and increasing frequency of cocaine use ($\beta = -.079$, $p = .028$).

To further investigate the role of African American race and cocaine use in suboptimal effort, we examined frequencies of cocaine use between race categories, as well as effort level between race categories (Table 5). Caucasians reported higher effort on the VAES (92.8%) as compared to African Americans (88.2%), regardless of Hispanic ethnicity ($F=21.39$, $P<.001$). African Americans also reported more frequent cocaine use and were over-represented in HAND diagnoses. To determine if the greater cocaine use among African Americans inflated this group's HAND prevalence, we removed all cocaine users and repeated the analyses. As shown in parentheses next to the aforementioned frequencies in Table 5, African Americans continued to have greater rates of HAND, confirming that both cocaine use and ethnicity are independent predictors of effort. Finally, when only the High effort group was used in these analyses, the rate of HAND among African Americans dropped from 41% to 25%, whereas rates for Caucasians dropped from 29% to 22%.

Discussion

While ANI may be a useful diagnostic category that designates some individuals as being in at risk for functional decline¹³, the low threshold required for diagnosis likely results in a large number of false-positive diagnostic errors^{7,8}. Additionally, as investigated here, when prevalence rates are determined via research cohorts, there is threat of inflated estimates due to suboptimal effort by study participants. Variable effort may also explain, in part, the apparent instability of HAND severity, such that a considerable proportion of individuals improve or decline across visits regardless of viral and immune factors^{6,30}. In this study, we examined the phenomenon of suboptimal effort in one of the largest HIV study cohorts, the MACS. The data derived from our novel measure, the VAES, indicate that while over 50% of participants reported suboptimal (<100%) effort, a much smaller number were considered

to exert low (N = 53, or 6%) or moderate (N = 175, or 19%) effort. Still the effect of these cases on HAND prevalence estimates was remarkable, with 58% of the low effort group having mild-to-moderate HAND, compared to 38% of the moderate effort group and 21% of the high effort group. Importantly, while African Americans were disproportionately represented in both the low effort group and as having HAND (41%, as compared to 25% of Caucasians), when those who indicated low or moderate effort were removed, the rates of HAND among African Americans and Caucasians nearly converged (25% and 22%, respectively). Finally, it is worth noting that the site with the best effort overall (Johns Hopkins) had the highest proportion of African Americans, and the site with the poorest effort overall (UCLA) had the lowest proportion. Johns Hopkins differed from the other three in that the neurocognitive testing is completed on a separate day then the other study procedures. At the other study sites, the neurocognitive testing is generally the last of up to 3 hours of other procedures, including physical exam, blood draw, and filling out numerous questionnaires. These results indicate that suboptimal effort is a significant factor behind the disproportionate number of African Americans meeting criteria for HAND in the MACS, and that a slight modification in study protocol can mitigate this.

When looking at effort overall, the strongest predictor was self-reported memory ability. That is, participants who rated their memory ability as poor also reported lower effort. This may reflect an attitude more than an accurate self-assessment, as the correlation between self-reported memory ability and effort was stronger than those with neurocognitive domain T-scores (results not shown). In this scenario, less effort is exerted because the individual does not believe they will be able to perform well due to their memory deficit. Alternatively, this finding may reflect an actual neurocognitive impairment that is not adequately detected by the MACS neuropsychological test battery. Another predictor of suboptimal effort was education attainment. Note that neurocognitive test scores are standardized according to age, race, and education, so the findings here are not due to confounds inherent in the normative data. One interpretation is that education attainment is an indicator of overall attitude about testing, or reflects the underlying motivation of participants to cognitive challenges.

The relationship between effort and HAND severity was significant. Furthermore, with regards to intra-individual variability in HAND severity over time, regression analysis revealed that it was only change in effort between baseline and follow-up visit that predicted change in HAND severity; depression, race, cocaine use, and self-reported memory ability (all significant predictors in cross-sectional analyses, save for depression), were not significant predictors. It is also notable that 33% were HIV-uninfected, so viral and immune factors were unlikely to explain the change in HAND severity. Indeed, chi-square analysis did not reveal significant differences in change in HAND severity or effort between HIV+ and HIV-uninfected participants (not shown). This finding underscores the significant contribution of effort to HAND variability over time. Furthermore, there has been recent interest in intra-individual neurocognitive variability in HIV as a behavioral marker of impending disability³⁴, functional deficits³⁵, and cognitive dyscontrol due to the combined effects of age and HIV in older (> 50 years old) adults³⁶. Those studies did not consider effort, leaving this question to be addressed in the future. Indeed, studies in traumatic brain injury have indicated that intra-individual variability in neurocognitive impairment increases both as a function of neuropathology and of suboptimal effort¹⁷.

One might argue that effort is related to neurocognitive functioning, and as such, it is not surprising that those with low effort also have higher risk for HAND. However, the data do not support this. First, the correlations between VAES and neurocognitive domain scores, while statistically significant, were not strong, and were similar across all effort groups. Secondly, if it were true, one would expect a relatively greater number of the Low Effort group to have severe HAND (i.e., HAD). In fact, no one in the Low Effort group had HAD, whereas 4.1% of the Moderate and 2.6% of the High Effort group members had HAD. This unusual finding requires further exploration. One possibility is that individuals who are truly impaired (e.g., with HAD) and aware of their deficits and put forth better effort because they are more invested in learning about their true level of functioning, whereas individuals who do not perceive themselves to have cognitive deficits are less interested in what the testing may reveal.

Together with the predictors identified here, the reasons provided by suboptimal responders may enable modification of study protocols to ensure adequate effort. The most common reasons provided were “tired/fatigued” (43%) and “poor concentration/distracted” (36%). Considering the aforementioned difference in site protocols, it would be expected that these reasons would be less common at Johns Hopkins. Indeed, further examination of the data reveal that only 19% and 12% of respondents at Johns Hopkins indicated “tired/fatigued” and “distracted/poor concentration” as reasons for suboptimal effort, respectively. As such, modifying study protocols such that neurocognitive testing is completed when participants are fresh is strongly indicated by these results.

We acknowledge several limitations of this study. Firstly, we did not employ embedded PVTs in our primary analyses, despite the fact that several are included in the MACS neurocognitive battery. Embedded measures are portions of standard neurocognitive tests further developed for detecting feigned effort. These are essentially aspects of standard neurocognitive tests that are fairly easy to obtain perfect or near perfect scores on. Like standard PVTs, these embedded measures have been studied for their utility detecting feigned impairment among groups considered more likely to feign impairment (e.g., mTBI patients in litigation), as well as among individuals instructed to complete the tests as if they were feigning impairment. Like other PVTs, these cutoffs would not be useful for detecting suboptimal effort, as they were developed for identifying feigned vs. true effort³⁷⁻⁴⁰. It could be argued that using the scores in a continuous fashion, rather than cutoff scores, may be a better approach. However, our data (not shown) indicate that such scores correlate much more strongly with other neurocognitive tests score than effort as measured with the VAES. Therefore, neither explicit PVTs or embedded measures are useful for assessing suboptimal effort in this context. Secondly, the VAES has not been tested for its validity or reliability. However, we point out that for adequate testing of this measure's psychometric qualities, a criterion measure of suboptimal effort is required. To our knowledge no such measure exists. As such, this study has generated important validity and reliability data for the VAES, which we hope can be used for assessing suboptimal effort in other contexts. Thirdly, the HAND classification system used by the MACS does not adequately assess for other causes of neurocognitive impairment. This is especially limiting considering that HAND is a diagnosis based on exclusion of other causes. However, like the MACS, other large cohort studies lack the resources to conduct the comprehensive diagnostic testing required to rule out other

causes. Finally, there are other factors likely influencing effort, or modifying neurocognitive performance as a function of effort. Examples include personality^{41,42} and acculturation^{43,44}. Thus, further investigation of factors that mitigate or augment effort in the context of research studies might consider factors not considered here.

One final point about inflated prevalence estimate; the finding reported here that almost as many HIV- as HIV+ MACS participants meet criteria for HAND underscores the poor specificity of current diagnostic criteria. Even when prevalence estimates are based only on those participants who reported full effort, 25.7% of HIV+ participants and 20.6% of the uninfected participants meet HAND criteria. This is especially troubling because prevalence estimates are generally cited from studies that use only HIV+ cases^{5,14,45,46}. While one more recent study included a HIV-uninfected comparison group³, a large proportion of uninfected individuals still met criteria for HAND. Further, the markedly higher rates of comorbidities (e.g., substance use and depression) among the HIV+ sample complicates interpretability of that study. Considering that the MACS cohort possesses comparatively fewer comorbidities and that its HIV+ and HIV-uninfected participants are more similar with regards to comorbidities, the data presented here more accurately reflect the reality of the inadequacies of the current diagnostic schema. Fortunately, growing awareness of this has led some to develop alternative strategies for classifying HAND⁴⁷.

To summarize, suboptimal effort is a true phenomenon in HIV research studies that has subsequent influence on HAND prevalence estimates, public policy, and designation of resources. Suboptimal effort is also a factor behind variability of HAND severity over time. We have identified a very simple solution for improving effort. We strongly recommend that future determination of HAND prevalence rates and progression over time consider the phenomenon of suboptimal effort and employ alternative statistical methods.

Acknowledgments

Funding: Data in this manuscript were collected by the Multicenter AIDS Cohort Study (MACS). MACS (Principal Investigators): Johns Hopkins University Bloomberg School of Public Health (Joseph Margolick), U01-AI35042; Northwestern University (Steven Wolinsky), U01-AI35039; University of California, Los Angeles (Roger Detels, Oto Martinez-Maza), U01-AI35040; University of Pittsburgh (Charles Rinaldo), U01-AI35041; the Center for Analysis and Management of MACS, Johns Hopkins University Bloomberg School of Public Health (Lisa Jacobson, Gypsyamber D'Souza), UM1-AI35043. The MACS is funded primarily by the National Institute of Allergy and Infectious Diseases (NIAID), with additional co-funding from the National Cancer Institute (NCI), the National Institute on Drug Abuse (NIDA), and the National Institute of Mental Health (NIMH). Targeted supplemental funding for specific projects was also provided by the National Heart, Lung, and Blood Institute (NHLBI), and the National Institute on Deafness and Communication Disorders (NIDCD). MACS data collection is also supported by UL1-TR001079 (JHU ICTR) from the National Center for Advancing Translational Sciences (NCATS) a component of the National Institutes of Health (NIH), and NIH Roadmap for Medical Research. The contents of this publication are solely the responsibility of the authors and do not represent the official views of the National Institutes of Health (NIH), Johns Hopkins ICTR, or NCATS. The MACS website is <http://aidscohortstudy.org>

References

1. Cysique LA, Maruff P, Brew BJ. Prevalence and pattern of neuropsychological impairment in human immunodeficiency virus-infected/acquired immunodeficiency syndrome (HIV/AIDS) patients across pre- and post-highly active antiretroviral therapy eras: a combined study of two cohorts. *J Neurovirol.* 2004; 10(6):350–357. [PubMed: 15765806]

2. Becker JT, Lopez OL, Dew MA, Aizenstein HJ. Prevalence of cognitive disorders differs as a function of age in HIV virus infection. *Aids*. 2004; 18(1):S11–18. [PubMed: 15075493]
3. Heaton RK, Franklin DR, Ellis RJ, et al. HIV-associated neurocognitive disorders before and during the era of combination antiretroviral therapy: differences in rates, nature, and predictors. *J Neurovirol*. 2011; 17(1):3–16. [PubMed: 21174240]
4. Bonnet F, Amieva H, Marquant F, et al. Cognitive disorders in HIV-infected patients: are they HIV-related? *AIDS*. 2013; 27(3):391–400. [PubMed: 23079813]
5. Simioni S, Cavassini M, Annoni JM, et al. Cognitive dysfunction in HIV patients despite long-standing suppression of viremia. *AIDS*. 2010; 24(9):1243–1250. [PubMed: 19996937]
6. Antinori A, Arendt G, Becker JT, et al. Updated research nosology for HIV-associated neurocognitive disorders. *Neurology*. 2007; 69(18):1789–1799. [PubMed: 17914061]
7. Gisslen M, Price RW, Nilsson S. The definition of HIV-associated neurocognitive disorders: are we overestimating the real prevalence? *BMC Infect Dis*. 2011; 11:356. [PubMed: 22204557]
8. Meyer AC, Boscardin WJ, Kwasa JK, Price RW. Is it time to rethink how neuropsychological tests are used to diagnose mild forms of HIV-associated neurocognitive disorders? Impact of false-positive rates on prevalence and power. *Neuroepidemiology*. 2013; 41(3-4):208–216. [PubMed: 24157541]
9. Schretlen DJ, Munro CA, Anthony JC, Pearlson GD. Examining the range of normal intraindividual variability in neuropsychological test performance. *J Int Neuropsychol Soc*. 2003; 9(6):864–870. [PubMed: 14632245]
10. Palmer BW, Boone KB, Lesser IM, Wohl MA. Base rates of “impaired” neuropsychological test performance among healthy older adults. *Arch Clin Neuropsychol*. 1998; 13(6):503–511. [PubMed: 14590634]
11. Schretlen DJ, Testa SM, Winicki JM, Pearlson GD, Gordon B. Frequency and bases of abnormal performance by healthy adults on neuropsychological testing. *J Int Neuropsychol Soc*. 2008; 14(3):436–445. [PubMed: 18419842]
12. Binder LM, Iverson GL, Brooks BL. To err is human: “abnormal” neuropsychological scores and variability are common in healthy adults. *Arch Clin Neuropsychol*. 2009; 24(1):31–46. [PubMed: 19395355]
13. Grant I, Franklin DR Jr, Deutsch R, et al. Asymptomatic HIV-associated neurocognitive impairment increases risk for symptomatic decline. *Neurology*. 2014; 82(23):2055–2062. [PubMed: 24814848]
14. Robertson KR, Smurzynski M, Parsons TD, et al. The prevalence and incidence of neurocognitive impairment in the HAART era. *AIDS*. 2007; 21(14):1915–1921. [PubMed: 17721099]
15. Heaton RK, Franklin DR Jr, Deutsch R, et al. Neurocognitive change in the era of HIV combination antiretroviral therapy: the longitudinal CHARTER study. *Clin Infect Dis*. 2015; 60(3):473–480. [PubMed: 25362201]
16. Burton RL, Enright J, O’Connell ME, Lanting S, Morgan D. RBANS embedded measures of suboptimal effort in dementia: effort scale has a lower failure rate than the effort index. *Arch Clin Neuropsychol*. 2015; 30(1):1–6. [PubMed: 25472686]
17. Hill BD, Rohling ML, Boettcher AC, Meyers JE. Cognitive intra-individual variability has a positive association with traumatic brain injury severity and suboptimal effort. *Arch Clin Neuropsychol*. 2013; 28(7):640–648. [PubMed: 23832096]
18. Armistead-Jehle P, Buican B. Comparison of select Advanced Clinical Solutions embedded Effort measures to the Word Memory Test in the detection of suboptimal effort. *Arch Clin Neuropsychol*. 2013; 28(3):297–301. [PubMed: 23535704]
19. Greve KW, Bianchini KJ, Black FW, et al. The prevalence of cognitive malingering in persons reporting exposure to occupational and environmental substances. *Neurotoxicology*. 2006; 27(6):940–950. [PubMed: 16904749]
20. Greve KW, Lotz KL, Bianchini KJ. Observed versus estimated IQ as an index of malingering in traumatic brain injury: classification accuracy in known groups. *Appl Neuropsychol*. 2008; 15(3):161–169. [PubMed: 18726736]
21. Iverson GL. Ethical issues associated with the assessment of exaggeration, poor effort, and malingering. *Appl Neuropsychol*. 2006; 13(2):77–90. [PubMed: 17009881]

22. Larrabee GJ. Detection of malingering using atypical performance patterns on standard neuropsychological tests. *Clin Neuropsychol*. 2003; 17(3):410–425. [PubMed: 14704892]
23. Woods SP, Conover E, Weinborn M, et al. Base rate of Hiscock Digit Memory Test failure in HIV-associated neurocognitive disorders. *Clin Neuropsychol*. 2003; 17(3):383–389. [PubMed: 14704888]
24. An KY, Zakzanis KK, Joordens S. Conducting research with non-clinical healthy undergraduates: does effort play a role in neuropsychological test performance? *Arch Clin Neuropsychol*. 2012; 27(8):849–857. [PubMed: 23027440]
25. DeRight J, Jorgensen RS. I just want my research credit: frequency of suboptimal effort in a non-clinical healthy undergraduate sample. *Clin Neuropsychol*. 2015; 29(1):101–117. [PubMed: 25494327]
26. Ross TP, Poston AM, Rein PA, Salvatore AN, Wills NL, York TM. Performance Invalidity Base Rates Among Healthy Undergraduate Research Participants. *Arch Clin Neuropsychol*. 2016; 31(1):97–104. [PubMed: 26490230]
27. Grant S, Aitchison T, Henderson E, et al. A comparison of the reproducibility and the sensitivity to change of visual analogue scales, Borg scales, and Likert scales in normal subjects during submaximal exercise. *Chest*. 1999; 116(5):1208–1217. [PubMed: 10559077]
28. Reips UD, Funke F. Interval-level measurement with visual analogue scales in Internet-based research: VAS Generator. *Behav Res Methods*. 2008; 40(3):699–704. [PubMed: 18697664]
29. Becker JT, Kingsley LA, Molsberry S, et al. Cohort Profile: Recruitment cohorts in the neuropsychological substudy of the Multicenter AIDS Cohort Study. *International journal of epidemiology*. 2014
30. Sacktor N, Skolasky RL, Seaberg E, et al. Prevalence of HIV-associated neurocognitive disorders in the Multicenter AIDS Cohort Study. *Neurology*. 2016; 86(4):334–340. [PubMed: 26718568]
31. Lawton MP, Brody EM. Assessment of older people: self-maintaining and instrumental activities of daily living. *Gerontologist*. 1969; 9(3):179–186. [PubMed: 5349366]
32. Radloff LS. The CES-D scale: A self-report depression scale for research in the general population. *Applied Psychological Measurement*. 1977; 1(1):385–401.
33. Foussias G, Siddiqui I, Fervaha G, et al. Motivated to do well: an examination of the relationships between motivation, effort, and cognitive performance in schizophrenia. *Schizophr Res*. 2015; 166(1-3):276–282. [PubMed: 26008882]
34. Morgan EE, Woods SP, Grant I. Group HIVNRP. Intra-individual neurocognitive variability confers risk of dependence in activities of daily living among HIV-seropositive individuals without HIV-associated neurocognitive disorders. *Arch Clin Neuropsychol*. 2012; 27(3):293–303. [PubMed: 22337933]
35. Thaler NS, Sayegh P, Arentoft A, Thames AD, Castellon SA, Hinkin CH. Increased Neurocognitive Intra-individual Variability Is Associated With Declines in Medication Adherence in HIV-Infected Adults. *Neuropsychology*. 2015
36. Morgan EE, Woods SP, Delano-Wood L, Bondi MW, Grant I. Group HIVNRP. Intraindividual variability in HIV infection: evidence for greater neurocognitive dispersion in older HIV seropositive adults. *Neuropsychology*. 2011; 25(5):645–654. [PubMed: 21574712]
37. Iverson GL, Lange RT, Green P, Franzen MD. Detecting exaggeration and malingering with the trail making test. *Clin Neuropsychol*. 2002; 16(3):398–406. [PubMed: 12607151]
38. Arentsen TJ, Boone KB, Lo TT, et al. Effectiveness of the Comalli Stroop Test as a measure of negative response bias. *Clin Neuropsychol*. 2013; 27(6):1060–1076. [PubMed: 23742292]
39. Reedy SD, Boone KB, Cottingham ME, et al. Cross validation of the Lu and colleagues (2003) Rey-Osterrieth Complex Figure Test effort equation in a large known-group sample. *Arch Clin Neuropsychol*. 2013; 28(1):30–37. [PubMed: 23232864]
40. Boone KB, Lu P, Wen J. Comparison of various RAVLT scores in the detection of noncredible memory performance. *Arch Clin Neuropsychol*. 2005; 20(3):301–319. [PubMed: 15797167]
41. Temple RO, McBride AM, David Horner MD, Taylor RM. Personality characteristics of patients showing suboptimal cognitive effort. *Clin Neuropsychol*. 2003; 17(3):402–409. [PubMed: 14704891]

42. Whiteside D, Clinton C, Diamonti C, et al. Relationship between suboptimal cognitive effort and the clinical scales of the Personality Assessment Inventory. *Clin Neuropsychol*. 2010; 24(2):315–325. [PubMed: 20077355]
43. Boone KB, Victor TL, Wen J, Razani J, Ponton M. The association between neuropsychological scores and ethnicity, language, and acculturation variables in a large patient population. *Arch Clin Neuropsychol*. 2007; 22(3):355–365. [PubMed: 17320344]
44. Razani J, Burciaga J, Madore M, Wong J. Effects of acculturation on tests of attention and information processing in an ethnically diverse group. *Arch Clin Neuropsychol*. 2007; 22(3):333–341. [PubMed: 17298874]
45. Tozzi V, Balestra P, Bellagamba R, et al. Persistence of neuropsychologic deficits despite long-term highly active antiretroviral therapy in patients with HIV-related neurocognitive impairment: prevalence and risk factors. *J Acquir Immune Defic Syndr*. 2007; 45(2):174–182. [PubMed: 17356465]
46. Heaton RK, Clifford DB, Franklin DR Jr, et al. HIV-associated neurocognitive disorders persist in the era of potent antiretroviral therapy: CHARTER Study. *Neurology*. 2010; 75(23):2087–2096. [PubMed: 21135382]
47. Su T, Schouten J, Geurtsen GJ, et al. Multivariate normative comparison, a novel method for more reliably detecting cognitive impairment in HIV infection. *AIDS*. 2015; 29(5):547–557. [PubMed: 25587908]

Table 1
Reasons for suboptimal effort

Reasons for Suboptimal Effort	HIV-	HIV+	Total	Chi-square
Low Motivation/Apathy	9.4%	6.3%	7.6%	1.52; p=.23
Tired/Fatigued	41.9%	44.6%	43.4%	.35; p=.57
Distracted/Poor concentration	37.9%	34.6%	36%	.56; p=.49
Under influence of drugs or alcohol	0.5%	1.1%	0.8%	.53; p=.64
Did not understand instructions	0.5%	0.7%	0.6%	.12; p=1.0
Other	31%	32%	31.6%	.05; p=.84

Table 2
Demographic and neurocognitive comparisons across effort groups

	Low Effort	Moderate Effort	High Effort	
Effort	50.5 (10.9)	79.3 (5.5)	97.4 (4)	--
Neurocognitive				
Domains	46.4 (8.6)	48 (10.1)	50.9 (9.7)	9.93, p<.0001
Executive	46.4 (8.8)	46.9 (9.8)	50.7 (10.1)	12.94, p<.0001
Speed of Processing	47.3 (11)	46.9 (9.9)	48.7 (8.7)	2.78, p=.06
Working Memory	44.6 (10.6)	45.9 (9.9)	51.5 (9.2)	34.12, p<.0001
Learning	45.6 (9.9)	45.9 (9.8)	51.3 (8.6)	32.27, p<.0001
Memory	43.3 (10.9)	44.6 (10.1)	45.8 (9.9)	1.95, p=.14
Motor				
Age	53.7 (12.8)	53.5 (12.1)	55.1 (12.4)	1.34, p=.26
Education	13.9 (2.5)	15.1 (2.7)	15.7 (2.6)	11.76, p<.0001
CES-D	13.5 (11.3)	11.7 (10.4)	10.5 (11.6)	2.0, p=.14
Ethnicity				X ² = 30.89, p<.0001
Caucasian	3.1%	15.4%	81.4%	
African American	9.9%	25.1%	64.9%	
Hispanic	8.2%	23.8%	68%	
Other	0%	14.3%	85.7%	

Table 3

Correlations between effort and neurocognitive domain T-scores

	Effort	Executive	Processing Speed	Working Memory	Learning	Memory
Executive	.153 *					
Processing Speed	.162 *	.683 *				
Working Memory	.065	.375 *	.417 *			
Learning	.223 *	.458 *	.389 *	.259 *		
Memory	.211 *	.413 *	.361 *	.230 *	.870 *	
Motor	.075	.372 *	.365 *	.214 *	.325 *	.261 *

* statistically significant ($P < .00001$)

Table 4
HAND ratings across effort groups

	Low Effort	Moderate Effort	High Effort
HAND Rating	$X^2 = 58.12, p < .0001$		
Normal	42.3%	57.9%	76.6%
ANI	25%	22.8%	12.2%
MCD	32.7%	15.2%	8.6%
HAD	0%	4.1%	2.6%

Table 5
Racial characterization in regards to cocaine use and HAND severity

	African American	Caucasian	
Effort	88.2%	92.8%	F=21.4, p<.001
Cocaine Use			X ² = 52.3, p<.001
Daily	0%	1.8%	
Weekly	0.2%	4.9%	
Monthly	0.5%	4%	
HAND			X ² = 42, p < .001
ANI	14.7% (14.6%) [11.4%]	15.7% (15.3%) [13%]	(X ² = 36.4, p <.001) *
MND	20.1% (18.8%) [12.8%]	8.1% (7.9%) [7.9%]	[X ² =13.1, p = .004] **
HAD	5.8% (6.3%) [5.4%]	1.1% (1.2%) [1.2%]	

*
 () = Repeat of analysis without cocaine users

**
 [] = Repeat of analysis with only High effort individuals