Published in final edited form as:

Bio Protoc. 2016 October 5; 6(19): . doi:10.21769/BioProtoc.1944.

ASC-particle-induced Peritonitis

Lucia de Almeida, Andrea Dorfleitner*, and Christian Stehlik*

Division of Rheumatology, Department of Medicine, Feinberg School of Medicine, Northwestern University, Chicago, USA

Abstract

In response to pathogen infection and tissue damage, inflammasome sensors such as NLRP3 and AIM2 are activated, which triggers PYRIN domain (PYD)-mediated ASC nucleation, followed by self-perpetuating ASC polymerization, which ultimately culminates in caspase-1 activation, interleukin (IL)-1 β and IL-18 processing and release and pyroptosis (Ratsimandresy *et al.*, 2013; Cai *et al.*, 2014). Inflammasomes release not only cytokines, but also the polymeric ASC danger particles (pASC) by pyroptosis, which perpetuate and propagate inflammasome responses to bystander cells to engage cell intrinsic ASC and caspase-1 (Baroja-Mazo *et al.*, 2014; Franklin *et al.*, 2014). In this protocol we describe intraperitoneal injection of polymeric ASC particles as a danger signal and measure neutrophil infiltration and levels of the pro-inflammatory cytokine IL-1 β by ELISA in the peritoneal lavage (de Almeida *et al.*, 2015).

Materials and Reagents

1. 5 ml syringes (BD, catalog number: 309646)
2. 25 G $\times$ 1 1/2 needles (BD, catalog number: 305127)
3. Conical tubes (15 ml) (Corning, Falcon[®], catalog number: 352095)
4. C57BL/6 mice, typically of 8–12 weeks old (male or female)
5. 1×10^5 pASC-GFP particles generated from stable or transiently expressing HEK293 cells and sorted by flow cytometry as described (Fernandes-Alnemri *et al.*, 2007; Fernandes-Alnemri and Alnemri, 2008)
6. $1 \times$ Dulbecco's phosphate-buffered saline (DPBS) (Corning, catalog number: 21-030-CV)
7. HEPES
8. KOH
9. Magnesium chloride (MgCl₂)
10. EGTA
11. CHAPS

*For correspondence: a-dorfleitner@northwestern.edu; c-stehlik@northwestern.edu.

12. cOmplete protease inhibitor cocktail (Roche Diagnostics, catalog number: 11697498001)
13. IL-1 β ELISA kit (BD, catalog number: 559603)
14. Lysis buffer (see Recipes)

Equipment

1. Centrifuge
2. Fluorescence microscope
3. Flow cytometry
4. Surgical instruments such as tweezers and scissors
5. Biosafety cabinet

Procedure

1. Isolate pASC-GFP particles from 10^7 HEK293 cells stable or transiently transfected with ASC-GFP as described previously (Fernandes-Alnemri *et al.*, 2007; Fernandes-Alnemri and Alnemri, 2008; Martín-Sánchez *et al.*, 2015) or by flow cytometry.
2. Remove media from cells, add DPBS and gently scrape cells. Spin down cells at $1,500 \times g$ for 5 min. Discard supernatant.
3. Prepare total cell lysates by hypotonic lysis of cell pellets in 20 mM HEPES-KOH, pH 7.5, 5 mM MgCl₂, 0.5 mM EGTA, 0.1% CHAPS, supplemented with protease inhibitors. Use a syringe with 25 G needle to lyse the cells by syringing several times. Centrifuge cell lysates full speed to obtain cell lysates supernatant.
4. Induce aggregation of ASC-GFP by incubation of cell lysates supernatant at 37 °C for 30 min as previously described (Fernandes-Alnemri *et al.*, 2007; Fernandes-Alnemri and Alnemri, 2008). Typical amount of ASC-GFP particles recovered is between 2×10^5 and 4×10^5 per million cells.
5. Add a small amount of supernatant to a microscope slide to confirm ASC polymerization into 1–3 μ m aggregates by fluorescence microscopy using a standard GFP excitation/emission filter set (excitation: 484 nm; emission: 507 nm) (Figure 1).
6. Sort ASC-GFP particles (1–2 μ m) by flow cytometry by gating for small particles based on forward scatter versus side scatter. Next start by gating for FITC positive particles. Later confirm ASC-GFP particles by fluorescent microscopy (Figure 1). Particles can be stored at 4 °C for one year.
7. Intraperitoneally inject mice with 1×10^5 pASC-GFP particles or the same volume DPBS for control mice (injection volume approximately 200 μ l).
8. 4 h later sacrifice mice, cut the skin and expose the peritoneal wall (Video 1).

9. Inject 4 ml of DPBS into the peritoneal cavity.
10. Shake the mouse to wash the peritoneal cavity.
11. Use the syringe to recover DPBS injected by gradually pulling out the plunge.
12. Remove the needle and add content to a conical tube.
13. Spin down the cells at $1,200 \times g$ for 5 min.
14. Transfer supernatant to fresh conical tubes and measure IL-1 β by ELISA. Typical IL-1 β concentrations are up to 2 ng/ml and a representative result is shown in our previous publication (de Almeida *et al.*, 2015).

Representative data

Figure 1. Total cell lysates from stable ASC-GFP-expressing HEK293 cells were GFP-sorted by flow cytometry after inducing ASC polymerization and control (Ctrl) and ASC-GFP (pASC)-containing fractions analyzed by fluorescence microscopy
HEK293 cell lysates were used as a negative control.

Notes

1. ASC-GFP particles should be protected from light until use.

Recipes

1. Lysis buffer
 - 20 mM HEPES-KOH, pH 7.5
 - 5 mM MgCl₂
 - 0.5 mM EGTA

0.1% CHAPS

Protease inhibitors

Acknowledgments

This protocol was adapted from a previously published study (de Almeida *et al.*, 2015). This work was supported by grants the National Institutes of Health (AI099009, AI120625, AI120618 and AR064349 to C.S., AR066739 to A.D., AI120625 and AI120618 to C.S. and A.D., T32AR007611 to L.d.A. and the American Heart Association 13GRNT17110117 to C.S.).

References

1. Baroja-Mazo A, Martin-Sanchez F, Gomez AI, Martinez CM, Amores-Iniesta J, Compan V, Barbera-Cremades M, Yague J, Ruiz-Ortiz E, Anton J, Bujan S, Couillin I, Brough D, Arostegui JJ, Pelegrin P. The NLRP3 inflammasome is released as a particulate danger signal that amplifies the inflammatory response. *Nat Immunol.* 2014; 15(8):738–748. [PubMed: 24952504]
2. Bondar VS, Puzyr AP. Use of nanodiamond particles for rapid isolation of recombinant apoobelin from *Escherichia coli*. *Dokl Biochem.* 2000; 373(1–6):129–131. [PubMed: 11002840]
3. Cai X, Chen J, Xu H, Liu S, Jiang QX, Halfmann R, Chen ZJ. Prion-like polymerization underlies signal transduction in antiviral immune defense and inflammasome activation. *Cell.* 2014; 156(6):1207–1222. [PubMed: 24630723]
4. de Almeida L, Khare S, Misharin AV, Patel R, Ratsimandresy RA, Wallin MC, Perlman H, Greaves DR, Hoffman HM, Dorfleutner A, Stehlik C. The PYRIN domain-only protein POP1 inhibits inflammasome assembly and ameliorates inflammatory disease. *Immunity.* 2015; 43(2):264–276. [PubMed: 26275995]
5. Fernandes-Alnemri T, Alnemri ES. Chapter thirteen assembly, purification, and assay of the activity of the ASC pyroptosome. *Methods Enzymol.* 2008; 442:251–270. [PubMed: 18662574]
6. Fernandes-Alnemri T, Wu J, Yu JW, Datta P, Miller B, Jankowski W, Rosenberg S, Zhang J, Alnemri ES. The pyroptosome: a supramolecular assembly of ASC dimers mediating inflammatory cell death via caspase-1 activation. *Cell Death Differ.* 2007; 14(9):1590–1604. [PubMed: 17599095]
7. Franklin BS, Bossaller L, De Nardo D, Ratter JM, Stutz A, Engels G, Brenker C, Nordhoff M, Mirandola SR, Al-Amoudi A, Mangan MS, Zimmer S, Monks BG, Fricke M, Schmidt RE, Espevik T, Jones B, Jarnicki AG, Hansbro PM, Busto P, Marshak-Rothstein A, Hornemann S, Aguzzi A, Kastenmuller W, Latz E. The adaptor ASC has extracellular and ‘prionoid’ activities that propagate inflammation. *Nat Immunol.* 2014; 15(8):727–737. [PubMed: 24952505]
8. Martín-Sánchez F, Gómez AI, Pelegrín P. Isolation of particles of recombinant ASC and NLRP3. *Bio-protocol.* 2015; 5(10):e1480.
9. Ratsimandresy RA, Dorfleutner A, Stehlik C. An update on PYRIN domaincontaining pattern recognition receptors: from immunity to pathology. *Front Immunol.* 2013; 4:440. [PubMed: 24367371]

Video 1. Peritoneal lavage

The video shows how to wash peritoneal cavity after injection of pASC-GFP particles to measure IL-1 β by ELISA.