Published in final edited form as:

Hepatology. 2017 February ; 65(2): 678–693. doi:10.1002/hep.28927.

Loss of the TGF- β Effector β 2SP Promotes Genomic Instability

Jian Chen^{1,*}, Vivek Shukla^{1,2,*}, Patrizia Farci³, Jaclyn Andricovich⁴, Wilma Jogunoori⁵, Lawrence N Kwong⁶, Lior H. Katz^{1,7}, Kirti Shetty⁸, Asif Rashid⁹, Xiaoping Su¹⁰, Jon White⁵, Lei Li¹¹, Alan Yaoqi Wang¹², Boris Blechacz¹, Gottumukkala S. Raju¹, Marta Davila¹, Bao-Ngoc Nguyen¹³, John R. Stroehlein¹, Junjie Chen¹¹, Sang Soo Kim¹⁴, Heather Levin¹³, Keigo Machida^{15,16}, Hidekazu Tsukamoto^{15,17,18}, Peter Michaely¹⁹, Alexandros Tzatsos⁴, Bibhuti Mishra^{5,13,20}, Richard Amdur^{13,20}, and Lopa Mishra^{5,13,20,†}

¹Department of Gastroenterology, Hepatology, and Nutrition, University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA

²Thoracic Oncology Section, Thoracic and Gastrointestinal Oncology Branch, National Cancer Institute, Bethesda, MD 20892, USA

³Hepatic Pathogenesis Section, Laboratory of Infectious Diseases, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD 20892, USA

⁴George Washington University, Department of Anatomy and Regenerative Biology, Washington, DC, 20052, USA

⁵Institute for Clinical Research, Veterans Affairs Medical Center, Washington, DC, 20422, USA

⁶Department of Translational Molecular Pathology, University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA

⁷Sheba Medical Center, Department of Gastroenterology, Tel Hashomer, 52621, Israel

⁸Division of Gastroenterology and Hepatology, Johns Hopkins University School of Medicine, Baltimore, MD 21287, USA

⁹Department of Pathology, University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA

¹⁰Department of Bioinformatics and Computational Biology, University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA

¹¹Department of Experimental Radiation Oncology, University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA

¹²Department of Cancer Biology, University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA

¹³Department of Surgery, George Washington University, Washington, DC, USA

¹⁴National Cancer Center, Radiation Medicine Branch, Goyang, 410-769, Korea

[†]Contact Information for Correspondence: Lopa Mishra, M.D., Director, Center for Translational Medicine, Professor, Department of Surgery, MFA, VA & George Washington University, 2300 Eye Street NW, Ross Hall #554, Washington, DC 20037, Tel: 202-401-2916, Fax: 202-462-2006, lopamishra2@gmail.com, lmishra@GWU.edu.

^{*}These authors contributed equally to this work

¹⁵Southern California Research Center for ALPD and Cirrhosis, University of Southern California, Los Angeles, CA 90089, USA

¹⁶Department of Molecular Microbiology and Immunology, Keck School of Medicine of the University of Southern California, Los Angeles, CA 90033 USA

¹⁷Department of Pathology, Keck School of Medicine of the University of Southern California, Los Angeles, CA 90089 USA

¹⁸Department of Veterans Affairs, Greater Los Angeles Healthcare System, Los Angeles, CA 90089, USA

¹⁹Department of Cell Biology, University of Texas Southwestern Medical Center, Dallas, Texas 75390, USA

²⁰Center for Translational Medicine, Department of Surgery, George Washington University, Washington DC 20037, USA

Abstract

Exposure to genotoxins such as ethanol-derived acetaldehyde leads to DNA damage and liver injury, and promotes the development of cancer. We report here a major role for the TGF- β /Smad3 adaptor β 2-Spectrin (β 2SP, gene *Sptbn1*) in maintaining genomic stability following alcohol-induced DNA damage. β 2SP supports DNA repair through β 2SP-dependent activation of Fancd2, a core component of the Fanconi anemia complex. Loss of β 2SP leads to decreased Fancd2 levels and sensitizes β 2SP mutants to DNA damage by ethanol treatment, leading to phenotypes that closely resemble those observed in animals lacking both *Aldh2* and *Fancd2*, and resemble human fetal alcohol syndrome. *Sptbn1*-deficient cells are hypersensitive to DNA cross-linking agents, and have defective DNA double-strand break repair that is rescued by ectopic Fancd2 expression. Moreover, *Fancd2* transcription in response to DNA damage/TGF- β stimulation is regulated by the β 2SP/Smad3 complex. Thus, dysfunctional TGF- β / β 2SP signaling impacts the processing of genotoxic metabolites by altering the Fanconi anemia DNA repair pathway.

Keywords

β 2-spectrin; liver cancer; genotoxicity; DNA repair; Fanconi anemia pathway

Introduction

Lifelong maintenance of genomic stability is essential for protection against the development of cancer. Alcohol-related genotoxicity, arising from DNA damage by metabolically-generated reactive aldehydes, has recently been observed in models with genetic inactivation of members of the Fanconi anemia pathway (1). *Fancd2*-mutant mice crossed with *Aldh2*-mutant mice are susceptible to ethanol teratogenicity and display defective DNA interstrand cross-link repair (1). Alcohol-treated *Fancd2*^{-/-} embryos, however, do not develop a fetal alcohol syndrome-like phenotype, implicating a critical protective role of *Aldh2* in acetaldehyde metabolism, and that this process is more complex than envisaged (2–5).

Transforming growth factor β (TGF- β) is critical for the regulation of several cancers (6) and functions as an extracellular sensor for ionizing radiation-induced cell damage. Keratinocytes isolated from TGF β 1-null mice exhibit marked genomic instability that accelerates tumor progression (7). Defective TGF- β signaling is implicated in multiple cancers owing to the frequent somatic mutations in, or deregulation of, its components, such as Smad3, Smad4, and TGF- β receptors 1 and 2 (TBR1 and TBR2). Smads are the intracellular mediators of TGF- β signaling, and their function is modulated by adaptor proteins such as the Smad anchor for receptor activation, filamin, microtubules, and β 2-spectrin (β 2SP) (8). Recent studies of Smad4-knockout mice, which develop head and neck cancers, demonstrate a significant role for Smad4 in promoting genomic stability through regulation of the Fanconi anemia/Breast Cancer (FANC/BRCA) DNA repair pathway (9). Despite these observations, precisely how the TGF- β pathway contributes to toxin-induced DNA damage repair remains unclear.

A crucial TGF- β /Smad3/4 adaptor and transcriptional co-factor, β 2-Spectrin (β 2SP, gene *Sptbn1*) confers multiple aspects of the tumor suppressor functions of Smad3, in hepatocellular and other cancer development (10–12). β 2-Spectrin is a dynamic intracellular non-pleckstrin homology (PH)-domain protein that belongs to a family of polypeptides implicated in conferring cell polarity. Through associated binding partners, such as ankyrin, spectrins target and stabilize membrane proteins, such as ion transporters, exchangers, and cell adhesion molecules in diverse tissues and cell types including erythrocytes, gut, liver, and brain cells (13). Spectrin dysfunction has previously been linked to abnormalities in mammalian physiology, including elliptocytosis, anemia, and cerebellar degeneration. More recently, spectrins have been linked to multiple signaling pathways, including cell cycle regulation, DNA repair, and TGF- β signaling (10, 14, 15). We have previously shown that heterozygous loss of *Sptbn1* in mice results in one of the more consistent genetic models of liver malignancies associated with loss of TGF- β signaling, with more than 40% of mice spontaneously developing liver tumors (8, 11, 16–18). β 2SP (*Sptbn1*) interacts with Smad3 on the cell nucleus in a TGF- β -dependent manner (12). The β 2SB/Smad3 complex then regulates transcriptional factors involved in stem-cell-associated liver tumorigenesis (12). Homozygous loss of *Sptbn1* results in mouse embryonic lethality due to multiple abnormalities of the liver, gut, and brain, demonstrating the essential nature of *Sptbn1* in embryogenesis (10).

Here, we report that more than half of *Sptbn1*-homozygous-null mouse embryos develop a spontaneous fetal alcohol syndrome-like phenotype, which becomes exacerbated upon exposure to alcohol. Analysis of *Sptbn1*-null cells indicates a marked defect in DNA repair in response to various exogenous mutagens including aldehyde. Importantly, the β 2SP/Smad3 complex directly regulates the transcription of *Fancd2*, as observed in both *in vivo* and *in vitro* systems. These findings are further discussed in the context of β 2SP-mediated TGF- β modulation, specifically the role of this signaling axis in the Fanconi anemia pathway for DNA damage repair, alcohol sensitivity, and liver tumorigenesis.

Experimental Procedures

Cell culture, transfection, and shRNA/siRNA-mediated silencing

Human liver cancer cell lines HepG2 (HB8065; ATCC, Rockville, MD) and SNU398 (CRL-2233, ATCC) were cultured in DME/F12 medium (D5671; Sigma Aldrich, St Louis, MO) supplemented with 10% fetal bovine serum (F2442; Sigma Aldrich). For transfection, *Sptbn1*^{-/-} MEFs were transfected with pDEST-SFB-FancD2, Myc-FancA, or Myc-FancM plasmids using Lipofectamine 2000 (Invitrogen, Carlsbad, CA). pDEST-SFB-FancD2, Myc-FancA, and Myc-FancM plasmids were kindly provided by Drs. Lei Li and Junjie Chen. TGF- β 1 (T1654; Sigma Aldrich) was added to a final concentration of 100 pM for cell culture. Lentiviral particles containing β 2SP (sc-36551) or control shRNA lentiviral particles (sc-108080) were used to infect HepG2 cells. shRNAs to β 2SP (sc-36551-v) and control shRNA (sc-108080) were purchased from Santa Cruz Biotechnology (Dallas, TX). siRNA for FANCD2 (L-016376-00-0005) and siRNA control Non-targeting Pool (# D-001810-10-05) were purchased from Dharmacon (Pittsburgh PA).

Animal exposure to ethanol

Matings of the male and female *Sptbn1*^{+/-} mice were set up, and the females were monitored daily for vaginal plugs. The day of vaginal plug detection was defined as E0.5. Pregnant mice were administered 5 g/kg 28% ethanol intraperitoneally from E6.5 to E12.5. The dose was divided into 2 separate injections of 2.5 g/kg 6 hours apart. On day E13.5, the mice were sacrificed and the fetuses were dissected for further analysis and genotyping (Table S1). For oral treatment with ethanol, 10- to 12-week-old mice of 2 genotypes (*Sptbn1*^{+/+}, n = 10; *Sptbn1*^{+/-}, n = 12) were exposed to continuous oral ethanol for 2 weeks (Supplementary Materials and Methods). Blood alcohol levels were determined using the ANALOX GM-7 Micro-Stat analyzer and its reagent (ANALOX Instruments USA, Lumburg, MA). All the experiments were approved by the Institutional Animal Care and Use Committee at MD Anderson Cancer Center, Houston, TX.

Lipid peroxidation assay

The concentrations of lipid peroxidation malondialdehyde (MDA) were measured in liver homogenates using kits from Abcom (ab118970, Cambridge, MA). Approximately 50 mg of liver tissue was homogenized in 300 μ L of MDA Lysis Buffer, and centrifuged at 13,000 $\times$ g for 10 minutes. Supernatants were mixed with Thiobarbituric Acid buffer, which was measured spectrophotometrically at 520 nm.

Homologous recombination (HR) repair analysis

U2OS cell clones stably expressing an HR repair reporter were kindly provided by Dr. Junjie Chen. These U2OS cells were infected with lentiviral shRNAs against luciferase or *Sptbn1* (β 2SP) to generate U2OS cells-shRNA- β 2SP stable cells. These cells were then transfected with mock or I-SceI plasmid along with MSCV-GFP as a control for transfection efficiency. After 2 days, the cells were assayed by flow cytometry to measure GFP-positive (HR-repaired) cells.

PCR for non-homologous end-joining (NHEJ) repair analysis

Genomic DNA was extracted from mock- or I-SceI-transfected DR-GFP stable expressed U2OS cells. PCR was then performed. Primers were as follows: DR-GFP-F, CTGCTAACCATGTTTCATGCC and DR-GFP-R, AAGTCGTGCTGCTTCATGTG. PCR products were subsequently digested with I-SceI or I-SceI + BclI and subjected to gel electrophoresis using 1.2% agarose gel.

Statistical analysis

Differences between 2 groups were evaluated using 2-tailed Student *t* tests. Results are expressed as mean $\pm$ standard deviation (SD) unless otherwise indicated. For multiple comparisons, 1-way ANOVA with post-hoc Bonferroni's test was used. Results are expressed as mean $\pm$ SD unless otherwise indicated. For all statistical analyses, $p < 0.05$ was considered statistically significant.

Results

Sptbn1-deficient mice exhibit a phenotype similar to human fetal alcohol syndrome and are sensitive to ethanol exposure

Using genetically engineered mouse models of *Sptbn1*, we found that no *Sptbn1*^{−/−} pups were born. Therefore, we examined E11.5 *Sptbn1*^{−/−} embryos and found not only a disruption in TGF- β signaling, but also a number of morphological abnormalities such as retarded growth, liver and gut hypoplasia, cardiac muscle hypertrophy, and neural defects (10). Later stage surviving *Sptbn1*^{−/−} embryos (52%, at E13.5–16.5) spontaneously develop exencephaly, edema, and anophthalmia, phenotypes that are also characteristic of human fetal alcohol syndrome (Fig. 1A–B and Fig. S1). Therefore, the role of β SP2 in the context of alcohol catabolism was further interrogated. Interestingly, a similar phenotype was recently described for double knockout *Aldh2*^{−/−}/*Fancd2*^{−/−} mice, in which the acetaldehyde-catabolizing enzyme aldehyde dehydrogenase 2 (*Aldh2*) was observed to be essential for embryonic viability of mice deficient in the Fanconi anemia complementation group D2 (*FancD2*) DNA repair gene (1).

Given the remarkable sensitivity of *Aldh2*^{−/−}/*Fancd2*^{−/−} mice to ethanol exposure, both *in utero* and postnatally, we asked whether *Sptbn1*-deficient mice are similarly ethanol sensitive. We treated *Sptbn1*^{+/−} females impregnated by *Sptbn1*^{+/−} males with a total dose of 5 g/kg 28% (v/v) ethanol twice per day by intraperitoneal injection from E6.5 to E12.5, and analyzed embryos at E13.5 (Fig. 1C). Fetal ablation occurred in 100% of the homozygote insufficient *Sptbn1*^{−/−} mutants from *Sptbn1*^{+/−} mothers treated with ethanol, whereas wild type *Sptbn1*^{+/+} offspring were unaffected by ethanol treatment (Fig. 1D–E and Table S1). A comparison of our results to those observed in *Aldh2*^{−/−}/*Fancd2*^{−/−} mice on the same C57BL/6 background reveals that both *Sptbn1*^{−/−} and *Aldh2*^{−/−}/*Fancd2*^{−/−} mice are markedly susceptible to ethanol toxicity and that both sets of mutants manifest similar morphological defects including exencephaly, anencephaly, and anophthalmia (Fig. S1D and Table S2).

Next, the effects of ethanol exposure were assessed in adult heterozygote *Sptbn1*^{+/-} mice by exposing six- to ten-week-old *Sptbn1*^{+/+} and *Sptbn1*^{+/-} mice to ethanol via their drinking water supply for two weeks—one week of 15% (v/v) followed by one week with 20% (v/v) ethanol/water (Fig. 1F). Blood alcohol levels in the ethanol-exposed mice ranged from 20 to 150 mg/dL, likely reflecting differing times of the last alcohol consumption prior to sacrifice (Fig. S2A). The hepatic concentration of malondialdehyde (MDA), the aldehydic product of lipid peroxidation, was also measured. Interestingly, increased MDA levels were only found in deficient *Sptbn1*^{+/-} mouse livers, and not wild type controls suggesting that *Sptbn1*^{+/-} mice are more susceptible to alcohol-induced oxidant damage compared to wild type mice (Fig. 1G). Moreover, we observed 3-fold increased levels of γ H2AX in mutant *Sptbn1*^{+/-} mouse livers after exposure to two weeks of alcohol (Fig. 1H).

Hematopoietic progenitor cells are known to be highly sensitive to ethanol-induced oxidant stress (1). Upon ethanol administration, these heterozygous mice presented a striking bone marrow aplasia, and blood analysis demonstrated dramatic decreases in hemoglobin content and platelet counts (Fig. S2B–C). Further analysis of the bone marrow of ethanol-treated *Sptbn1*^{+/-} mice revealed a marked reduction of nucleated cells, as well as the inability to form colonies of all hematopoietic lineages when plated on methylcellulose (Fig. S2D) (1, 19). Collectively, the data suggest that deficiency in TGF- β -signaling/ β 2SP predisposes mice to DNA damage associated with ethanol exposure, similar to the observations in *Aldh2/Fancd2* mutant mice.

***Sptbn1* deficiency leads to genomic instability and hypersensitivity to DNA damage**

In *Aldh2*^{-/-}/*Fancd2*^{-/-} mice, ethanol sensitivity has been attributed to an underlying genomic instability resulting from loss of Fancd2-dependent DNA repair activity. We therefore hypothesized that loss of β 2SP similarly hinders DNA repair after exposure to alcohol/genotoxic stress and possibly promotes genomic instability. To further investigate the role of β 2SP in genotoxic stress and genomic stability, *Sptbn1*^{-/-} mouse embryonic fibroblasts (MEFs) were irradiated to induce chromosomal aberrations. We found that *Sptbn1*^{-/-} MEFs exhibited a significant increase in chromatid breaks and complex radial structures compared to wild-type MEFs (Fig. 2A–C). The chromosomal instability of *Sptbn1*^{-/-} MEFs portrays a similarity to Fanconi anemia patients and animal models which are known to be hypersensitive to DNA cross-linking agents (20, 21). These data suggest a common, underlying defect in DNA damage repair. To test this hypothesis, *Sptbn1*^{-/-} MEFs were treated with the cross-linking agent mitomycin C (MMC) or with ionizing irradiation (IR) to induce DNA double strand breaks. Despite their distinct mechanisms of DNA damage, both treatments resulted in a greater dose-dependent decrease in mutant *Sptbn1*^{-/-} cell viability compared to wild-type (Fig. 2D–E). Furthermore, we used splenic B cells to test whether *Sptbn1*^{+/-} cells have decreased resistance to acetaldehyde. Indeed, the viability of B cells from *Sptbn1*^{+/-} mice significantly decreased after exposure to acetaldehyde in a dose-dependent manner when compared to wild type B cells (Fig. 2F). Consistently, a significant increase (2–3 fold) in chromosomal aberrations was observed in *Sptbn1*^{-/-} MEFs treated with MMC compared to untreated *Sptbn1*^{-/-} MEFs and *Sptbn1*^{+/+} MEFs with or without MMC treatment (Fig. 2G–H). The results suggest that *Sptbn1*-deficient cells exhibit

DNA repair defects across a range of insults including DNA cross-linking, DNA double strand breakage, and acetaldehyde adducts.

To determine if the observed hypersensitivity of mutant *Sptbn1*^{-/-} MEFs to DNA damage is a repercussion from a defective DNA repair, foci formation was assessed through microscopy using phospho-H2AX (γ H2AX), 53BP1, phospho-CDK2 (pCDK2), Rad51, and NBS1 as markers (22). We found that *Sptbn1*^{-/-} MEFs treated with 1 μ M MMC have increased residual γ H2AX (4 fold), 53BP1 (2.4 fold), and pCDK2 (3 fold) foci, and decreased Rad51 (4.5 fold) and NBS1 (3 fold) foci compared to wild type, indicating defective repair of DNA double-strand breaks (DSBs) by the homologous recombination pathway (Fig. 3A–E).

Loss of β 2SP results in a defect in homologous recombination (HR) DNA repair

To further assess the impact of *Sptbn1* deficiency on DNA damage repair pathways, we used a well-established homologous recombination repair system, DR-GFP, that is composed of two differentially-mutated green fluorescent protein (GFP) genes oriented as direct repeats: the upstream repeat contains the recognition site for the rare-cutting I-SceI endonuclease (SecGFP) and the downstream repeat is a truncated GFP fragment (iGFP) (Fig. 4A) (23–25). Briefly, transfection of DR-GFP-integrated U2OS cells with an I-SceI endonuclease expression vector can generate a site-specific DSB in the SceGFP coding region. If this DSB is repaired by HR, GFP expression is restored and GFP-positive cells can be quantitated by flow cytometry (Fig. 4A). We first knocked down β 2SP, Fancd2, or β 2SP/Fancd2 in DR-GFP-integrated U2OS cells (Fig. 4B, left panel). In the absence of I-SceI expression, very few GFP-positive cells were detected among these cells (Fig. 4B, right panel). After I-SceI expression, β 2SP- and/or Fancd2-knockdown cells exhibited ~3–4-fold decreases in the percentage of GFP-positive cells compared to control cells, indicative of defective HR repair (Fig. 4B, right panel). These results suggest that both β 2SP- and Fancd2-knockdown cells have impaired HR, indicating a role for both β 2SP and Fancd2 in this repair pathway. Interestingly, double knockdown of Fancd2 and β 2SP in U2OS cells resulted in further increased defective repair by HR, suggesting that while β 2SP is important for proper Fancd2 function, it is likely that other regulators are involved in this complex process. Next, a PCR-based method was used to further investigate both HR and non-homologous end joining (NHEJ) (Fig. 4C, left panel) (26). We used PCR primers that would amplify products of HR and NHEJ repair surrounding the I-SceI site in DR-GFP-integrated U2OS cells that had been transfected with I-SceI endonuclease. We then cleaved the PCR products with I-SceI to determine how much of the amplified fragment consisted of detectable DSB repair products. To specifically detect NHEJ products, we cleaved the PCR products with I-SceI and with BcgI, which cleaves only the homologous repair product (Fig. 4C, left panel). Without transfection of the I-SceI expression vector, the PCR product is completely cleaved by I-SceI (0.5kb) (Fig. 4C, right panel, first line). With transfection of the I-SceI expression vector in control cells, PCR product was not cleaved by I-SceI. The amount of uncleaved DNA (0.65kb) reflects an HR repair process, which results in I-SceI site loss (replacement of BcgI on I-SceI site). For β 2SP- and/or Fancd2-knockdown cells, the amount of uncleaved DNA (0.65kb) is markedly reduced (~5–10 fold), indicating that the chromosomal I-SceI site was not repaired by HR and was clearly cleaved by I-SceI (Fig. 4C, right panel, upper blot). To

differentiate NHEJ from homologous repair, PCR products were additionally treated with BclI, which as mentioned only cleaves HR products but not NHEJ products (Fig. 4C, right panel, lower blot). Thus, the amount of uncleaved DNA after digestion with both BclI and I-SceI indicates the level of NHEJ. In all of the cells, whether knockdown control or β 2SP and/or Fancd2 knockdown, the amount of uncleaved DNA is not altered, demonstrating that the level of repair of a chromosomal DSB by NHEJ is not reduced by either β 2SP or Fancd2 deficiency. These assays confirm that β 2SP knockdown reduces HR repair, but has no effect on NHEJ repair efficiency. Taken together, the results indicate that loss of β 2SP and/or Fancd2 results in fundamental repair defects by homologous recombination.

TGF- β / β 2SP/Smad3 upregulates Fancd2 transcriptionally

To gain further understanding of how TGF- β 1 impacts DNA repair, we extended previous studies in our mutant cells. TGF- β 1-induced Smad3 phosphorylation is inhibited in β 2SP-knockdown HepG2 cells, confirming prior studies that β 2SP is required for activation of the TGF- β pathway (Fig. S3) (10, 12). We observe that MMC activates TGF- β signaling, as evidenced by increased Smad3 phosphorylation. This TGF- β activating effect was reduced in mutant *Sptbn1*^{-/-} MEFs (Fig. 5A) and β 2SP-knockdown HepG2 cells (Fig. 5B), suggesting that β 2SP may play a critical role on MMC-activated TGF- β pathway.

The nuclear translocation of β 2SP in response to TGF- β , and β 2SP's known role as a transcriptional co-activator with Smad3/4 (12), led us to hypothesize that β 2SP may similarly be involved in transcriptional programs of the DNA damage repair response. To gain an unbiased view of transcriptomic changes caused by β 2SP perturbation, we analyzed whole-transcriptome RNA sequencing data of MEFs isolated from wild-type and *Sptbn1*^{-/-} embryos. *Fancd2* mRNA levels are significantly reduced in *Sptbn1*^{-/-} MEFs (Fig. S4A). Since *Fancd2* is involved in DNA interstrand cross-link repair, we validated the altered expression of Fanconi anemia genes by Q-RT-PCR and confirmed markedly suppressed *Fancd2* expression (4 fold) in mutant *Sptbn1*^{-/-} MEFs (Fig. S4B).

Previous reports have demonstrated that β 2SP interacts with Smad3 in the cell nucleus in a TGF- β -dependent manner. The β 2SP/Smad3 complex then transcriptionally upregulates its targets, such *TERT* and *CTCF* (12). To determine whether β 2SP and TGF- β facilitate DNA-damage-induced *Fancd2* expression, we compared *Fancd2* mRNA levels in wild type *Sptbn1*^{+/+} with mutant *Sptbn1*^{-/-} MEFs. Treatment with TGF- β 1 or MMC increased *Fancd2* mRNA levels in wild type *Sptbn1*^{+/+} MEFs but not in mutant *Sptbn1*^{-/-} MEFs, and simultaneous treatment with both agents produced an additive increase in *Fancd2* mRNA only in the wild type MEFs (Fig. 6A). Treatment with the TGF- β inhibitor SB431542 significantly inhibited MMC-, TGF- β -, or TGF- β /MMC-induced *Fancd2* mRNA levels (Fig. 6A). These results demonstrate that TGF- β / β 2SP is required for *Fancd2* levels to increase in response to MMC or TGF- β , suggesting that the β 2SP/Smad3 complex may directly regulate *Fancd2* transcription. To address this hypothesis, we examined *Fancd2* mRNA levels in HepG2 cells after Smad3 knockdown. Knockdown of Smad3 in HepG2 cells significantly prevented the TGF- β /MMC-induced *Fancd2* mRNA levels, indicating that Smad3 is required for *Fancd2* upregulation in response to MMC or TGF- β (Fig. 6B).

Smad3 transcriptionally regulates downstream target genes via the Smad binding element (SBE) motif, 5'-GTCT-3' (27). We examined human *Fancd2* and identified five potential SBE motifs within the -1,000 bp promoter proximal region. Of note, only two evolutionarily conserved sites, at ⁻³¹⁶GTCT-⁻³¹³ and ⁻⁵³⁰AGAC-⁻⁵²⁷, were found to be significantly enriched for β 2SP and Smad3 binding in HepG2 cells after treatment with TGF- β 1, suggesting a TGF- β -dependent mechanism (Fig. 6C). We then identified two evolutionarily-conserved SBE motifs within a 1,000 bp promoter region proximal to the mouse *Fancd2* gene. Examination of protein binding to the 2 elements using chromatin immunoprecipitation (ChIP) assays revealed significant enrichment of both β 2SP and Smad3 at one of the elements, the promoter proximal ⁻²⁹⁸AGAC-⁻²⁹⁵ motif. Importantly, Smad3 binding was completely lost in mutant *Sptbn1*^{-/-} MEFs, suggesting the β 2SP/Smad3 complex is necessary for *Fancd2* transcriptional regulation (Fig. 6D). We further demonstrated that knockdown of β 2SP with shRNA in HepG2 cells significantly reduced TGF- β /Smad3-induced *Fancd2* transcriptional activity (Fig. 6E). Mutation of either of these two conserved SBE motifs on the human *Fancd2* promoter compromised TGF- β /Smad3/ β 2SP-induced *Fancd2* transcription, while simultaneous mutation of both SBE motifs virtually abrogated transcription (Fig. 6F). Knockdown of Smad3 in HepG2 cells markedly suppressed TGF- β - or MMC-induced total FancD2 protein levels (Fig. S5A). Depletion of β 2SP with shRNA in HepG2 cells resulted in a significant reduction of TGF- β /Smad3-induced FancD2 expression levels (Fig. S5B), but did not alter FancD2 protein stability (Fig. S5C), thus confirming transcriptional control of *Fancd2* levels by TGF β / β 2SP/Smad3 (Fig. 6). Cellular defects in DNA damage repair observed in *Sptbn1*^{-/-} cells were rescued only by ectopic expression of FancD2, but not by FancM or FancA, suggesting the specific role of FancD2 transcription in *Sptbn1*-deficiency-induced DNA damage (Fig. 7). Overall, these data suggest that β 2SP, together with Smad3, is a critical regulator of DNA repair by mediation of *Fancd2* transcription.

β 2SP and FancD2 are clinically correlated

We investigated the clinical relevance of altered β 2SP and FancD2 function using immunohistochemical analyses of 20 human liver specimens from alcoholic hepatitis (n=5), alcoholic cirrhosis (n=5), and alcohol-associated liver cancer (n=5) patients, as well as normal controls (n=5). Increased nuclear localization of β 2SP was detected in patients with alcoholic hepatitis and cirrhosis, while nuclear expression of β 2SP and FancD2 was dramatically decreased in alcohol-associated liver tumor cells (Fig. 8A–B). Overall, nuclear β 2SP positively correlated with nuclear FancD2 (Fig. 8C), consistent with the hypothesis that β 2SP and FancD2 are activated in response to alcohol-induced liver injury. Furthermore, loss of this response in alcohol-associated liver cancer is consistent with a selection for events that contribute to genomic instability, though this remains to be proven. Finally, detailed analysis of a 91-case hepatitis C virus (HCV)-associated HCC cohort available from the Gene Expression Omnibus database (#GSE9843) was performed to characterize expression levels of *β 2SP* and *Fancd2* in HCC. Indeed, the mRNA expression of *β 2SP* and *Fancd2* are positively correlated in this independent HCC cohort (Fig. 8D). Recent studies have also demonstrated that *Sptbn1* gene expression positively correlates with *E-cadherin* and *kallistatin* gene expression in both HCV-associated HCC (Gene Expression Omnibus GSE6764) and hepatitis B virus (HBV)-associated HCC (Gene Expression Omnibus

GSE14520) (28). The data suggest that β 2SP functions as a tumor suppressor in HCC through multiple different signaling pathways. Moreover, it is possible that the TGF- β / β 2SP/Smad3/Fancd2 axis may also be directly activated in response to virus-induced DNA damage and genomic instability, but this remains to be investigated further. On the basis of our findings, we propose a model in which activation of the TGF- β / β 2SP/Smad3 pathway and the Fancd2 protein protects against toxin- or alcohol-induced DNA damage and promotes DNA repair. Alternatively, dysfunctional TGF- β signaling profoundly decreases protection against genotoxic metabolites by affecting the Fanconi anemia pathway.

Discussion

In this study, a novel role for the TGF- β pathway was uncovered in mediating genomic stability and homologous recombination through the Smad3 adaptor protein β 2SP, which is required for the transcriptional activation of a component of the Fanconi anemia pathway—Fancd2. Thus, TGF- β signaling is pivotal not only for regulation of hepatocyte proliferation and extracellular matrix deposition in the liver as previously reported (29–32), but also for maintenance of genomic stability in the context of toxin-induced DNA damage. Indeed, TGF- β is activated in response to tissue damage (33). Interestingly, only the double knockout *Fancd2/Aldh2* mice display fetal alcohol syndrome (FAS), indicating that elevated acetaldehyde was required for the FAS phenotype. When there is a deficiency of β 2SP, it is possible that the loss of the TGF- β pathway has additional sensitizing effects besides Fancd2 expression, thus rendering the mice more susceptible to DNA damage. Moreover, recent discoveries indicate that cytoskeletal proteins belonging to the spectrin/ankyrin family can regulate key signal transduction pathways by functioning as adaptor molecules (13).

The complex detoxifying function of the liver clearly exposes the organ to a significant degree of genotoxic stress; this stress must be overcome to prevent DNA damage, cirrhosis, and the onset of liver cancer. Our model proposes that, in response to toxins such as alcohol, the TGF- β signaling pathway upregulates expression of *Fancd2*, mediated by β 2SP and Smad3, to confer genomic stability and proper DNA repair. These studies highlight that further characterization of the TGF- β pathway in alcohol-induced injury could prove beneficial in therapies that target reactive aldehydes and DNA repair.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial Support

This work was supported by National Institute of Health grants R01AA023146 (L. Mishra), R01CA042857 (L. Mishra), P01CA130821 (L. Mishra), R01AA019392 (H. Tsukamoto, L. Mishra), Multidisciplinary Research Program Proposal (L. Mishra), and Science & Technology Acquisition and Retention Funding (L. Mishra).

We thank Drs. Walter Hittelman and KJ. Patel for critical review and helpful suggestions with the manuscript; Drs. Deborah Berry, Yun Seong Jeong, Liem Phan, Bibhuti Mishra, Bhaskar V.S. Kallakury, Xifeng Wu, Milind Javle, Premkumar Reddy, Nina M. Muñoz, Shareen J. Iqbal and Young Jin Gi, Suchin Kundra, Jiun-Sheng Chen for technical support. This work was supported by National Institute of Health grants R01AA023146 (L. Mishra), R01CA042857 (L. Mishra), P01CA130821 (L. Mishra), R01AA019392 (H. Tsukamoto, L. Mishra),

List of Abbreviations

β2SP	β2-Spectrin
ChIP	Chromatin immunoprecipitation
DSBs	Double-strand breaks
FAS	Fetal alcohol syndrome
GFP	Green fluorescent protein
HBV	Hepatitis B Virus
HCV	Hepatitis C Virus
HR	Homologous recombination
MEFs	Mouse Embryonic Fibroblasts
MDA	Malondialdehyde
MMC	Mitomycin C
NHEJ	non-homologous end-joining
SBE	Smad-binding element
TGF-β	Transforming growth factor β

References

1. Langevin F, Crossan GP, Rosado IV, Arends MJ, Patel KJ. Fancd2 counteracts the toxic effects of naturally produced aldehydes in mice. *Nature*. 2011; 475:53–58. [PubMed: 21734703]
2. Seitz HK, Stickel F. Molecular mechanisms of alcohol-mediated carcinogenesis. *Nat Rev Cancer*. 2007; 7:599–612. [PubMed: 17646865]
3. Moldovan GL, D'Andrea AD. To the rescue: the Fanconi anemia genome stability pathway salvages replication forks. *Cancer Cell*. 2012; 22:5–6. [PubMed: 22789533]
4. Joo W, Xu G, Persky NS, Smogorzewska A, Rudge DG, Buzovetsky O, Elledge SJ, et al. Structure of the FANCI-FANCD2 complex: insights into the Fanconi anemia DNA repair pathway. *Science*. 2011; 333:312–316. [PubMed: 21764741]
5. Oberbeck N, Langevin F, King G, de Wind N, Crossan GP, Patel KJ. Maternal aldehyde elimination during pregnancy preserves the fetal genome. *Mol Cell*. 2014; 55:807–817. [PubMed: 25155611]
6. Muraoka-Cook RS, Kurokawa H, Koh Y, Forbes JT, Roebuck LR, Barcellos-Hoff MH, Moody SE, et al. Conditional overexpression of active transforming growth factor beta1 in vivo accelerates metastases of transgenic mammary tumors. *Cancer Res*. 2004; 64:9002–9011. [PubMed: 15604265]
7. Glick A, Popescu N, Alexander V, Ueno H, Bottinger E, Yuspa SH. Defects in transforming growth factor-beta signaling cooperate with a Ras oncogene to cause rapid aneuploidy and malignant transformation of mouse keratinocytes. *Proc Natl Acad Sci U S A*. 1999; 96:14949–14954. [PubMed: 10611318]
8. Mishra L, Derynck R, Mishra B. Transforming growth factor-beta signaling in stem cells and cancer. *Science*. 2005; 310:68–71. [PubMed: 16210527]

9. Bornstein S, White R, Malkoski S, Oka M, Han G, Cleaver T, Reh D, et al. Smad4 loss in mice causes spontaneous head and neck cancer with increased genomic instability and inflammation. *J Clin Invest.* 2009; 119:3408–3419. [PubMed: 19841536]
10. Tang Y, Katuri V, Dillner A, Mishra B, Deng CX, Mishra L. Disruption of transforming growth factor-beta signaling in ELF beta-spectrin-deficient mice. *Science.* 2003; 299:574–577. [PubMed: 12543979]
11. Kitisin K, Ganesan N, Tang Y, Jogunoori W, Volpe EA, Kim SS, Katuri V, et al. Disruption of transforming growth factor-beta signaling through beta-spectrin ELF leads to hepatocellular cancer through cyclin D1 activation. *Oncogene.* 2007; 26:7103–7110. [PubMed: 17546056]
12. Chen J, Yao ZX, Chen JS, Gi YJ, Munoz NM, Kundra S, Herlong HF, et al. TGF-beta/beta2-spectrin/CTCF-regulated tumor suppression in human stem cell disorder Beckwith-Wiedemann syndrome. *J Clin Invest.* 2016; 126:527–542. [PubMed: 26784546]
13. Bennett V, Healy J. Organizing the fluid membrane bilayer: diseases linked to spectrin and ankyrin. *Trends Mol Med.* 2008; 14:28–36. [PubMed: 18083066]
14. Husi H, Ward MA, Choudhary JS, Blackstock WP, Grant SG. Proteomic analysis of NMDA receptor-adhesion protein signaling complexes. *Nat Neurosci.* 2000; 3:661–669. [PubMed: 10862698]
15. Sridharan D, Brown M, Lambert WC, McMahon LW, Lambert MW. Nonerythroid alphaII spectrin is required for recruitment of FANCA and XPF to nuclear foci induced by DNA interstrand cross-links. *J Cell Sci.* 2003; 116:823–835. [PubMed: 12571280]
16. Tang Y, Kitisin K, Jogunoori W, Li C, Deng CX, Mueller SC, Ransom HW, et al. Progenitor/stem cells give rise to liver cancer due to aberrant TGF-beta and IL-6 signaling. *Proc Natl Acad Sci U S A.* 2008; 105:2445–2450. [PubMed: 18263735]
17. Kanzler S, Meyer E, Lohse AW, Schirmacher P, Henninger J, Galle PR, Blessing M. Hepatocellular expression of a dominant-negative mutant TGF-beta type II receptor accelerates chemically induced hepatocarcinogenesis. *Oncogene.* 2001; 20:5015–5024. [PubMed: 11526486]
18. Romero-Gallo J, Sozmen EG, Chytil A, Russell WE, Whitehead R, Parks WT, Holdren MS, et al. Inactivation of TGF-beta signaling in hepatocytes results in an increased proliferative response after partial hepatectomy. *Oncogene.* 2005; 24:3028–3041. [PubMed: 15735717]
19. Smith C, Gasparetto M, Jordan C, Pollyea DA, Vasilou V. The effects of alcohol and aldehyde dehydrogenases on disorders of hematopoiesis. *Adv Exp Med Biol.* 2015; 815:349–359. [PubMed: 25427917]
20. Chen M, Tomkins DJ, Auerbach W, McKelvie C, Youssoufian H, Liu L, Gan O, et al. Inactivation of Fac in mice produces inducible chromosomal instability and reduced fertility reminiscent of Fanconi anaemia. *Nat Genet.* 1996; 12:448–451. [PubMed: 8630504]
21. Sasaki MS, Tonomura A. A high susceptibility of Fanconi's anemia to chromosome breakage by DNA cross-linking agents. *Cancer Res.* 1973; 33:1829–1836. [PubMed: 4352739]
22. Deans AJ, West SC. DNA interstrand crosslink repair and cancer. *Nat Rev Cancer.* 2011; 11:467–480. [PubMed: 21701511]
23. Peng G, Yim EK, Dai H, Jackson AP, Burgt I, Pan MR, Hu R, et al. BRIT1/MCPH1 links chromatin remodelling to DNA damage response. *Nat Cell Biol.* 2009; 11:865–872. [PubMed: 19525936]
24. Pierce AJ, Johnson RD, Thompson LH, Jasin M. XRCC3 promotes homology-directed repair of DNA damage in mammalian cells. *Genes Dev.* 1999; 13:2633–2638. [PubMed: 10541549]
25. Weinstock DM, Nakanishi K, Helgadottir HR, Jasin M. Assaying double-strand break repair pathway choice in mammalian cells using a targeted endonuclease or the RAG recombinase. *Methods Enzymol.* 2006; 409:524–540. [PubMed: 16793422]
26. Nakanishi K, Yang YG, Pierce AJ, Taniguchi T, Digweed M, D'Andrea AD, Wang ZQ, et al. Human Fanconi anemia monoubiquitination pathway promotes homologous DNA repair. *Proc Natl Acad Sci U S A.* 2005; 102:1110–1115. [PubMed: 15650050]
27. Massague J, Seoane J, Wotton D. Smad transcription factors. *Genes Dev.* 2005; 19:2783–2810. [PubMed: 16322555]

28. Zhi X, Lin L, Yang S, Bhuvaneshwar K, Wang H, Gusev Y, Lee MH, et al. betaII-Spectrin (SPTBN1) suppresses progression of hepatocellular carcinoma and Wnt signaling by regulation of Wnt inhibitor kallistatin. *Hepatology*. 2015; 61:598–612. [PubMed: 25307947]
29. Moses HL, Yang EY, Pietenpol JA. TGF-beta stimulation and inhibition of cell proliferation: new mechanistic insights. *Cell*. 1990; 63:245–247. [PubMed: 2208284]
30. Michalopoulos GK, DeFrances MC. Liver regeneration. *Science*. 1997; 276:60–66. [PubMed: 9082986]
31. Im YH, Kim HT, Kim IY, Factor VM, Hahm KB, Anzano M, Jang JJ, et al. Heterozygous mice for the transforming growth factor-beta type II receptor gene have increased susceptibility to hepatocellular carcinogenesis. *Cancer Res*. 2001; 61:6665–6668. [PubMed: 11559531]
32. Thenappan A, Shukla V, Abdul Khalek FJ, Li Y, Shetty K, Liu P, Li L, et al. Loss of transforming growth factor beta adaptor protein beta-2 spectrin leads to delayed liver regeneration in mice. *Hepatology*. 2011; 53:1641–1650. [PubMed: 21520177]
33. Ewan KB, Henshall-Powell RL, Ravani SA, Pajares MJ, Arteaga C, Warters R, Akhurst RJ, et al. Transforming growth factor-beta1 mediates cellular response to DNA damage in situ. *Cancer Res*. 2002; 62:5627–5631. [PubMed: 12384514]

Figure 1. *Sptbn1*^{-/-} mice exhibit a phenotype similar to human fetal alcohol syndrome and are markedly susceptible to ethanol toxicity

(A) Developmental abnormalities of fetal alcohol syndrome are observed in *Sptbn1*^{-/-} mutants at E13.5. Black arrowheads highlight microcephaly, exencephaly, anencephaly, edema, and anophthalmia. (B) *Sptbn1*-deficient embryos were observed during E13.5 to E16.5 at Mendelian ratio. Developmental abnormalities are found in *Sptbn1*^{-/-} mutants. χ^2 test relative to a Mendelian segregation, confidence interval 5%. *: Fisher's exact test, confidence interval 1%. (C) Diagrammatic representation of the experimental protocol to assess the effects of ethanol consumption in *Sptbn1*^{+/-} adult mice. (D) *In utero* exposure to ethanol-ablated *Sptbn1*^{-/-} homozygote fetuses. Scale bar, 2mm. (E) Quantification of developmental abnormalities observed in *Sptbn1*^{-/-} mice. Fisher's exact test was used, with a confidence interval of 1%. (F) Diagrammatic representation of the experimental protocol. *Sptbn1*^{+/+} and *Sptbn1*^{+/-} mice were exposed to continuous oral ethanol in their drinking water supply for 2 weeks. (G) Increased lipid peroxidation levels in mouse livers were

observed in alcohol-fed *Sptbn1*^{+/-} mice, compared to wild type mice. Hepatic lipid peroxidation levels were assessed by the concentration of MDA after 2 weeks oral ethanol drinking water administration. 5 mice for each group were used. (H) Increased expression levels of γ H2AX were observed in ethanol-fed *Sptbn1*^{+/-} mice livers. γ H2AX levels were detected by immunohistochemical analysis in mouse livers (n = 5 for each group). Scale bars, 25 μ m. Quantification of the immunohistochemical staining is shown. Error bars are shown as standard deviation (G and H). *: $p < 0.01$, Student *t*-test.

Figure 2. *Sptbn1*^{-/-} cells show genomic instability and sensitivity to DNA damaging agents as determined by clonogenic survival and metaphase chromosome aberrations analysis
 (A) Representative metaphase spreads from (i) *Sptbn1*^{+/+} and (ii) *Sptbn1*^{-/-} MEFs. Red arrows point to chromatid breaks and black arrows point to radial structures. (B) Quantification of aberrant radial figures from 200 metaphase spreads. (C) Quantification of aberrant chromatid breaks from 200 metaphase spreads. Means ± standard deviations of 3 independent experiments are shown in B and C. *: $p < 0.01$, Student t -test. (D, E) *Sptbn1*^{-/-} MEFs are significantly sensitive to MMC (D) and γ -irradiation (E) compared to *Sptbn1*^{+/+} MEFs. (F) *Sptbn1*^{+/-} mouse primary B cells are sensitive to exogenous acetaldehyde. Mouse primary B cells obtained from the spleens of *Sptbn1*^{+/+} and *Sptbn1*^{+/-} mice were stimulated with 40 μ g/ml lipopolysaccharide and treated with increasing concentrations of acetaldehyde, and cell viability was measured using trypan blue exclusion. For D-F, each point represents the mean ± standard deviation of the results obtained from 3 independent experiments, each carried out in triplicate. *: $p < 0.05$, one-way ANOVA. (G, H) Mutant

Sptbn1^{-/-} MEFs treated with MMC display significantly more chromosome breaks (G) and radial (H) per metaphase than *Sptbn1*^{+/+} MEFs. Means $\pm$ standard deviations of 3 independent experiments are shown. *: $p < 0.05$, 1-way ANOVA with post-hoc Bonferroni's test.

Figure 3. Impaired DNA damage response in absence of Sptbn1

(A to E) Mitomycin C (MMC) induced significantly more γ H2AX (A), 53BP1 (B), and pCDK2 (C) foci and fewer Rad51 (D) and NBS1 (E) foci in *Sptbn1*^{-/-} than *Sptbn1*^{+/+} MEFs. Foci were analyzed in five pairs of MEFs at passage 3 and exposed to 1 μ M MMC for different time points. Error bars represent standard deviations, 3 independent experiments were performed. *: $p < 0.01$, Student *t*-test.

Figure 4. β 2SP knockdown results in a defect in HR repair

(A) Diagrammatic representation of a HR repair system that is composed of two differentially-mutated GFP genes oriented as direct repeats: the upstream repeat contains the recognition site for the rare-cutting I-SceI endonuclease (SceGFP) and the downstream repeat is a truncated GFP fragment (iGFP). Cells were transfected with DR-GFP expression vector to generate a DR-GFP-integrated stable cell line. Transfection of DR-GFP-integrated U2OS cells with an I-SceI endonuclease expression vector generates a site-specific DSB in the SceGFP coding region. HR repairs DSB, and restored GFP-positive cells are detected and quantitated by flow cytometry. (B) β 2SP knockdown results in a defect in HR repair as measured by GFP-positive cells. DR-GFP-integrated U2OS cells were infected with

lentiviral shRNA-Ctrl, shRNA- β 2SP (left upper panel), siRNA-Ctrl, siRNA-Fancd2, or shRNA- β 2SP/siRNA-Fancd2 to generate knockdown cells (left lower panel). Then, the cells were transfected with mock or I-SceI plasmid (right panel) along with MSCV-GFP as an internal control for transfection efficiency. After 2 days of transfection, the cells were assayed by flow cytometry to measure GFP-positive (HR-repaired) cells. Means $\pm$ standard deviations of 3 independent experiments are shown. *: $p < 0.001$, 1-way ANOVA with post-hoc Bonferroni's test. (C) β 2SP knockdown does not affect NHEJ repair. A diagrammatic representation of the PCR-based method is shown (left panel). HR repair but NHEJ repair results in replacement of the I-SceI site with a BclI restriction site. PCR was performed using the primers F and R. The DR-GFP-integrated U2OS cells were infected with indicated lentiviral shRNAs, and then were transfected with mock or I-SceI plasmid. 48 hours post-transfection, genomic DNA was extracted for PCR amplification, followed by I-SceI or I-SceI plus BclI digestion. PCR products were subjected to gel electrophoresis (right panel).

Figure 5. MMC activates the TGF-β/Smad3 pathway
Sptbn1^{+/+} or *Sptbn1*^{-/-} MEFs (A) and HepG2-shRNA-Ctrl or shRNA-β2SP cells (B) were treated with 100 pM TGF-β1 for the indicated time points. Protein levels were detected by immunoblot analysis using indicated antibodies. Levels of Smad3 phosphorylation were quantitated by integrating optical density from Image J and are demonstrated as bar graphs. Total Smad3 was used for density normalization, and tubulin or actin were used for loading control.

Figure 6. β2SP/Smad3 complex transcriptionally regulates Fancd2 in a TGF-β-dependent manner

(A) Depletion of β2SP results in reduction of TGF-β- and/or MMC-induced *Fancd2* expression levels. mRNA expression of *Fancd2* was significantly increased (six fold) in *Sptbn1*^{+/+} MEFs after treatment with MMC and TGF-β, but not in *Sptbn1*^{-/-} MEFs, as determined by Q-RT-PCR. To block the TGF-β pathway, cells were pre-treated with 50 μM of TGFBR1 inhibitor SB431542 overnight. Cells were then exposed to 1 μM MMC and/or 100 pM TGF-β1 for 2 hours. *: $p < 0.001$, 1-way ANOVA with post-hoc Bonferroni's test. (B) Knockdown of Smad3 inhibits TGF-β- and/or MMC-induced *Fancd2* expression levels. mRNA expression of *Fancd2* was significantly decreased in HepG2-shRNA-Smad3 cells

compared to HepG2-shRNA-Ctrl cells after treatment with MMC and TGF- β . NS: not significant. *: $p < 0.001$, 1-way ANOVA with post-hoc Bonferroni's test. (C) Two conserved sites (-316 GTCT -313 and -530 AGAC -527) were significantly enriched for β 2SP/Smad3 binding on the human *Fancd2* promoter region. Diagrammatic representation shows 5 potential SBE motifs on the human *Fancd2* promoter region. Chromatin immunoprecipitation (ChIP) assays were performed to detect binding activity of Smad3 on the *Fancd2* promoter region. Genomic DNA was isolated from HepG2 cells treated with TGF- β 1 (100 pM). Enrichment of β 2SP and Smad3 was measured using Q-PCR. *: $p < 0.001$ versus indicated, 1-way ANOVA with post-hoc Bonferroni's test. Error bars indicate standard deviations. (D) Smad3 binding to *Fancd2* on its promoter region is completely lost in *Sptbn1* $^{-/-}$ MEFs. ChIP assays were performed. Diagrammatic representation shows 2 potential Smad binding element (SBE) motifs on the mouse *Fancd2* promoter region. Genomic DNA was isolated from *Sptbn1* $^{+/+}$ and *Sptbn1* $^{-/-}$ MEFs and treated with TGF- β 1 (100 pM). Enrichment of β 2SP and Smad3 was measured using Q-PCR. *: $p < 0.001$ versus indicated, 1-way ANOVA with post-hoc Bonferroni's test. (E) β 2SP knockdown results in reduction of TGF- β /Smad3-induced *Fancd2* transcriptional activity. Gaussia luciferase reporter plasmids containing a -1.3kb of *Fancd2* promoter region were transfected with indicated plasmids into HepG2 shRNA-Ctrl and HepG2 shRNA- β 2SP cells. Cells were treated with 200 pM TGF- β 1 for 2 hours. The secondary reporter, secreted Alkaline Phosphatase (SEAP), serves as an internal control. *: $p < 0.001$, 1-way ANOVA with post-hoc Bonferroni's test. **: $p < 0.01$, Student *t*-test. (F) Mutation of SBE motifs on *Fancd2* promoter compromises TGF- β /Smad3/ β 2SP-induced *Fancd2* transcriptional activity as determined by secreted Gaussia luciferase assays. HepG2 cells were co-transfected with indicated plasmids. Cells were treated with 200 pM TGF- β 1 for 2 hours. Mut SBE-A: GTCT $\rightarrow$ CATA; Mut SBE-B: AGAC $\rightarrow$ GCGT; Mut SBE-A+B: GTCT $\rightarrow$ CATA---AGAC $\rightarrow$ GCGT. NS: not significant. *: $p < 0.001$, 1-way ANOVA with post-hoc Bonferroni's test. Results presented are mean $\pm$ standard deviations of 3 independent experiments (A-F).

Figure 7. Ectopic expression of FancD2 rescues the DNA repair defect in *Sptbn1*-null cells
Sptbn1^{+/+} or *Sptbn1*^{-/-} MEFs were transfected with FancD2, FancA or FancM expression vector. 48 hours post-transfection, MEFs were exposed to 1μM MMC for different time points. Foci were determined by immunofluorescence analyses. Ectopic expression of FancD2, FancA or FancM were detected by immunoblot analyses (insets). Error bars represent standard deviations, 3 independent experiments were performed. *: $p < 0.01$, Student *t*-test.

Figure 8. β2SP and FancD2 are activated in response to alcohol-induced liver injury

(A) Immunohistochemical staining for β2SP, FancD2, Smad3 and phosph-Smad3 expression in 20 human tissue specimens of normal liver ($n = 5$), alcoholic hepatitis ($n = 5$), alcoholic cirrhosis ($n = 5$), and alcohol-associated hepatocellular carcinoma ($n = 5$). The arrows point to nuclear staining for β2SP, FancD2, Smad3 and phosph-Smad3. Magnification × 20; insets magnification × 60. Scale bar, 200 μm. (B) Quantification of β2SP and FancD2 nuclear staining. The numbers of positive β2SP and FancD2 nuclear staining hepatocytes were counted in 5 random different fields for each patient (>50 hepatocytes per field). Error bars are shown as standard deviation. NS: not significant; *: $p < 0.01$, Student's *t*-test. (C) Nuclear staining for β2SP and FancD2 in human tissue specimens is positively correlated. SigmaPlot was used to analyze correlative relationships. (D) mRNA expression of β2SP and FancD2 are positively correlated in HCC. The z cores were calculated using Nexus Expression v3.0 (BioDiscovery) on the basis of transcriptomics analysis in HCC patient

cohort #GSE9843 (n = 91) (Gene Expression Omnibus database). The correlation graph and test were done using GraphPad Prism v6.0e.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript