Differential efficacy profile of aldosterone receptor antagonists, depending on the type of chronic heart failure, whether with reduced or preserved left ventricular ejection fraction—results of a meta-analysis of randomized controlled trials

Renato De Vecchis¹, Carmelina Ariano²

¹Cardiology Unit, Presidio Sanitario Intermedio “Elena d’Aosta”, ASL Napoli 1 Centro, Napoli, Italy; ²Geriatric Department, Casa di Cura “S.Maria del Pozzo”, Somma Vesuviana (Napoli), Italy

Contributions: (I) Conception and design: R De Vecchis; (II) Administrative support: R De Vecchis; (III) Provision of study materials or patients: All authors; (IV) Collection and assembly of data: All authors; (V) Data analysis and interpretation: All authors; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors

Correspondence to: Dr Renato De Vecchis, via P.Gaurico 21, 80125 Napoli, Italy. Email: devechis.erre@virgilio.it; r.de.vecchis@alice.it.

Background: Because of renin-angiotensin-aldosterone system (RAAS) activation, the patients with chronic heart failure (CHF) manifest increased ventricular stress, with impaired left ventricular function, and a slowing down in systemic venous drainage. More importantly, a reduction of the patient's life expectancy has been proven in the case of RAAS overstimulation. For these reasons, huge efforts have been made to obtain molecules able to efficaciously antagonize the RAAS overstimulation, such as aldosterone receptor antagonists (ARAs). These drugs have been shown to improve clinical outcomes in patients with heart failure with reduced left ventricular ejection fraction (HFREF), but not in those with preserved left ventricular ejection fraction (HFpEF). In order to study this topic more deeply, we carried out a meta-analysis of selective and nonselective ARAs in HFREF and HFpEF.

Methods: Only randomized controlled trials (RCTs) were incorporated in our meta-analysis. Studies were included if they satisfied the following criteria: experimental groups included patients with CHF treated with ARAs in addition to the conventional therapy; control groups included patients with CHF receiving conventional therapy without ARAs. Outcomes of interest were all-cause mortality, cardiovascular hospitalizations, hyperkalemia, or gynecomastia.

Results: Overall, 15 RCTs including a total of 15,671 patients were eligible for inclusion in the meta-analysis. ARA use in patients with heart failure was associated with a significant reduction in adverse outcomes. Indeed, a significant reduced odds of all-cause death among CHF patients treated with ARAs compared to controls was found [odds ratio (OR) =0.79; 95% CI: 0.73–0.87]. Subgroup analysis based on the heart failure (HF) type revealed a statistically significant benefit as regards all-cause death for patients with HFREF (OR =0.77; 95% confidence interval (CI): 0.69–0.84), but not for those with HFpEF (OR =0.91; 95% CI: 0.76–1.1). Furthermore reduced odds of CV hospitalizations was detected in the entire group of CHF patients under treatment with ARAs (OR =0.73; 95% CI: 0.61–0.89) as well as among HFREF patients treated with ARAs, compared to controls (OR =0.66; 95% CI: 0.51–0.85). Hyperkalemia was significantly more frequent with ARA use. Besides, ARA use was shown to be associated with the occurrence of gynecomastia. In particular, selective ARAs didn't induce significant amounts of gynecomastia compared to controls (OR =0.74; 95% CI: 0.43–1.27), while nonselective ARAs did (OR =8.22; 95% CI: 4.9–13.81).

Conclusions: Based on this meta-analysis, ARAs should be systematically used in patients with HFREF, in whom these drugs proved to reduce all-cause mortality and hospitalizations from cardiac cause. Conversely, ARA usage in HFpEF patients is questionable since in this CHF setting no significant improvement in clinical endpoints has been demonstrated so far. New selective ARAs are devoid of the risk of gynecomastia,

while are similar to nonselective ARAs with regard to the efficacy profile as well as to the risk of eliciting hyperkalemia.

Keywords: Heart failure (HF); aldosterone receptor antagonists (ARAs); meta-analysis

Submitted Nov 23, 2016. Accepted for publication Jan 08, 2017.

doi: 10.21037/cdt.2017.03.04

View this article at: <http://dx.doi.org/10.21037/cdt.2017.03.04>

Introduction

Background

The renin-angiotensin-aldosterone system (RAAS) is activated during chronic heart failure (CHF) but this activation implies unfavorable changes of hemodynamics and cardiac work. In short, the pathophysiologic mechanisms underlying the overactivation of the RAAS share some elements with the pathophysiologic dynamics involved in the stimulation of antidiuretic hormone release from the posterior pituitary (1). In fact, a common trigger, as regards the stimulation of the two systems, is represented by the relative condition of underfilling of the arterial vascular compartment with a simultaneous increase in the volume of blood contained in the systemic venous compartment. Underfilling of the arterial circulation due to reduced cardiac output results in unloading of the arterial baroreceptors, which will increase non-osmotic vasopressin release and activate the sympathetic nervous system (SNS) and RAAS. This adrenergic surge promotes additional stimulation of the RAAS.

Aldosterone, produced by the zona glomerulosa of the adrenal cortex, promotes avid reabsorption of sodium and water at the level of the collecting duct in the distal nephron. In CHF, increased volume of extracellular fluids, elicited by aldosterone through the increase in sodium and water reabsorption in the distal nephron, would be aimed at retrieving an effective arterial blood volume, so as to potentiate the cardiac pump function through the Frank-Starling mechanism (2-5). However, retrieval of the pump effectiveness is not really achieved, because increased preload causes a deleterious aggravation of the cardiac work in the decompensated left ventricle, which entails a progressively increasing defaillance of the working myocardial fibers. So in the failing heart an increase in left ventricular end-diastolic pressure occurs as a result of the increase in circulating fluid volume, induced by aldosterone; however, this does not result in any increase in cardiac output but rather causes a state of hemodynamic

congestion (6), initially subclinical, then clinically evident (dyspnea on minimal efforts, peripheral edema, etc.).

Because of RAAS activation, the CHF patient manifests increased ventricular stress, with impaired left ventricular systolic function, and impaired systemic venous drainage, i.e., slowing down of systemic venous flow.

Thus, RAAS overactivation is mirrored by clinical worsening of heart failure with aggravation of dyspnea and edema; more importantly, a reduction of the patient's life expectancy has been proven in the case of RAAS overstimulation. For these reasons, huge efforts have been made to obtain molecules that have the property to efficaciously antagonize RAAS overstimulation. ACE inhibitors, angiotensin receptor blockers (ARBs), and aldosterone receptor antagonists (ARAs) are the three classes of drugs that have been proven to effectively counter the RAAS overstimulation typically found in CHF.

Several studies showed that despite maximal RAAS blockade with ACE inhibitors and/or ARBs, plasma aldosterone levels remain elevated in patients with heart failure (7-9). Thus ARAs were introduced in the therapy of CHF, following the demonstration of their protective effect on the failing heart. Indeed, several trials documented improved survival for patients treated with ARAs, compared with controls who received only beta-blockers, diuretics, and ACE inhibitors (10-12). The aldosterone receptor blockers include both nonselective (spironolactone, canrenone) and selective antagonists (eplerenone, finerenone). Eplerenone was synthesized through chemical modification of spironolactone in order to enhance binding of mineralocorticoid receptors while reducing off-target binding to progesterone or androgen receptors (13). Eplerenone is associated with lower rates of impotence, gynecomastia, or breast pain in comparison to spironolactone (11,12).

Aims

The aim of this meta-analysis of randomized controlled

trials (RCTs) was to verify the impact of ARAs on some hard endpoints (all-cause death and hospitalizations from cardiovascular cause), making a comparative evaluation of these outcomes in CHF patients with reduced left ventricular ejection fraction (HFREF) and in those with preserved left ventricular ejection fraction (HFpEF), respectively. The meta-analysis was then extended to the comparison between nonselective and selective ARAs, with regard to their respective effects on the above-mentioned clinical endpoints, and also as regards the respective side effects, such as hyperkalemia or gynecomastia.

Methods

Eligibility criteria

In our meta-analysis we considered exclusively RCTs. These studies were included if they met the following criteria: experimental groups included patients with CHF treated with ARAs in addition to the conventional therapy; control groups included patients with CHF receiving conventional therapy without ARAs. In addition, the studies selected for the meta-analysis only included patients older than 18 years. Animal experimental studies as well as case reports of ARA administration in patients with CHF were eliminated from the meta-analysis. Similarly, all studies not written in English, duplicated studies, non-randomized studies, review articles, editorials, and expert opinions were excluded.

Outcomes of interest were all-cause mortality, cardiovascular hospitalizations, hyperkalemia, or gynecomastia. The efficacy endpoints, i.e., mortality and hospitalizations, as well as the safety endpoints, namely episodes of hyperkalemia or occurrence of gynecomastia, were evaluated in both patients with heart failure with HFREF and those with HFpEF as compared to control groups. Moreover, a comparison was made between nonselective ARAs (e.g., spironolactone, canrenone) and selective ARAs (e.g., eplerenone or finerenone) with regard to efficacy outcomes as well as side effects.

Study selection

A systematic search using some related terms was conducted using the PubMed and Scopus electronic archives (from June 1995 to June 30 2016).

The study was performed according to the guidelines and recommendations expressed in the PRISMA (Preferred Reporting Items for Systematic reviews and Meta-

Analyses) (14) statement. Search terms included “ARAs”, “mineralocorticoid receptor antagonists”, “spironolactone”, “canrenone”, “eplerenone”, “all-cause mortality”, “hospitalization”, “hyperkalemia”, and “gynecomastia” variously combined by means of the Boolean operators AND and odds ratio (OR). Titles and abstracts of all identified citations were reviewed independently by the two authors (R.D.V. and C.C.) and any candidate study was selected for further screening of the full text. We extracted the following items from each study: population (type of heart failure, study size), intervention (ARA type), control (placebo, none, other), and outcomes (all-cause mortality, cardiovascular mortality, hospitalizations, hyperkalemia, and gynecomastia).

Statistical analysis

Statistical analysis was performed using Review Manager 5.0.4 software (available from Cochrane Collaboration at <http://www.cochrane.org>) and Stata version 10 (Stata Corp LP, College station, TX, USA). For all of the considered efficacy or safety endpoints, the effect size, i.e., the effect of therapy with ARAs plus conventional therapy (ACE inhibitors, beta-blockers, and loop diuretics) versus conventional therapy alone was presented as an OR with a 95% confidence interval (CI). We used a random effects model to pool results, and inverse variance as the weighting method. For quantifying statistical heterogeneity, the calculation of the I^2 statistic (I^2) was also used, to represent the percentage of variability due to between-study variability. We rated I^2 of less than 25%, 25–50%, and more than 50% as low, moderate, and high amounts of heterogeneity, respectively. Publication bias was assessed using Begg's funnel plot. Results were regarded as statistically significant if P was less than 0.05.

Results

One thousand seven hundred sixty citations were found; from these, in total, 36 studies were selected. Among them, 21 were eliminated after reading the article in full, due to incomplete or unavailable data, or because of ascertained inconsistency with our inclusion criteria. In particular, we excluded articles from the meta-analysis because of treatment in a non-HF setting (four studies), lack of relevant outcomes (13 studies), study duplication (three studies), and not an RCT design (one study; see *Figure 1*).

Overall, 15 RCTs comprising a total of 15,671 patients

Figure 1 Flow diagram for meta-analysis. Overview of process used to identify studies suitable for inclusion in the meta-analysis.

were eligible for inclusion in the meta-analysis. ARA use in patients with heart failure was associated with a significant reduction in adverse outcomes. Indeed, significantly reduced odds of all-cause death among CHF patients treated with ARAs compared to controls was found (OR =0.79; 95% CI: 0.73–0.87; I^2 =30.67 [moderate heterogeneity]; see *Figure 2*). Subgroup analysis based on the HF type—whether HFREF or HFpEF—revealed a statistically significant benefit for patients with HFREF (OR =0.77; 95% CI: 0.69–0.84), whereas a protective effect against all-cause death was not attained by ARAs in the HFpEF subset (OR =0.91; 95% CI: 0.76–1.1; see *Figure 3*). Moreover, use of either a nonselective or selective ARA yielded a similarly significant reduction in all-cause mortality (*Figure 4*).

Furthermore, as regards hospitalizations from cardiovascular cause (CV hosp), *Figures 5* and *6* illustrate that significantly reduced odds of CV hosp were detected in the entire group of CHF patients under treatment with ARAs (OR =0.73; 95% CI: 0.61–0.89) as well as among HFREF patients treated with ARAs, compared to controls (OR =0.66; 95% CI: 0.51–0.85). High heterogeneity was found as an accompanying feature (I^2 =64.20%). Our *a priori* subgroup analysis partially explained the heterogeneity within this outcome, as a significant reduction in CV hosp was found in the HFREF (*Figure 6*) and nonselective ARA subgroups (*Figure 7*), whereas reduction in CV hosp in the selective ARA subset did not reach statistical significance

(*Figure 7*). Hyperkalemia was significantly more common with ARA use (*Figure 8*).

In addition, subgroup analysis by ARA type documented that both nonselective and selective ARAs were similarly associated with increased odds of episodes of hyperkalemia compared to controls (*Figure 9*).

ARA use was shown to be associated with the occurrence of gynecomastia (*Figure 10*). In particular, selective ARAs proved not to produce significant amounts of gynecomastia compared to controls (OR =0.74; 95% CI: 0.43–1.27), while nonselective ARAs did (OR =8.22; 95% CI: 4.9–13.81; see *Figure 11*).

Discussion

In this meta-analysis we tried to assess the impact of ARAs on several efficacy and safety endpoints by maintaining distinct the outcomes detected in HFREF from those associated with HFpEF. Furthermore, for both efficacy and safety outcomes in the setting of CHF, separate meta-analyses were performed for RCTs centered on nonselective ARAs and for those which had investigated selective ARAs.

ARA use in patients with heart failure was associated with a significant reduction in all-cause death (OR =0.79; 95% CI: 0.73–0.87; *Figure 2*) and CV hosp (OR =0.73; 95% CI: 0.61–0.89; *Figure 5*). However, judging by our subgroup analysis, the favorable effects of ARAs on the efficacy endpoints were limited to HFREF; conversely, ARA-related reductions concerning all-cause mortality and CV hosp in HFpEF patients did not reach statistical significance (*Figures 3* and *6*, respectively). Both selective and nonselective ARAs increased the risk of hyperkalemia in a similar manner. Instead, gynecomastia was limited to nonselective ARAs; indeed, nonselective ARAs exhibited an OR of 8.22 (95% CI: 4.9–13.81), as opposed to selective ARAs which showed a non-significant OR of 0.74 (95% CI: 0.43–1.27; *Figure 11*).

Overall, the evidence supporting ARA use in HFREF was based on a large number of trials with significant effect sizes for reducing adverse cardiac events. In contrast, the evidence for ARA use in HFpEF was based on a relatively small number of trials, only one of which had a significant reduction in CV hosp but no significant effect on all-cause mortality (15). Finally, our conclusions supporting ARA use in HFREF are in keeping with current American Heart Association guidelines which recommend ARAs for patients with HFREF and NYHA class II–IV symptoms or following acute MI complicated by HF and LVEF ≤40% (16).

Figure 2 Forest plot of all-cause mortality with ARA use in HF. Eight trials reported all-cause mortality rates with ARA use in HF patients compared to controls. ARA, aldosterone receptor antagonist; HF, heart failure.

On the whole, no significant effects of ARAs on efficacy outcomes (all-cause mortality, CV hosp) concerning HFpEF were demonstrated by the present meta-analysis. Furthermore, ARAs were associated with a significantly increased risk of some harmful side-effects, such as hyperkalemia and gynecomastia. In this meta-analysis, selective ARAs proved to offer a slight benefit in terms of no significant gynecomastia while having equivalent reductions in adverse cardiac outcomes with respect to nonselective ARAs. Based on our meta-analysis, we would advise continued usage of ARAs in HFREF, where there is a significant reduction in adverse cardiac outcomes,

such as all-cause mortality and CV hosp. Instead, based on the present meta-analysis, in our opinion the choice of administering an ARA to patients with HFpEF is rather questionable, because in this type of CHF the ARAs-related reductions in adverse cardiovascular outcomes did not reach statistical significance.

Hyperkalemia during ARA therapy: a challenging issue

Our meta-analysis was not specifically aimed at evaluating the approach with ARAs in the setting of cardio-renal syndromes, i.e., the conditions characterized by the

Heterogeneity: $I^2=28.10\%$; $Q=6.954$, $P=0.224$

Heterogeneity: $I^2=0.00\%$; $Q=0.52$, $P=0.4705$

Figure 3 The forest plot of all-cause mortality with ARA use has been subdivided according to HF type. ARA, aldosterone receptor antagonist; HF, heart failure.

Figure 4 The forest plot of all-cause mortality has been subdivided according to ARA type (whether nonselective or selective). ARA, aldosterone receptor antagonist.

Figure 5 Forest plot of hospitalizations from cardiovascular cause with ARA use in HF. Ten trials reported cardiovascular hospitalization rates with ARA use in HF patients compared to controls. ARA, aldosterone receptor antagonist; HF, heart failure.

Heterogeneity: $I^2=72.88\%$; $Q=22.12$, $P=0.0011$

Heterogeneity: $I^2=19.57\%$; $Q=2.486$, $P=0.2884$

Figure 6 The forest plot of cardiovascular hospitalizations has been subdivided according to HF type. HF, heart failure.

Heterogeneity: $I^2=62.21\%$; $Q=15.879$, $P=0.0144$

Heterogeneity: $I^2=69.16\%$; $Q=6.484$, $P=0.039$

Figure 7 The forest plot of cardiovascular hospitalizations has been subdivided according to ARA type (whether nonselective or selective). ARA, aldosterone receptor antagonist.

Figure 8 Forest plot of hyperkalemia with ARA use in HF. Fifteen trials reported hyperkalemia rates with ARA use in HF patients compared to controls. ARA, aldosterone receptor antagonist; HF, heart failure.

Heterogeneity: $I^2=0.00\%$; $Q=5.907$, $P=0.8229$ Heterogeneity: $I^2=0.00\%$; $Q=2.3337$, $P=0.5053$

Figure 9 The forest plot of hyperkalemia has been subdivided according to ARA type (whether nonselective or selective). ARA, aldosterone receptor antagonist.

Figure 10 Forest plot of gynecomastia with ARA use in HF. Eight trials reported gynecomastia rates with ARA use in HF patients compared to controls. ARA, aldosterone receptor antagonist; HF, heart failure.

Heterogeneity: $I^2=0.00\%$; $Q=0.228$, $P=0.9987$

Heterogeneity: $I^2=0.00\%$; $Q=0.0129$, $P=0.9095$

Figure 11 The forest plot of gynecomastia has been subdivided according to ARA type (whether nonselective or selective). ARA, aldosterone receptor antagonist.

coexistence of cardiac and renal failure (17–19). However, some points would deserve more thorough awareness in the medical community. In particular, current rough epidemiological calculations (20) indicate that about a third of the general population aged >70 years is affected by a reduction in the estimated glomerular filtration rate (eGFR) of <60 mL/min/1.73 m² and therefore has a chronic kidney disease (CKD) by definition (21). However, a stable low eGFR in the elderly, when it is physiologically sufficient to satisfy the homeostatic demands, is not a disease per se and seldom progresses up to an advanced renal failure (21). Moreover, it would be appropriate to underline that a large proportion of patients with CHF and concomitant renal insufficiency may have a simple overlapping of the two dysfunctions, cardiac and renal, with no possibility of documenting a cardiorenal syndrome type 2.

Notably, the most important RCTs incorporated in our meta-analysis did not exclude CHF patients with concomitant mild- to moderate renal insufficiency. For instance, in the EMPHASIS-HF study by Zannad *et al.* (12) as well as in the TOPCAT study by Pitt *et al.* (15), only severe renal dysfunction was excluded, as opposed to mild-to-moderate renal insufficiency. Indeed, in the Methods of both studies it is affirmed that severe renal dysfunction (an eGFR of <30 mL/min/1.73 m² of body surface area or a serum creatinine ≥2.5 mg/dL) was an exclusion criterion. Instead, the recruitment of CHF patients with a lesser degree of renal dysfunction was not banned. So it is reasonable to affirm that the populations of these studies did not consist of “enclaves” isolated from the real world. In fact in these patients, the comorbidity “renal failure” (due to atherosclerotic ischemic renal disease, hypertensive or diabetic nephropathy, etc.) was adequately represented.

Nevertheless, an effective and safe dosing modulation concerning evidence—based therapies for CHF, i.e., beta-blockers, ACE-inhibitors, ARBs and ARAs, remains a challenging problem, which requires wise clinical choices, in addition to the adherence to few inclusion and exclusion criteria. In particular, with regard to the conditions in which ARA treatment has to be banned, an important one is avoiding ARA therapy in CHF patients with advanced renal insufficiency (GFR <30 mL/min/1.73 m²), in whom even the use of reduced doses of ACE-inhibitor, e.g., enalapril at a dose of 2.5–5 mg per day, cannot be regarded as a sufficient precaution to avert the risk of hyperkalemia. Thus, in advanced CKD as well as in any case of acute kidney injury superimposed on a condition of chronic cardiac decompensation, ARA use should be mandatorily

avoided in order to prevent the impending risk of dangerous episodes of hyperkalemia.

However, in CHF patients with only mild or moderate CKD [GFR calculated between 30 and 60 mL/min/1.73 m² by using the modified diet in renal disease (MDRD) equation (22)] the simultaneous use of various types of drugs blocking the RAAS—in particular, ACE-inhibitors plus ARAs or, alternatively, ARBs plus ARAs—should not be impeded but rather it should be adequately modulated and carefully monitored (17).

Proper patient selection, including patients with diminished LV ejection fraction and excluding ones with moderate CKD (creatinine level ≥2.5 mg/dL) or serum K⁺ >5 mEq/L, would help minimize potential life-threatening hyperkalemia (23).

Conclusions

In addition to already existing supportive data, our meta-analysis provides further evidence that ARAs should be systematically used in patients with HFREF, where they improve some hard clinical endpoints, such as all-cause mortality and hospitalizations from cardiac cause. Conversely, based on the present meta-analysis, ARA usage in HFpEF patients is questionable since in this CHF setting no significant improvement in clinical endpoints has been demonstrated so far, in the face of the well-known risks of hyperkalemia and/or gynecomastia that chronic ARA therapy entails. Furthermore, new selective ARAs are not burdened by significant risk of gynecomastia, while they are similar to nonselective ARAs with regard to the efficacy profile as well as to the risk of eliciting hyperkalemia.

Acknowledgements

None.

Footnote

Conflicts of Interest: The authors have no conflicts of interest to declare.

References

1. Masoumi A, Ortiz F, Radhakrishnan J, et al. Mineralocorticoid receptor antagonists as diuretics: Can congestive heart failure learn from liver failure? *Heart Fail Rev* 2015;20:283–90.

2. Schrier RW. Body fluid volume regulation in health and disease: a unifying hypothesis. *Ann Intern Med* 1990;113:155-9.
3. Schrier RW. Water and sodium retention in edematous disorders: role of vasopressin and aldosterone. *Am J Med* 2006;119:S47-53.
4. Schrier RW. Decreased effective blood volume in edematous disorders: what does this mean? *J Am Soc Nephrol* 2007;18:2028-31.
5. Schrier RW, Abraham WT. Hormones and hemodynamics in heart failure. *N Engl J Med* 1999;341:577-85.
6. Gheorghiade M, Filippatos G, De Luca L, et al. Congestion in acute heart failure syndromes: an essential target of evaluation and treatment. *Am J Med* 2006;119:S3-S10.
7. McKelvie RS, Yusuf S, Pericak D, et al. Comparison of candesartan, enalapril, and their combination in congestive heart failure: randomized evaluation of strategies for left ventricular dysfunction (RESOLVD) pilot study. The RESOLVD Pilot Study Investigators. *Circulation* 1999;100:1056-64.
8. Struthers A, Krum H, Williams GH. A comparison of the aldosterone-blocking agents eplerenone and spironolactone. *Clin Cardiol* 2008;31:153-8.
9. Pitt B. Aldosterone blockade in patients with chronic heart failure. *Cardiol Clin* 2008;26:15-21.
10. Pitt B, Zannad F, Remme WJ, et al. The effect of spironolactone on morbidity and mortality in patients with severe heart failure. Randomized Aldactone Evaluation Study Investigators. *N Engl J Med* 1999;341:709-17.
11. Pitt B, Remme W, Zannad F, et al. Eplerenone, a selective aldosterone blocker, in patients with left ventricular dysfunction after myocardial infarction. *N Engl J Med* 2003;348:1309-21.
12. Zannad F, McMurray JJ, Krum H, et al. Eplerenone in patients with systolic heart failure and mild symptoms. *N Engl J Med* 2011;364:11-21.
13. de Gasparo M, Joss U, Ramjoué HP, et al. Three new epoxy-spirolactone derivatives: characterization in vivo and in vitro. *J Pharmacol Exp Ther* 1987;240:650-6.
14. Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* 2009;339:b2535.
15. Pitt B, Pfeffer MA, Assmann SF, et al. Spironolactone for heart failure with preserved ejection fraction. *N Engl J Med* 2014;370:1383-92.
16. Yancy CW, Jessup M, Bozkurt B, et al. 2013 ACCF/AHA guideline for the management of heart failure: executive summary: a report of the American College of Cardiology Foundation/American Heart Association Task Force on practice guidelines. *Circulation* 2013;128:1810-52.
17. Ronco C, Haapio M, House AA, et al. Cardiorenal syndrome. *J Am Coll Cardiol* 2008;52:1527-39.
18. Boerrigter G, Hoher B, Lapp H. Changes in renal function in congestive heart failure. *Curr Heart Fail Rep* 2013;10:285-95.
19. De Vecchis R, Baldi C. Cardiorenal syndrome type 2: from diagnosis to optimal management. *Ther Clin Risk Manag* 2014;10:949-61.
20. National Kidney Foundation. K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. National Kidney Foundation. *Am J Kidney Dis* 2002;39:S1-266.
21. Winearls CG, Glassock RJ. Classification of chronic kidney disease in the elderly: pitfalls and errors. *Nephron Clin Pract* 2011;119 Suppl 1:c2-4.
22. Levey AS, Bosch JP, Lewis JB, et al. A more accurate method to estimate glomerular filtration rate from serum creatinine: a new prediction equation. Modification of Diet in Renal Disease Study Group. *Ann Intern Med* 1999;130:461-70.
23. Ko DT, Juurlink DN, Mamdani MM, et al. Appropriateness of spironolactone prescribing in heart failure patients: a population-based study. *J Card Fail* 2006;12:205-10.

Cite this article as: De Vecchis R, Ariano C. Differential efficacy profile of aldosterone receptor antagonists, depending on the type of chronic heart failure, whether with reduced or preserved left ventricular ejection fraction—results of a meta-analysis of randomized controlled trials. *Cardiovasc Diagn Ther* 2017;7(3):272-287. doi: 10.21037/cdt.2017.03.04