Users' guide to detecting misleading claims in research

Misleading claims may be symptom of even more serious flaws

EDITOR—Montori et al justifiably draw attention to misleading claims in published clinical trials.¹ This is a serious and common problem.^{2,3} However, I object to their radical proposal that only the methods and results sections should be read, while the remainder of the paper should be ignored.

The proposal is inconsistent with the title of their paper, "Users' guide to detecting misleading claims in clinical research reports."¹ How are these misleading claims to be identified if the sections containing them are omitted? Furthermore, anyone capable of critically appraising a trial solely on the basis of the methods and results is unlikely to be fooled by misleading claims in the discussion.

But, more importantly, the proposal would deprive the discerning reader of witnessing the conflict between the results and the unwarranted conclusions. If researchers are willing to disseminate misleading claims then their integrity is brought into question. But if so, then all aspects of the trial—including the methods and results sections—are also brought into question. Misleading claims should be identified and broadcast loudly for they signal doubts about the entire study.

Large scale randomised trials create the ideal conditions for data manipulation. Yet this is merely one of a multitude of problems stemming from a flawed method.³ The remedy? When reading a study, ignore everything except the number of patients recruited. If this is large proceed no further as there is little prospect of encountering any data of genuine benefit to patients.

James Penston *consultant physician*
Scunthorpe General Hospital, Scunthorpe, North
Lincolnshire DN15 7BH
James.Penston@nig.nhs.uk

Competing interests: None declared.

1 Montori VM, Jaeschke R, Schünemann HJ, Bhandari M, Brozek JA, Devereaux PJ, et al. User's guide to detecting misleading claims in clinical research reports. *BMJ* 2004;329:1093-6. (6 November.)

2 Wennberg R, Zimmermann C. The PROGRESS trial three years later: time for a balanced report of effectiveness. *BMJ* 2004;329:968-70. (23 October.)

3 Penston J. *Fiction and fantasy in medical research: the large-scale randomised trial*. London: London Press, 2003.

Discussion section helps redress the balance

EDITOR—Montori et al produce a users' guide to detecting misleading claims in reports of clinical research.¹ In the previous letter Penston raises objections to the proposal to read only the methods and results section and to bypass the discussion section, from a reader's point of view. The implications of this for some authors of clinical research papers is also unnerving.

Every research paper that is produced is imperfect. Authors are usually aware of the bias and pitfalls of their paper, some of which are inevitable. The discussion section gives them an opportunity to address these issues and explain their obligatory position.

Researchers have spent valuable time and resources on the work and are well informed about the methods and results. In contrast, readers who scan the methods and results section can easily overlook important information because critical appraisal of a clinical trial is a truly absorbing and daunting task.

Some research papers may well mislead readers towards believing the result, but whether just reading the methods and results warrants the number one slot in the users' guide is debatable.

S Anuradha *master of public health student*
Heath Park, Cardiff CF14 4XN
sanuradha@ntlworld.com

Competing interests: None declared.

1 Montori VM, Jaeschke R, Schünemann HJ, Bhandari M, Brozek JA, Devereaux PJ, et al. User's guide to detecting misleading claims in clinical research reports. *BMJ* 2004;329:1093-6. (6 November.)

Meta-analyses have led to misleading claims in the past

EDITOR—Montori et al cite the Cochrane albumin review in the table comparing two systematic reviews of albumin for fluid resuscitation.^{1,2} This review is an egregious example of a review that made a misleading claim—in this case a 6% additional mortality when albumin was compared with saline in a meta-analysis.

The users' guide of Montori et al is designed for assessing a single randomised controlled trial; when many randomised con-

trolled trials are amalgamated into a review, the compilers demand a degree of trust from those who use their review for guidance. In dealing with a complex diverse subject, such as critical illness, no ordinary reader, editor, or peer reviewer, will examine critically all the randomised controlled trials from which the review's authors derived their conclusions.

In the Cochrane albumin review, seven randomised controlled trials were adduced to support the mistaken contention that albumin, used in the treatment of hypovolaemia, would result in one additional death for every 17 patients compared with saline. I made criticisms of each of these randomised controlled trials, and the authors of the review did not answer any of them.³

Reviews such as this are usually assessed by the Centre for Research and Development, whose aim is to raise the general standard of reviews in the NHS. Its comment on this review was that Cochrane reviews are of a high standard and are not evaluated by one of its reviewers.⁴ It has taken six years for the SAFE study to disprove the erroneous allegation of the albumin review,⁵ and so far there has been no retraction by any member of the Cochrane Collaboration.

Peter J Horsey *honorary consultant anaesthetist*
Ashley, Stockbridge, Hampshire SO20 6RH
Horsey@lineone.net

Competing interests: None declared.

1 Montori VM, Jaeschke R, Schünemann HJ, Bhandari M, Brozek JA, Devereaux PJ, et al. User's guide to detecting misleading claims in clinical research reports. *BMJ* 2004;329:1093-6. (6 November.)

2 Cochrane Injuries Group Albumin Reviewers. Human albumin administration in critically ill patients: systematic review of randomised controlled trials. *BMJ* 1998;317:235-40.

3 Horsey P, Roberts I. Albumin and hypovolaemia. *Lancet* 2002;359:70-3.

4 NHS Centre for Reviews and Dissemination. Human albumin administration in critically ill patients: systematic reviews of randomised trials. Database of Abstracts of Reviews (DARE) document 124582, accessed 22 June 2002.

5 Saline versus albumin study evaluation. A comparison of saline and albumin for fluid resuscitation in the intensive care unit. The SAFE Study Investigators. *N Engl J Med* 2004;350:2247-56.

bmj.com

Letters appearing here are an edited selection of rapid responses originally posted on bmj.com

We ask for all letters to the editor to be submitted as rapid responses via bmj.com

For advice see: bmj.com/rapidresponses

Authors' reply

EDITOR—The tone and scope of the rapid responses to our users' guides confirm the central assumption of our guides: that there are emotional and economic interests that affect the objectivity of science. We think that these guides will help readers form their own opinions about the evidence.

Penston pointed out an inconsistency between our title and our first guide—how can one detect misleading claims if one is not exposed to the discussion section, where those claims are often made? Perhaps our original title (Guides for clinicians to defend themselves from misleading claims) was more appropriate. Others have seen in our guides a validation of their concerns about specific studies and scientists' behaviours and justification for a broader evaluation of research protocols in ethics boards.

It seems that the process that our guides are intended to foster—frank dialogue on more effective ways for decision makers to deal with data that comes wrapped in conflict—is ongoing.

Victor M Montori *assistant professor*
McMaster University, Hamilton, ON, Canada
Department of Medicine

Roman Jaeschke *clinical professor of medicine*
Mayo Clinic College of Medicine, Rochester, MN
55905, USA
montori.victor@mayo.edu

Holger J Schünemann *associate professor of medicine*
Department of Clinical Epidemiology and
Biostatistics, McMaster University

The four other authors of this letter are Mohit Bhandari, assistant professor of medicine, P J Devereaux, assistant professor of medicine, Gordon H Guyatt, professor of medicine, all of Department of Clinical Epidemiology and Biostatistics, McMaster University; and Jan L Brozek, lecturer in medicine, Polish Institute for Evidence Based Medicine, Jagiellonian University Medical School, Krakow, Poland.

Competing interests: VMM, RJ, HJS, PJD, and GHG are associate editors of the *ACP Journal Club* and *Evidence-Based Medicine*. JLB and RJ edit an evidence based medicine journal in Poland. MB is editor of the evidence based orthopaedic trauma section in the *Journal of Orthopaedic Trauma*.

Issues in reporting epidemiological studies

Can't we all agree on confidence intervals?

EDITOR—Pocock et al discuss many valuable caveats regarding published epidemiological analyses.¹ The authors, however, perpetuate two misconceptions.

The first is that post hoc power calculations help the reader interpret already collected data. Power calculations are a guide to study design—namely, to consider what the investigators might observe if they do the study. Once they collect the data, the confidence interval contains all the relevant information because it is based on what the investigators actually observed.²

The second misconception is that the P value contains useful information. In almost

all cases the confidence interval (along with the point estimate) is the better choice. But not, as many aver, to see whether it crosses some arbitrary null value, such as a relative risk of 1. Rather readers should ask whether the confidence interval excludes important effects. For example, if a reduction of cardiovascular mortality of at least 5% is meaningful, check to see if the upper limit of the confidence interval is less than 0.95.

Matthew W Gillman *associate professor*
Department of Ambulatory Care and Prevention,
Harvard Medical School, Boston, MA 02215, USA
Matthew_gillman@hms.harvard.edu
Competing interests: None declared.

- 1 Pocock SJ, Collier TJ, Dandreo KJ, de Stavola BL, Goldman MB, Kalish LA, et al. Issues in the reporting of epidemiological studies: a survey of recent practice. *BMJ* 2004;329:883-98. (16 October.)
- 2 Goodman SN, Berlin JA. The use of predicted confidence intervals when planning experiments and the misuse of power when interpreting results. *Ann Intern Med* 1994;121:200-6 [erratum: *Ann Intern Med* 1995;122:478].

No data are given to support generalisation

EDITOR—The article by Pocock et al outlined deficiencies in epidemiological research.¹ Undoubtedly, the deficiencies they reported will be generalised to all observational studies, even though they specifically excluded papers examining treatments or patients with diseases. Even the accompanying editorial revisits the hormone replacement therapy debacle.² Interestingly, a paper about hormone replacement therapy would not have been eligible for this review as studies using pharmacological treatments were excluded. However, the authors dedicate only two paragraphs to the methods section so this fact may not be obvious at first glance, or not at all if the abstract is reviewed in isolation.

Before this study is used as another reason to discredit all observational research, its limitations should be appreciated. The authors say that clinical epidemiology in people with disease and studies in pharmacoeconomics raise similar issues,¹ but they provide no data to support this generalisation. On the basis of their inclusion and exclusion criteria, they would have examined studies such as those linking coffee and diabetes.³ I am neither surprised nor concerned that papers such as this may contain some residual confounding, and I think it is a reasonable expectation that healthcare professionals interpret these reports appropriately. Often all that is required is to read beyond the abstract.

David F Blackburn *assistant professor*
College of Pharmacy and Nutrition, University of
Saskatchewan, Saskatoon, SK, Canada S7N 5C9
d.blackburn@usask.ca

Competing interests: None declared.

- 1 Pocock SJ, Collier TJ, Dandreo KJ, de Stavola BL, Goldman MB, Kalish LA, et al. Issues in the reporting of epidemiological studies: a survey of recent practice. *BMJ* 2004;329:883-98. (16 October.)
- 2 Von Elm E, Egger M. The scandal of poor epidemiological research. *BMJ* 2004;329:868-9. (16 October.)
- 3 Salazar-Martinez E, Willett WC, Ascherio A, Manson JE, Leitzmann MF, Stampfer MJ, et al. Coffee consumption and risk for type 2 diabetes mellitus. *Ann Intern Med* 2004;140:1-8.

Study raises several concerns

EDITOR—Pocock et al surveyed some recent epidemiological publications.¹ We agree with the general idea that the practice of reporting in epidemiologic publications needs improvement, but we are concerned about their methodological approach and conclusions.

Firstly, they say that few studies gave any power calculations to justify the study sizes. Ironically, Pocock et al's survey does not give any justification for the chosen study size of 73 epidemiological papers.

Secondly, Pocock et al's opinion on studies too small for meaningful analysis can be seen from a different point of view. Studies that are statistically underpowered are obviously less well planned and are often disregarded because they more often suffer from type I and type II errors compared with larger studies. However, from a public health perspective small studies may indicate true increased risks for exposures of interest that may have a substantial population-wide effect if the exposure is prevalent. These studies could be regarded as early warnings.

Thirdly, we are generally concerned about the discussion section. Pocock et al nearly completely omitted any critical discussion of their own results. It is not our purpose to persuade the reader that every paper published in 2001 eligible with respect to the study by Pocock et al is appropriate from a methodological point of view. However, we are convinced that this study adds little to our (those working in the field) knowledge and to science in general. The study lacks a clear hypothesis and design, has an insufficient study size in view of the high "bibliometric" variability, and is prone to misinterpretation for those with little knowledge of epidemiology.

Karl-Heinz Jöckel *professor of medical informatics, biometry and epidemiology*
Institute of Medical Informatics, Biometry and
Epidemiology, University Hospital, University of
Duisburg-Essen, Hufelandstrasse 55, D-45147
Essen, Germany
k-h.joeckel@uni-essen.de

Andreas Stang *professor of clinical epidemiology*
Institute of Medical Epidemiology, Biometry and
Informatics, Medical Faculty,
Martin-Luther-University of Halle-Wittenberg,
Magdeburger Strasse 27, D-06097 Halle, Germany
andreas.stang@medizin.uni-halle.de

Competing interests: None declared.

- 1 Pocock SJ, Collier TJ, Dandreo KJ, de Stavola BL, Goldman MB, Kalish LA, et al. Issues in the reporting of epidemiological studies: a survey of recent practice. *BMJ* 2004;329:883-98. (16 October.)

Authors' reply

EDITOR—We agree with Gillman that estimates and their confidence intervals contain the key information when reporting results. However, our observation that many epidemiological studies lack statistical power is still highly pertinent.

The relevance of P values in epidemiologic publications will continue to be debated. Our view is that they are of value if used sparingly and interpreted wisely.

We are surprised that Blackburn questions our premise that studies in clinical epidemiology in people with disease and pharmacoepidemiology are liable to have similar reporting concerns. After all, they use the same types of study designs and analysis methods. We would welcome another survey in those fields, and are not optimistic that consistent high quality would shine through.

Regarding issues of residual confounding and other biases, we do not share Blackburn's view that healthcare professionals at large are sufficiently astute in critical appraisal of published epidemiology.

We accept the suggestion of Jöckel and Stang that a larger survey would be desirable. Nevertheless, we think that one month's survey does provide a sufficiently informative sample of epidemiological publications to make useful observations on current reporting practices.

We were deliberately not studying a specific epidemiological hypothesis. Rather, we explored several well recognised features of epidemiological research and assessed how they were tackled in current publications. We believe that our recommendations have a sound basis and are a valuable stepping stone in raising reporting standards in epidemiology.

Like any professional group, epidemiologists may not be universally receptive to constructive critical appraisal of their research output, but we are hopeful that guidance from STROBE (Standards for the Reporting of Observational Studies in Epidemiology) currently under development (www.strobe-statement.org) will have a wide acceptance in enhancing the quality of publications.

Stuart J Pocock *professor*
Stuart.Pocock@lshtm.ac.uk

Timothy J Collier *lecturer*
Department of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, London WC1E 7HT

Kimberley J Dandreo *epidemiologist*
New England Research Institutes, 9 Galen Street, Watertown, MA 02472, USA

The five other authors of this letter are Bianca L de Stavola, reader, and Valerie A McCormack, lecturer, Department of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine; Marlene B Goldman, associate professor, Department of Obstetrics and Gynaecology, Harvard Medical School and Beth Israel Deaconess Medical Centre, Boston, MA 02215, USA; Leslie A Kalish, associate professor, Clinical Research Program, Children's Hospital, Boston, MA 02115, USA; Linda E Kasten, senior biostatistician, Prometrika LLC, Cambridge, MA 02138, USA.

Competing interests: None declared.

HIV in injecting drug users in Asian countries

Naltrexone implants are set to supersede harm minimisation

EDITOR—Wodak et al draw attention to intravenous drug use as a source of HIV transmission.¹ They make the case for harm

minimisation strongly but omit to mention that this is consistent with ongoing and even increasing drug use.² Harm minimisation remains a hypothesis that has never been shown to fulfil Hill's principles of causation³; neither have simultaneous confounding factors been controlled for.

Such advocacy overlooks discordant data that include epidemic levels of the non-condompreventable human papillomavirus (including the commonest tumour in the developing world, cervical cancer); epidemic levels of herpes and rising rates of chlamydia; holoendemic hepatitis C among drug users; the unpopularity of methadone both among patients who take it and who don't; the long term effects of methadone including mortality of 50%, and 590 deaths implicating it in Australia 1997-2001; severe erosive dental disease; osteoporosis and osteopenia rates of 87%; the high prevalence of methadone use among major heroin dealers; and the failure rate of the "exchange" component of needle programmes with 10 times more needles given away annually than we have Australians. In three sub-Saharan nations HIV has decreased despite non-adherence to harm minimisation (J Museveni, address, Common Ground conference, Medical Institute of Sexual Health, Washington, DC, June 2004; www.medinstitute.org/Museveni.htm).

Contrariwise naltrexone has been called "psychiatry's most effective drug." Poor compliance has now been overcome by the development of long lasting subcutaneous depot preparations including that from Perth, Western Australia, which lasts up to 11 months.⁴ Surely the HIV dilemma necessitates regulatory fast tracking of this device, and the implant's emergent dissemination across the developing world?

Albert S Reece *general practitioner*
Brisbane, QLD 4101, Australia
sreece@bigpond.net.au

- 1 Wodak A, Ali R, Farrell M. HIV in injecting drug users in Asian countries. *BMJ* 2004;329:697-8. (25 September.)
- 2 Wodak A. Harm reduction: Australia as a case study. *Bull N Y Acad Med* 1995;P339.
- 3 Hill AB. The environment and disease: association or causation? *Proc R Soc Med* 1965; 58:295-300.
- 4 Kim TW, Alford DP, Malabanan AO, Samet JH. Bone health in methadone maintenance treatment. College of the Problems of Drug Dependence meeting, Puerto Rico, June 2004. Abstracts booklet 279-95.
- 5 Hulse GK, Arnold-Reed DE, O'Neil G, Chan CT, Hansson RC. Achieving long term continuous blood naltrexone and 6-beta-naltrexol coverage following sequential naltrexone implants. *Addict Biol* 2004;9:67-72.

Authors' reply

EDITOR—Reece concedes the critical importance of controlling the epidemic of HIV among and from injecting drug users, especially in Asia. This is a useful start. However, his subsequent comments are less than helpful.

Reece argues that the high incidence of hepatitis C indicates that harm minimisation is ineffective. A recent study estimated that needle syringe programmes between 1988 and 2000 avoided 25 000 cases of HIV and 21 000 cases of hepatitis C in injecting drug users in Australia.¹ How does Reece explain the excellent control of HIV in injecting

drug users achieved in Australia, which explicitly accepted harm minimisation and vigorously implemented harm minimisation programmes, whereas high rates of HIV infection occurred in the United States where harm reduction was explicitly rejected?

Reece claims that methadone treatment is unsafe: yet the risk of death is three to four times less for patients continuing in treatment than in those who discontinue treatment.² His claim that harm minimisation lacks public support is irreconcilable with official survey data. A solid majority (59%) of respondents to the 2001 national drug strategy household survey supported needle syringe programmes,³ representing an 18% increase in support from the 1998 survey,⁴ whereas 58% of respondents in the 1998 survey supported or strongly supported methadone compared with 19% opposed or strongly opposed.⁴

Reece argues that naltrexone implants are effective and safe. It is conventional to assume that all new treatments are ineffective and unsafe until data are available from multiple, rigorous, randomised controlled trials with reasonable durations of follow up. No exception should be made for naltrexone implants.

Alex Wodak *director*
Alcohol and Drug Service, St Vincent's Hospital, Darlinghurst, NSW 2010, Australia
awodak@stvincents.com.au

Robert Ali *director of clinical policy and research*
University of Adelaide
robert.ali@adelaide.edu.au

M Farrell *senior lecturer in addictions*
PO Box 48, National Addiction Centre, Institute of Psychiatry, London SE5 8AF
m.farrell@iop.kcl.ac.uk

Competing interests: None declared.

- 1 Commonwealth Department of Health and Ageing. *Return on investment in needle and syringe programs in Australia*. 2002. www.drugpolicy.org/docUploads/ROIfinal.pdf (accessed 10 Nov 2004).
- 2 Bell J, Zador D. A risk-benefit analysis of methadone maintenance treatment. *Drug Saf* 2000;22:179-90.
- 3 National drug strategy household survey: first results. Canberra: Australian Institute of Health and Welfare, 2001.
- 4 National drug strategy household survey: first results. Canberra: Australian Institute of Health and Welfare, 1998.

Paternal age and schizophrenia

One swallow does not make a summer

EDITOR—Sipos et al set themselves the task of showing the presence of a higher than average paternal age in schizophrenia and that this phenomenon is restricted only to cases with a low family history for the disease.¹ Both hypotheses are confirmed with statistical significance in their study.

This is interesting, but the sample studied is biased, which casts serious doubts on the validity of the results. The study includes only subjects with an age at onset between 16 and 38. The literature on schizophrenia shows that to generalise the

results obtained from a limited age sample is dangerous. Often differences that seem highly significant in age biased samples even up when the cohorts are followed up throughout their lifetime. The aetiological factors in schizophrenia are probably the same at all ages qualitatively but seem to vary greatly quantitatively.²

The male to female ratio in the incidence and prevalence of schizophrenia is a cautionary tale. The incidence is significantly higher in men than in women in age groups up to the mid-30s, but the ratio of men to women becomes inverted with age, reaching 1:2 for onset after 40 years of age and 1:6 after 60. Eventually the lifetime morbid risk is the same in both sexes.³

The results of the study by Sipos et al are therefore intriguing but must be interpreted cautiously. No stratification can correct the above bias. Only the follow up of this cohort for at least a further 15 years can offer reliable results.

Marco Procopio *consultant psychiatrist*
Priory Hospital Hove, Hove, East Sussex BN3 4FH
marcoprocopio00@hotmail.com

Competing interests: None declared.

- 1 Sipos A, Rasmussen F, Harrison G, Tynelius P, Lewis G, Leon DA, Gunnell D. Paternal age and schizophrenia: a population based cohort study. *BMJ* 2004;329:1070-3. (6 November.)
- 2 Kendler KS, Karkowski-Shuman L, Walsh D. Age at onset in schizophrenia and risk of illness in relatives. *Br J Psychiatry* 1996;169:213-8.
- 3 Jablenski A, Sartorius N, Ernberg G, Anker M, Korten A, Cooper JE, et al. Schizophrenia: manifestations, incidence and course in different cultures. A World Health Organisation 10-country study. *Psychol Med* 1992;20(suppl):1-97.

Authors' reply

EDITOR—Procopio highlights the fact that our study included only subjects with a younger age of onset ranging from 16 to 28 and suggests that our finding of an association between paternal age and schizophrenia therefore needs to be interpreted cautiously.

This association has already been replicated in a number of studies covering much wider ranges of age of onset: the Jerusalem cohort covered ages of onset until 33,¹ the case control study using the Stockholm inpatient register had ages of onset until 34,² and Zammit et al showed such an association in their cohort with ages of onset up to 45.³ A large Danish case-control study of 7704 patients found no evidence for any interaction with age of onset.⁴

We also investigated possible age of onset effects by dividing our cohort according to the length of the follow up period for each individual and found no marked difference in the hazard ratios per 10 year increase in paternal age for those with less than nine years of follow up (hazard ratio 1.51, 95% confidence interval 1.25 to 1.82) and those with more than nine years of follow up (hazard ratio 1.77, 1.10 to 2.86).

One swallow doesn't make a summer, but this association has now been reported in several studies, in different settings, and

over a wide range of ages. It seems to be a robust finding that may shed some light on the aetiology of schizophrenia.

Attila Sipos *honorary senior clinical lecturer in psychiatry*
Attila.Sipos@bristol.ac.uk

Glynn Harrison *professor of mental health*
Academic Unit of Psychiatry, University of Bristol, Bristol BS6 6JL

Finn Rasmussen *senior clinical lecturer and associate professor of epidemiology*
Department of Public Health Sciences, Karolinska Institute, Norrbacka, SE-17176 Stockholm, Sweden

The four other authors of this letter are Per Tynelius, senior statistician, Department of Public Health Sciences, Karolinska Institute; Glyn Lewis, professor of psychiatric epidemiology, Academic Unit of Psychiatry, University of Bristol; David A Leon, professor of epidemiology, Department of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, London WC1E 7HT; and David Gunnell, professor of epidemiology, Department of Social Medicine, University of Bristol, Bristol BS8 2PR.

Competing interests: None declared.

- 1 Malaspina D, Harlap S, Fennig S, Heiman D, Nahon D, Feldman D, et al. Advancing paternal age and the risk of schizophrenia. *Arch Gen Psychiatry* 2001;58:361-7.
- 2 Dalman C, Allebeck P. Paternal age and schizophrenia: further support for an association. *Am J Psychiatry* 2002;159:1591-2.
- 3 Zammit S, Allebeck P, Dalman C, Lundberg I, Hemmingson T, Owen MJ, et al. Paternal age and risk for schizophrenia. *Br J Psychiatry* 2003;183:405-8.
- 4 Byrne M, Agerbo E, Ewald H, Eaton WW, Mortensen PB. Parental age and risk of schizophrenia: a case-control study. *Arch Gen Psychiatry* 2003;60:673-8.

Self monitoring of blood pressure at home

Informed self regulation of drug treatment could be next step

EDITOR—While highlighting that home blood pressure readings can be lower and more reliable than those taken in the clinic, Stergiou et al seem reluctant to recommend extensive use of the practice or take the next step of empowering patients to take action on the basis of these results.¹ However, patients adjust their drug treatment, or even stop taking it altogether, without consulting a doctor or looking at their latest blood pressure readings. An estimated 50% of drugs for long term conditions are not taken as prescribed, and patients' beliefs about drugs are the biggest predictor of compliance, the necessity of the drugs being weighed against concerns about side effects and dependency.²

Research to date has focused on self monitoring, with comparatively few studies investigating the potential for informed self regulation of treatment. However, small scale studies on self monitoring and self medication in hypertension and anticoagulation management already show that this is an avenue to pursue.^{3,4} A recent Cochrane review reported better lung function outcomes in patients who adjusted their drug treatment by using a written plan than in those whose treatment was adjusted by a doctor.⁵

The Medicines Partnership is currently developing a pilot with Hillingdon Hospital and Hillingdon Primary Care Trust to test the approach in a UK setting, including patients with diabetes who have high blood pressure. This approach may be particularly suitable for patients with diabetes, who are already accustomed to self management and adjustment of insulin or oral treatment on the basis of self testing. This approach will not be for everybody, a regimen must not be imposed that will add to somebody's anxiety. However, self regulation on the basis of home monitoring could be an effective option when control is critical but current methods are not always effective.

Caroline L Kelham *project manager*
Medicines Partnership, London SE1 7JN
ckelham@medicines-partnership.org

Competing interests: None declared.

- 1 Stergiou G, Megden T, Padfield PL, Parati G, O'Brien E, Working Group on Blood Pressure Monitoring of the European Society of Hypertension. Self monitoring of blood pressure at home. *BMJ* 2004;329:870-1. (16 October.)
- 2 Horne R, Weinman J. Patients' beliefs about prescribed medicines and their role in adherence to treatment in chronic illness. *J Psychosom Res* 1999;47:555-67.
- 3 Zarnke KB, Feagan BG, Mahon JL, Feldman RD. A randomized study comparing a patient-directed hypertension management strategy with usual office-based care. *Am J Hypertens* 1997;10:58-67.
- 4 Murray E, Fitzmaurice D, McCahon D, Fuller C, Sandhur H. Training for patients in a randomised controlled trial of self management of warfarin treatment. *BMJ* 2004;328:437-8.
- 5 Gibson PG, Powell H, Coughlan J, Wilson AJ, Abramson M, Haywood P, et al. Self-management education and regular practitioner review for adults with asthma. *Cochrane Database Syst Rev* 2003;(1):CD001117.

Primary care professionals remain sceptical

EDITOR—The review by Stergiou et al of home blood pressure monitoring sets out recommendations that are similar to those published some years ago as a result of reviewing the literature and a study confirming the feasibility and acceptability of the procedure.¹⁻³

The problem is not in deciding what should be done but in getting recommendations implemented. Primary care, where most British patients with high blood pressure are managed exclusively, remains sceptical about this procedure, although the use of semiautomatic devices is now commonplace. Now, in my dotage, I am on the receiving end of the sphygmomanometer, and I sadly find that my carers are only interested in isolated office readings.

Malcolm Aylett *retired general practitioner*
Bath BA2 5EJ
Malcolm.aylett@smrl.demon.co.uk

Competing interests: None declared.

- 1 Stergiou G, Megden T, Padfield PL, Parati G, O'Brien E, Working Group on Blood Pressure Monitoring of the European Society of Hypertension. Self monitoring of blood pressure at home. *BMJ* 2004;329:870-1. (16 October.)
- 2 Aylett M. Pressure for change: unresolved issues in blood pressure measurement. *Br J Gen Pract* 1999;49:136-9.
- 3 Aylett MJ. The use of home blood pressure measurements to diagnose white coat hypertension in general practice. *J Hum Hypertens* 1996;10:17-20.