safety equipment at baseline. Consequently, the absolute differences between treatment arms in the percentages of families with safety practices were relatively small, and a greater difference in safety practices may be required to affect the occurrence of injury. It is also possible that safety practices are not associated with reduced injury rates, but this is less plausible as several observational studies have shown a lower risk of injury among people with a range of safety practices.

Several explanations are possible for the higher attendance rate in primary care among intervention arm children. Firstly, participation in the intervention arm of the trial may have raised parents' awareness and changed their consulting behaviour for more minor injuries. Secondly, risk compensation⁸ may have occurred.

Implications for injury prevention practice and future research

Our findings in relation to safety practices and degrees of satisfaction are encouraging. However, our findings also highlight the importance of rigorously evaluating the widespread provision of equipment not only in terms of safety practices but also in terms of injury outcomes and uptake of schemes by those most at risk. Very large trials are required, or well designed case-control studies or cohort studies to examine the protective effect of specific items of equipment, followed by randomised controlled trials.

We thank the families who took part in the study and the health visitors, nursery nurses, and doctors who supported the study. We are also indebted to Sarah Ross for tracking trial participants and identifying new births.

Contributors: See bmj.com

Funding: NHS Executive, Trent. DK was funded by a Department of Health Public Health Career Scientist Award. AW was funded by a Department of Health Primary Care Researcher Development Award.

Competing interests: None declared.

Ethical approval: Queen's Medical Centre Research Ethics Committee.

- 1 Towner E, Dowswell T, Mackereth C, Jarvis S. *What works in preventing unintentional injuries in children and young adolescents? An updated systematic review*. London: Health Development Agency, 2001.
- 2 Lyons R, Sander L, Weightman A, Patterson J, Jones SL, Rolfe B, et al. Modification of the home environment for the reduction of injuries. *Cochrane Database Syst Rev* 2003;4:CD003600.
- 3 DiGuseppi C, Roberts IG. Individual-level injury prevention strategies in the clinical setting. *Future Child* 2000;10:53-82.
- 4 DiGuseppi C, Higgins JP. Systematic review of controlled trials of interventions to promote smoke alarms. *Arch Dis Child* 2000;82:341-8.
- 5 King WJ, Klassen TP, LeBlanc J, Bernard-Bonnin AC, Robitaille Y, Pham B, et al. The effectiveness of a home visit to prevent childhood injury. *Pediatrics* 2001;108:382-8.
- 6 Kendrick D, Marsh P, Fielding K, Miller P. Preventing injuries in children: cluster randomised controlled trial in primary care. *BMJ* 1999;318:980-3.
- 7 Woolf AD, Saperstein A, Forjuoh S. Poisoning prevention knowledge and practices of parents after a childhood poisoning incident. *Pediatrics* 1992;90:867-70.
- 8 Hedlund J. Risky business: safety regulations, risk compensation, and individual behavior. *Injury Prev* 2000;6:82-9.

(Accepted 9 November 2004)

doi 10.1136/bmj.38309.664444.8F

Complicity

After qualification and the obligatory two house jobs, I had been called up for national service in the Royal Army Medical Corps. In 1958 I was in Cyprus with a small team at a medical centre and living in the mess of an infantry battalion. It was August, very hot, and Eoka snipers were shooting at us; not at me personally, but on occasion I had to administer to those who had been shot or peppered with shrapnel from a pencil bomb. Apart from this, life was pleasant: work in the morning, a very strong gin and tonic at lunchtime (the duty-free gin being very cheap and the tonic extremely expensive), a siesta, and an evening of conversation and developing friendship with the men who were actually out there being shot at.

Sometimes the afternoon was interrupted by a telephone call prefaced by a bizarre military codeword; one I remember was "flying duck." There were two types of call: one involved a long drive in the Landrover (with armed escort) to some remote hill village, there to decide whether an aged parent was sufficiently near to death that a visit from an imprisoned Eoka suspect son was warranted. This was almost impossible to determine, so the answer was invariably yes.

Far more disturbing was the call to the nearby police station. There I found four or five burly policemen, who regarded me with bored and slightly hostile contempt, and two very frightened young Cypriots. The question was whether they were fit for "further questioning." They had a number of injuries, mainly

bruising but some apparent joint damage. These were, I was told, a result of "resisting arrest." Standing in a bare room in rather threatening surroundings, my examination was not even adequate. I persuaded myself that one might have a broken rib and the other a shoulder injury and that both required x ray examination at the hospital.

It was not heroic. Yes, I was intimidated, but also the camaraderie of the previous three weeks under threat as against the clear suggestion of physical abuse made me ambivalent about both police and prisoners.

It was, as these things go, a trivial incident, but when I hear of medical complicity in torture I know how easy it would be to be drawn in and how difficult then to get free.

John Dewhurst *retired pharmaceutical physician and former principal in general practice, Finchampstead*

We welcome articles up to 600 words on topics such as *A memorable patient*, *A paper that changed my practice*, *My most unfortunate mistake*, or any other piece conveying instruction, pathos, or humour. Please submit the article on <http://submit.bmj.com> Permission is needed from the patient or a relative if an identifiable patient is referred to. We also welcome contributions for "Endpieces," consisting of quotations of up to 80 words (but most are considerably shorter) from any source, ancient or modern, which have appealed to the reader.