Published in final edited form as:

J Int Neuropsychol Soc. 2017 March ; 23(3): 214–222. doi:10.1017/S1355617716001089.

The Effect of Retrieval Practice on Short-Term and Long-Term Retention in HIV+ Individuals

G. Avci¹, S.P. Woods^{1,2}, M. Verduzco³, D.P. Sheppard¹, J.F. Sumowski^{4,5}, N. Chiaravalloti^{4,5}, J. DeLuca^{4,5}, and The HIV Neurobehavioral Research Program (HNRP) Group

¹Department of Psychology, University of Houston, TX, USA

²School of Psychology, University of Western Australia, Perth, AUSTRALIA

³Joint Doctoral Program in Clinical Psychology, San Diego State University, University of California–San Diego, San Diego, CA, USA

⁴Neuropsychology and Neuroscience, Kessler Foundation, West Orange, NJ, USA

⁵Rutgers, New Jersey Medical School, Newark, NJ, USA

Abstract

Objective—Episodic memory deficits are both common and impactful among persons infected with HIV; however, we know little about how to improve such deficits in the laboratory or in real life. Retrieval practice, by which retrieval of newly learned material improves subsequent recall more than simple restudy, is a robust memory boosting strategy that is effective in both healthy and clinical populations. In this study we investigated the benefits of retrieval practice in 52 people living with HIV and 21 seronegatives.

Method—In a within-subjects design, all participants studied 48 verbal paired associates in 3 learning conditions: Massed-Restudy, Spaced-Restudy, and Spaced-Testing. Retention of verbal paired associates was assessed after short- (30 minutes) and long- (30 days) delay intervals.

Results—After a short delay, both HIV+ persons and seronegatives benefited from retrieval practice more so than massed and spaced restudy. The same pattern of results was observed specifically for HIV+ persons with clinical levels of memory impairment. The long-term retention interval data evidenced a floor effect that precluded further analysis.

Conclusions—This study provides evidence that retrieval practice improves verbal episodic memory more than some other mnemonic strategies among HIV+ persons.

Keywords

Cognition; Learning; Memory; Episodic; Memory Disorders; Neuropsychological Tests; Rehabilitation

Topic Areas

Learning & Memory; Rehabilitation; Cognitive & Other

Introduction

Episodic memory impairment is quite common among HIV+ persons, with most estimates ranging between 40 and 60% (e.g., Heaton et al., 1995; Rippeth et al., 2004). The etiology of the episodic memory impairment in HIV is likely multifactorial, likely reflecting the adverse effects of the virus on frontostriatal circuits (Ellis, Langford, & Masliah, 2007; Ernst & Chang, 2004) as well as psychiatric (e.g., substance use disorders) and medical (e.g., hepatitis C co-infection) (Antinori et al., 2007) comorbidities. Of clinical relevance, HIV+ persons with episodic memory deficits are at heightened risk for experiencing problems with a variety of different everyday activities, including health behaviors. For example, Hinkin et al. (2004) reported that impairment in memory was associated with poor adherence to combination anti-retroviral therapy (cART). In that study, approximately two-thirds of HIV+ participants with memory impairment were classified as poor adherers. Moreover memory functioning in HIV+ persons has been identified as an independent predictor of real-world tasks such as return to work (van Gorp et al., 2006) and household shopping (Heaton et al., 2004). Thus, research to date clearly shows that episodic memory deficits are both common and impactful among HIV+ persons. However, we know much less about how to improve or restore memory deficits in this complex clinical population (see Weber et al., 2013 for a review).

One potentially viable approach is to translate promising findings from the rich cognitive psychology literature in healthy older adults into effective memory rehabilitation techniques for neuropsychological populations. This approach has been successfully implemented with people living with HIV. For example, Weber et al. (2011) tested the self-generation effect (i.e., information generated by an individual is learned more effectively than material that is presented didactically) in HIV+ persons via paired associate learning task. Results revealed that HIV+ persons with clinical memory impairment benefitted more from self-generation as opposed to simply reading the materials. Thus, it may be possible to improve memory functions in persons living with HIV disease using approaches that have shown success in basic cognitive psychology experiments.

The testing effect, also known as retrieval practice effect, is well-established memory strategy that produces particularly robust memory improvements in both healthy (see Roediger & Butler, 2011 for a review) and clinical populations (see Balota, Duchek, & Logan, 2007) and thus may be another effective means of improving memory in HIV+ persons. The testing effect is a cognitive phenomenon by which the act of retrieval of newly learned material improves subsequent recall more than other learning strategies such as massed study and spaced restudy (Carpenter & DeLosh, 2005; Roediger & Karpicke, 2006).

Both practical and theoretical reasons exist for examining the testing effect in adults with HIV. It has been reported that the testing effect has improved memory performance in clinical groups (for reviews see Dunlosky, Rawson, Marsh, Nathan, & Willingham, 2013;

Middleton & Schwartz, 2012). In fact, the effect has been observed in a variety of neuropsychological populations including multiple sclerosis (e.g., Basso, Lowery, Ghormley, Combs, & Johnson, 2006; Chiaravalloti & DeLuca, 2002; Sumowski et al., 2010a; 2013), traumatic brain injury (e.g., O'Brien, Chiaravalloti, Arango-Lasprilla, Lengenfelder, & DeLuca, 2007; Sumowski et al., 2010b), and mild dementia in early Alzheimer's disease (e.g., Lipinska et al., 1994). For example, Sumowski et al. (2010a) in a study with healthy adults and persons with multiple sclerosis (MS) found a large main effect of learning condition on memory ($\eta^2 = .54$) revealing large mnemonic advantages of tests for both groups. In fact, testing was the best learning condition for 90% of the participants (MS: 91% and healthy adults: 88%). Practically, if we are to encourage and expand the use of effective and simple manipulations such as the testing effect, then we need to be confident that the approach works in a variety of clinical groups with various types of memory deficits (e.g., HIV+ persons). Indeed, the cognitive rehabilitation literature shows that the relative magnitude of benefit from memory manipulations can vary across different clinical populations who may have different memory profiles (i.e., strengths and weaknesses). As such, from a scientific perspective, it is important to do the groundwork of demonstrating that the basic testing effect can be observed in HIV+ persons under controlled laboratory conditions before beginning the process of integrating the manipulation into a more comprehensive rehabilitation program.

Theoretical motivation for examining the testing effect in HIV-infected persons is more complex than the practical motivation given the lack of consensus among researchers regarding the factors that best explain this robust cognitive phenomenon. The testing effect has been explained in terms of changes in semantic representations through elaboration or the restriction of the search-set to relevant associations, effortful retrieval processes that become easier with repetition, and the potentiation of subsequent encoding (Carpenter, 2011; Karpicke, Lehman, & Aue, 2014; Pyc & Rawson, 2010; Roediger & Butler, 2011). Thus, several cognitive theories exist explaining the nature of the testing effect and its essential processes.

In this study, we do not directly test any of these theories; rather we seek to apply the testing effect in an effort to bolster memory performance among HIV+ persons. We argue that the testing effect will benefit HIV+ persons due to the nature of the neurocognitive impairment that is commonly observed in this population. It is well established that HIV preferentially (but not exclusively) impacts frontostriatal circuits (Ellis et al., 2007; Ernst & Chang, 2004) and that HIV+ persons show a corresponding pattern of impairment that includes slowed information processing efficiency, executive dysfunction, and deficient episodic memory encoding and retrieval (e.g., Gongvatana et al., 2007; Murji et al., 2003; Woods et al., 2006). For example, episodic memory deficits—specifically diminished free recall but broadly spared retention and recognition—are evident in HIV (Woods et al., 2005). Hence, given the encoding and retrieval deficits experienced by this population, we would expect that the testing effect could improve memory due to: (1) enhancement of encoding via a number of cognitive processes (e.g., strengthening semantic representations through elaboration or through effortful retrieval processes, the restriction of the search-set to relevant associations) and/or (2) multiple retrieval routes, which may be particularly beneficial in a population with strategic retrieval impairment.

To our knowledge no study to date has investigated the testing effect in HIV+ persons. Still, given the importance of intact memory functioning for downstream health behaviors in individuals living with HIV (Hinkin et al., 2004) and previous studies suggesting that individuals with HIV can benefit from some memory improvement techniques (Weber et al., 2011), the present study aimed to evaluate the effectiveness of retrieval practice for improving memory in HIV+ adults following short (i.e., 30 min) and long retention intervals (i.e., 30 days). We expected that the participants, regardless of HIV serostatus, would perform better in testing condition than in restudy conditions.

Method

Participants

The study sample consisted of 73 participants aged 50 years or older recruited from the San Diego community and local HIV clinics as part of an NIH-funded study on aging and memory in HIV disease. HIV serostatus was confirmed using standard Western blot and/or a point-of-care test (MedMira Inc., Nova Scotia, Canada) and yielded 52 HIV-seropositive (HIV+) individuals and 21 HIV-seronegative comparison individuals (HIV-). Exclusion criteria included having a severe psychiatric disorder (e.g., schizophrenia) or neuromedical condition including an active central nervous system opportunistic infection, a seizure disorder, head injury with loss of consciousness for more than 30 minutes, stroke with persistent neurological sequelae, non-HIV-associated dementia, and an estimated verbal IQ score < 70 on the Wechsler Test of Adult Reading (WTAR; Psychological Corporation, 2001). Individuals were also excluded if they tested positive on a breathalyzer or urine toxicology screen for illicit drugs (except marijuana) on the day of testing. The participants' demographic, psychiatric, and HIV disease and treatment characteristics are displayed in Table 1. Although the groups were matched on the majority of demographic characteristic, HIV+ group was younger than HIV- group ($p < .05$) and included significantly higher proportions of men than the seronegative group ($p < .05$). Additionally memory impairment and depression were more common in the HIV+ group than in the HIV- group ($p < .05$).

Materials and Procedure

The university's human subjects committee approved this study. All participants provided written, informed consent prior to completing comprehensive medical, psychiatric, and neuropsychological research evaluations for which they received financial compensation.

Retrieval practice paradigm—This task was developed by Sumowski et al. (2010a, 2010b). In a within-subjects design, participants were presented with 48 weakly associated verbal paired associates (VPAs), which were divided across three learning conditions: Massed-Restudy, Spaced-Restudy, and Spaced-Testing. Each pair was individually presented on a computer screen three times in accordance with condition-specific schedules and VPAs were counterbalanced across learning conditions. Each presentation lasted for 6 seconds. The initial presentation of VPAs in the Massed-Restudy condition was immediately followed by two restudy trials; hence the participants were shown the same word pair in three consecutive trials. In the Spaced-Restudy condition, the initial VPA presentation was followed by three filler trials (i.e., other VPAs), a restudy trial, six filler trials, and a second

restudy trial. In the Spaced-Testing condition, the initial presentation was followed by three filler trials (i.e., other VPAs), a test trial, six filler trials, and a second test trial. Each test trial consisted of a 5-s cued-recall test (e.g., party - ____) followed by a 1-s feedback screen showing the completed VPA (e.g., party - serve). The paradigm is illustrated in Figure 1.

The paradigm isolated the effect of retrieval practice relative to restudy on memory via different types of cognitive process required by Spaced-Restudy and Spaced-Testing conditions. The only difference between the Spaced-Restudy and Spaced-Testing conditions was the type of cognitive reprocessing, which isolated the effect of retrieval practice during testing. The task also isolated the effect of spaced learning via different presentation schedules in Massed-Restudy and Spaced-Restudy conditions. The only difference between the Massed-Restudy and Spaced-Restudy conditions was the presentation schedule, which isolated the effect of spaced learning on memory (See Sumowski et al., 2010a See Sumowski et al., 2010b for further information on the development of the task).

The dependent variables were the performance on (1) short-delay cued recall (i.e., 30-min) and (2) long-delay cued recall (i.e., 30-day) tests. During the cued-recall task, the first word of each VPA was presented on the computer screen, and participants were asked to orally provide the second word of the pair. No feedback was provided in the delayed cued-recall tests. The order of VPAs during the test was quasi-randomized so that VPAs from each learning condition were equally represented at the beginning, middle, and end of the task.

Medical evaluation—Participants were administered a brief medical evaluation conducted by a research nurse and included a review of systems, medications, history, urine toxicology, and a blood draw.

Psychiatric evaluation—Current (i.e., within the last 30 days) and lifetime affective disorders (i.e., Major Depressive Disorder, Generalized Anxiety Disorder) were evaluated via the Composite International Diagnostic Interview (CIDI; version 2.128). The CIDI also provided lifetime diagnoses of Substance Use Disorders. Participants also completed the Profile of Mood States (POMS; McNair, Lorr, & Droppleman, 1971), which assesses symptoms of anxiety, depression, and other mood states over the past week.

Picture Sequence Memory—The Picture Sequence Memory Test in NIH Toolbox was used for the assessment of episodic memory. It involved recalling increasingly long series of illustrated objects and activities that have been presented in a particular order on the computer screen. The participants were asked to recall the sequence of pictures that were presented over two learning trials (for a detailed description of the test see Weintraub et al., 2013). Consistent with the recommendations of the Frascati criteria (Antinori et al., 2007), we used demographically adjusted normative standards (Casaletto et al., 2015) to quantify performance and used a 1 SD cut-point for identifying impaired memory function.

Data Analysis

To investigate our hypotheses regarding the performance on Verbal Paired Associate Learning paradigm as a function of learning conditions and HIV serostatus in short (30 min) retention interval, a 2 (HIV+ and HIV-) X 3 (Massed-Restudy, Spaced-Restudy, and Spaced-

Testing) mixed-design analysis of variance (ANOVA) was performed. To determine covariates for this model, we considered only variables that related to both independent and dependent variables at a critical alpha of .05. We analyzed comorbid variables (memory impairment, generalized anxiety disorder, major depressive disorder, substance abuse, and hepatitis C) and demographic variables (age, gender, estimated premorbid IQ, diabetes, hyperlipidemia). Of the variables (i.e., age, gender, memory impairment, and depression) that were associated with our fixed effect (see Table 1), memory impairment and depression scores were also associated with the short-delay memory performance outcomes; therefore these two variables were included in the models as fixed effects factor.

When appropriate, Cohen's d or partial eta squared (η_p^2) was used to report effect size estimates. Of note, the Shapiro-Wilk test of normality revealed that the data collected in cued-recall tests (i.e., short delay) failed the assumption of normality of parametric tests ($p < .05$). Since ANOVA-based tests are considered robust tests against normality assumption and there is no non-parametric corresponding test of ANOVA, the mixed-design ANOVA was determined to be the most appropriate statistical approach to analyze data in short-delay condition.

The analyses were conducted using the JMP 11.2 statistical program and the critical alpha was set at .05.

Results

The model with HIV, memory impairment, and depression as between-subject factors and learning conditions (Massed-Restudy, Spaced-Restudy, and Spaced-Testing conditions) as within-subjects factors yielded a large effect of learning condition (Pillai's Trace = .17, $F(2,68) = 32.76$, $p < .001$) where participants recalled 7.91 ($SE = .08$) of VPAs studied through Spaced-Testing compared with only 3.39 ($SE = .58$) through Massed-Restudy and 5.26 through Spaced-Restudy ($SE = .02$) with medium to large effect sizes between Massed-Restudy and Spaced-Testing ($d = .98$) and between Spaced-Restudy and Spaced-Testing ($d = .58$). The main effect of HIV status was not significant ($p = .22$) (See Figure 2a). The main effect of depression was marginally significant ($F(1, 69) = 3.35$, $p = .07$, $\eta_p^2 = .05$). Participants with history of major depression tended to remember fewer words pairs ($M = 4.77$, $SE = .81$) than participants with no history of major depression ($M = 6.81$, $SE = .56$). HIV status and depression did not interact with learning conditions ($p > .05$). The analysis revealed a significant main effect of memory impairment ($F(1, 69) = 5.36$, $p = .02$, $\eta_p^2 = .08$) such that individuals with memory impairment were less likely to remember VPAs. The interaction between memory impairment and learning conditions was marginally significant ($F(2, 68) = 2.86$, $p = .06$, $\eta_p^2 = .05$). Since there was only 1 HIV- participant with memory impairment, we sought to inspect the effect of memory impairment on outcome variables by performing within group comparisons only in the HIV+ group. We categorized participants in the HIV+ group into those with memory impairment (HIV+Imp) ($n = 14$) and those without memory impairment (HIV+Normal) ($n = 38$) according to their performance on the memory task from NIH Toolbox: As noted above, the cut-point for identifying impaired memory function has been established at 1 SD of the normative mean (Antinori et al., 2007). The analysis of variance showed that Spaced-Testing generated the superior memory

performance than Massed-Restudy ($d = 1.13$ and $d = .52$, for HIV+Normal and HIV+Imp, respectively) and than Spaced-Restudy ($d = .75$ and $d = .39$, HIV+Normal and HIV+Imp, respectively) ($p < .05$). The difference between Massed-Restudy and Spaced-Restudy was significant in HIV+Normal ($d = .39$, $p = .004$) but not in HIV+Imp group ($d = .15$, $p = .11$) (See Figure 2b).

Preliminary analysis of long-delay memory performance (i.e., number of pairs recalled in the cued-recall test) revealed that the participants scored poorly with very little variance ($M = .12$, $SE = .03$). Therefore, floor effects prevented further analysis of these data.

Discussion

Although memory impairment is common and functionally impactful in HIV+ persons, there are few established ways to improve it. The present study examined the testing effect via a verbal paired associate learning paradigm in an HIV+ sample as a potential memory improvement strategy. The results revealed that retrieval practice generated a strong and reliable memory gain for both people with and without HIV disease.

Findings showed that the testing condition improved memory in short-delay memory test, which was associated with medium to large effect sizes. Even spaced study, which is a well-established method to help people with dementia or other memory impairments (Balota et al., 2007), did not generate comparable learning rates. This point is important to mention because there is a large body of experimental work investigating spaced practice and these studies have generally reported a robust beneficial effect of distributed (over massed) presentation of to-be-remembered material (for meta-analysis, see Cepeda, Pashler, Vul, Wixted, & Rohrer, 2006) as was reported in the present study. Based on the findings of our study, we further suggest that spacing and testing can be combined—that is, spaced retrieval practice—in order to obtain the benefits of both.

Incidentally, the medium-to-large effect sizes observed in Spaced-Testing over the other two learning conditions (d between .58 and 1.13) deserve particular attention. The magnitude of these effects attests to the potential use of retrieval practice strategy for improving real-life memory performance of people with memory impairment (Sumowski et al., 2010a; 2010b) especially considering the scarcity of cognitive rehabilitation studies producing such large effect sizes. For example, methodical review of the literature concerning the effectiveness of cognitive rehabilitation of people with traumatic brain injury or stroke revealed that neurocognitive rehabilitation following brain injury are associated with improvements in attention, memory, and executive function (Cicerone et al. 2000; 2005; 2011). However only small fraction of improvement was a directly attributable to cognitive rehabilitation, which was evidenced by small treatment effect size ($d = .30$) (for meta-analyses see Rohling, Faust, Beverly, & Demakis, 2009). The medium-to-large effect sizes in this study and a similar study (Sumowski et al., 2010b) suggest that retrieval practice might be an effective memory improvement tool and should incite further exploration of the strategy in rehabilitation literature.

Another interesting finding was the absence of an interaction between HIV serostatus group and learning conditions. The serostatus groups performed similarly across all three learning conditions: they recalled more word pairs in the Spaced-Testing condition than in the restudy conditions. Moreover the testing effect was broadly comparable across the groups with medium to high effect sizes (i.e., d between .58 and .98). This finding indicates that the testing effect is robust cognitive phenomenon and can be used for memory improvement in HIV+ persons. It is equally important to note that the magnitude of the memory advantage of the testing effect relative to restudying in the HIV+ group with memory impairment was approximately half of that observed in the HIV+ group without memory impairment. The difference in effect sizes is noteworthy, especially since a liberal 1 SD cutoff was used to determine memory impairment. This finding might indicate that the magnitude of the testing effect may be slightly dampened, but still robust among individuals with fairly subtle memory impairment. Indeed, while the HIV+ group with memory impairment demonstrated somewhat inferior memory performance (i.e., recalled fewer number of word pairs) they still benefitted from retrieval practice as compared to the Spaced-Restudy and Massed-Restudy (d between .39 and .52) conditions (See Figure 2b). This might indicate that the effect was present in the individuals for whom such an intervention was most needed. Indeed, spaced testing in the memory impaired group was comparable to that of massed study in the normal group, suggesting that the strategy may be of some clinical benefit.

Although it was not the main focus of the present study, the marginal main effect of a lifetime history of major depression on the paired associates deserves brief discussion. Specifically, participants with a history of depression tended to perform poorly on the memory task than their non-depressed counterparts, irrespective of HIV serostatus ($p = .07$). Follow-up post-hoc tests showed that current symptoms of depression as measured by the POMS depression/dejection scale were significantly associated with poor cued recall memory performance ($r_s = -.29$, $p = .01$). The literature on the associations between depression and verbal memory is inconsistent (for a review, see Castaneda et al., 2008), particularly in the setting of HIV disease (e.g., Cysique et al., 2006). Some studies have observed verbal memory impairments among depressed participants compared to healthy adults (Smith et al., 2006), yet other studies have reported no deficits in this domain (Fossati et al., 1999; Wang et al., 2006). Although our results point to verbal memory impairment associated with depression, the trend level effect, cross-sectional design, small effect size, and small sample size prevent us from drawing any firm conclusions.

This study is not without its limitations. The first limitation concerns the paradigm used in the study. Specifically, one may argue that the absence of a Massed-Testing condition is a weakness of the task, since such a condition could allow one to conclude whether testing alone elicited an advantage relative to spaced restudying in adults with HIV, or if this advantage was only achieved when combined with spacing. However, we argue that the paradigm isolated the effect of retrieval practice during testing relative to restudy on memory: the only difference between Spaced-Testing and Spaced-Restudy was the cognitive processes required by the condition. Hence, the higher learning rates in the spaced testing generated by the retrieval practice. The task also isolated the effect of spaced learning via different presentation schedules in Massed-Restudy and Spaced-Restudy conditions, since the only difference between Massed-Restudy and Spaced-Restudy conditions was the

schedule of presentation, which isolated the effect of spaced learning on memory. Hence, the higher learning rates generated via spaced restudy as compared to massed restudy was due to the spacing effect. Since the effect of retrieval and spacing are captured separately by the task, we argue that higher learning rates in Spaced-Testing are due to the retrieval practice.

One might argue that the 1 SD cut-off for memory impairment is another limitation of the study. Indeed, there is some controversy in the neuroAIDS literature (as well as other conditions) as to what the optimal cut-points are to maximize sensitivity and specificity. However, to date, there is no persuasive evidence to support the benefits of using a 1.5 or 2 SD cut-point versus a 1 SD cut-point, with the former improving specificity but also reducing sensitivity (and in all cases, the literature is lacking rigorous gold standards against which to determine true classification accuracy). At present, the most widely used diagnostic cut-points are those offered by the Frascati criteria, and thus we anchor our approach to this literature. If we were to follow the more conservative 2 SD cut-points recommended by other groups (e.g., Gisslén, Price, & Nilsson, 2011) only 2% ($n = 1$) of the sample would meet the criteria, which would prevent us from conducting statistical analyses.

Another source of weakness in this study is the long-term retention task (i.e., cued-recall task). The long-term retention performance of the participant indicated that cued-recall test following a long delay interval (i.e., 30 days) was difficult for our participants, which manifested through floor effects. An easier task, such as recognition, or a shorter delay interval might have been more suitable for the current population.

In addition, the generalizability of these results is subject to certain limitations. First, the small sample size limits the generalization of our findings. The attrition rate of the participants in the second session limits the generalizability of the testing effect even further. Second, further investigation of the effect in other modalities and ecologically relevant materials (e.g., health related information) is needed.

To sum up, our findings indicate that the testing effect can enhance episodic verbal memory performance among persons living with HIV, including those with clinical memory deficits. Such findings are consistent with research in other neurological conditions that experience mild-to-moderate memory deficits such as multiple sclerosis (e.g., Basso et al., 2006; Chiaravalloti & DeLuca, 2002;), traumatic brain injury (e.g., O'Brien et al., 2007), and mild dementia in early AD (e.g., Lipinska et al., 1994). The present research provides the first known evidence for the mnemonic value of the testing effect relative to different study schedules among individuals with HIV. Considering the notable adverse affects that memory deficits have on everyday functioning, health behaviors, and quality of life in HIV, we argue that it is particularly important to identify effective ways to improve such deficits. Hence, a theoretically-driven and empirically supported neurocognitive rehabilitation approach is imperative to this long-term goal to improve quality of life among persons with HIV disease.

Acknowledgments

This work was supported by the National Institutes of Health under grants R01-MH073419 and P30-MH62512. The authors have no financial conflicts of interest related to this work. The views expressed in this article are those of the authors and do not reflect the official policy or position of the Department of the Navy, Department of Defense, nor the United States Government. The authors thank Donald Franklin, Stephanie Corkran, Jessica Beltran, and Dr.

Erin E. Morgan for their help with data processing and study coordination. The San Diego HIV Neurobehavioral Research Program [HNRP] group is affiliated with the University of California, San Diego, the Naval Hospital, San Diego, and the Veterans Affairs San Diego Healthcare System, and includes: Director: Igor Grant, M.D.; Co-Directors: J. Hampton Atkinson, M.D., Ronald J. Ellis, M.D., Ph.D., and J. Allen McCutchan, M.D.; Center Manager: Thomas D. Marcotte, Ph.D.; Jennifer Marquie-Beck, M.P.H.; Melanie Sherman; Neuromedical Component: Ronald J. Ellis, M.D., Ph.D. (P.I.), J. Allen McCutchan, M.D., Scott Letendre, M.D., Edmund Capparelli, Pharm.D., Rachel Schrier, Ph.D., Debra Rosario, M.P.H., Neurobehavioral Component: Robert K. Heaton, Ph.D. (P.I.), Mariana Cherner, Ph.D., Jennifer E. Iudicello, Ph.D., David J. Moore, Ph.D., Erin E. Morgan, Ph.D., Matthew Dawson; Neuroimaging Component: Terry Jernigan, Ph.D. (P.I.), Christine Fennema-Notestine, Ph.D., Sarah L. Archibald, M.A., John Hesselink, M.D., Jacopo Annese, Ph.D., Michael J. Taylor, Ph.D.; Neurobiology Component: Eliezer Masliah, M.D. (P.I.), Cristian Achim, M.D., Ph.D., Ian Everall, FRCPsych., FRCPPath., Ph.D. (Consultant); Neurovirology Component: Douglas Richman, M.D., (P.I.), David M. Smith, M.D.; International Component: J. Allen McCutchan, M.D., (P.I.); Developmental Component: Cristian Achim, M.D., Ph.D.; (P.I.), Stuart Lipton, M.D., Ph.D.; Participant Accrual and Retention Unit: J. Hampton Atkinson, M.D. (P.I.); Data Management Unit: Anthony C. Gamst, Ph.D. (P.I.), Clint Cushman (Data Systems Manager); Statistics Unit: Ian Abramson, Ph.D. (P.I.), Florin Vaida, Ph.D., Reena Deutsch, Ph.D., Anya Umlauf, M.S.

References

- Antinori A, Arendt G, Becker JT, Brew BJ, Byrd DA, Cherner M, et al. Updated research nosology for HIV-associated neurocognitive disorders. *Neurology*. 2007; 69:1789–1799. DOI: 10.1212/01.wnl.0000287431.88658.8b [PubMed: 17914061]
- Balota, DA., Duchek, JM., Logan, JM. Is expanded retrieval practice a superior form of spaced retrieval? A critical review of the extant literature. In: Nairne, JS., editor. *The foundations of remembering*. New York: Psychology Press; 2007. p. 83-105.
- Basso MR, Lowery N, Ghormley C, Combs D, Johnson J. Self-generated learning in people with multiple sclerosis. *Journal of International Neuropsychology Society*. 2006; 12(05):640–648. DOI: 10.1017/s1355617706060759
- Carpenter SK. Semantic Information Activated During Retrieval Contributes to Later Retention: Support for the Mediator Effectiveness Hypothesis of the Testing Effect. *Journal of Experimental Psychology: Learning, Memory, & Cognition*. 2011; 37:1547–1552. DOI: 10.1037/e520602012-165
- Carpenter SK, DeLosh EL. Application of the testing and spacing effects to name learning. *Applied Cognitive Psychology*. 2005; 19(5):619–636. DOI: 10.1002/acp.1101
- Casaleto KB, Umlauf A, Beaumont J, Gershon R, Slotkin J, Akshoomoff N, Heaton RK. Demographically Corrected Normative Standards for the English Version of the NIH Toolbox Cognition Battery. *Journal of International Neuropsychology Society*. 2015; 21(05):378–391. DOI: 10.1017/s1355617715000351
- Castaneda AE, Tuulio-Henriksson A, Marttunen M, Suvisaari J, Lönnqvist J. A review on cognitive impairments in depressive and anxiety disorders with a focus on young adults. *Journal of Affective Disorders*. 2008; 106(1-2):1–27. DOI: 10.1016/j.jad.2007.06.006 [PubMed: 17707915]
- Cepeda NJ, Pashler H, Vul E, Wixted JT, Rohrer D. Distributed practice in verbal recall tasks: A review and quantitative synthesis. *Psychological Bulletin*. 2006; 132(3):354–380. DOI: 10.1037/0033-2909.132.3.354 [PubMed: 16719566]
- Chiaravalloti ND, DeLuca J. Self-generation as a means of maximizing learning in multiple sclerosis: An application of the generation effect. *Archives of Physical Medicine and Rehabilitation*. 2002; 83(8):1070–1079. DOI: 10.1053/apmr.2002.33729 [PubMed: 12161827]
- Cicerone KD, Dahlberg C, Kalmar K, Langenbahn DM, Malec JF, Bergquist TF, et al. Morse PA. Evidence-based cognitive rehabilitation: Recommendations for clinical practice. *Archives of Physical Medicine and Rehabilitation*. 2000; 81(12):1596–1615. DOI: 10.1053/apmr.2000.19240 [PubMed: 11128897]
- Cicerone KD, Dahlberg C, Malec JF, Langenbahn DM, Felicetti T, Kneipp S, et al. Catanese J. Evidence-Based Cognitive Rehabilitation: Updated Review of the Literature From 1998 Through 2002. *Archives of Physical Medicine and Rehabilitation*. 2005; 86(8):1681–1692. DOI: 10.1016/j.apmr.2005.03.024 [PubMed: 16084827]
- Cicerone KD, Langenbahn DM, Braden C, Malec JF, Kalmar K, Fraas M, et al. Ashman T. Evidence-Based Cognitive Rehabilitation: Updated Review of the Literature From 2003 Through 2008.

- Archives of Physical Medicine and Rehabilitation. 2011; 92(4):519–530. DOI: 10.1016/j.apmr.2010.11.015 [PubMed: 21440699]
- Cysique LA, Deutsch R, Atkinson JH, Young C, Marcotte TD, Dawson L, et al. Heaton RK. Incident major depression does not affect neuropsychological functioning in HIV-infected men. *Journal of the International Neuropsychological Society*. 2006; 13(01)doi: 10.1017/s1355617707070026
- Dunlosky J, Rawson KA, Marsh EJ, Nathan MJ, Willingham DT. Improving students' learning with effective learning techniques: promising directions from cognitive and educational psychology. *Psychological Science in the Public Interest*. 2013; 14:4–58. DOI: 10.1177/1529100612453266 [PubMed: 26173288]
- Ellis R, Langford D, Masliah E. HIV and antiretroviral therapy in the brain: neuronal injury and repair. *Nature Reviews Neuroscience*. 2007; 8(1):33–44. DOI: 10.1038/nrn2040 [PubMed: 17180161]
- Ernst T, Chang L. Effect of aging on brain metabolism in antiretroviral-naïve HIV patients. *AIDS*. 2004; :61–67. DOI: 10.1097/00002030-200418001-00009
- Fossati P, Amar G, Raoux N, Ergis AM, Allilaire JF. Executive functioning and verbal memory in young patients with unipolar depression and schizophrenia. *Psychiatry Research*. 1999; 89(3):171–187. DOI: 10.1016/s0165-1781(99)00110-9 [PubMed: 10708264]
- Heaton RK, Grant I, Butters N, White DA, Kirson D, Atkinson JH, et al. The HNRC 500—Neuropsychology of HIV infection at different disease stages. *Journal of the International Neuropsychological Society*. 1995; 1:231–251. DOI: 10.1017/s1355617700000230 [PubMed: 9375218]
- Heaton RK, Marcotte TD, Mindt MR, Sadek J, Moore DJ, Bentley H, et al. Grant I. The impact of HIV-associated neuropsychological impairment on everyday functioning. *Journal of International Neuropsychology Society*. 2004; 10(03)doi: 10.1017/s1355617704102130
- Hinkin CH, Hardy DJ, Mason KI, Castellon SA, Durvasula RS, Lam MN, Stefaniak M. Medication adherence in HIV-infected adults. *AIDS*. 2004; :19–25. DOI: 10.1097/00002030-200418001-00004
- Gisslén M, Price RW, Nilsson S. The definition of HIV-associated neurocognitive disorders: are we overestimating the real prevalence? *BMC Infectious Diseases*. 2011; 11(1)doi: 10.1186/1471-2334-11-356
- Gongvatana A, Woods SP, Taylor MJ, Vigil O, Grant I. Semantic Clustering Inefficiency in HIV-Associated Dementia. *The Journal of Neuropsychiatry and Clinical Neurosciences*. 2007; 19(1): 36–42. DOI: 10.1176/jnp.2007.19.1.36 [PubMed: 17308225]
- Karpicke, JD., Lehman, M., Aue, WR. Retrieval-based learning: An episodic context account. In: Ross, BH., editor. *The Psychology of Learning and Motivation*. Vol. 61. San Diego, CA: Elsevier Academic Press; 2014. p. 237-284.
- Lipinska B, Bäckman L, Mäntylä T, Viitanen M. Effectiveness of self-generated cues in early Alzheimer's disease. *Journal of Clinical and Experimental Neuropsychology*. 1994; 16:809–819. DOI: 10.1080/01688639408402695 [PubMed: 7890817]
- McNair, DM., Lorr, M., Droppleman, LF. *Manual for the Profile of Mood States*. San Diego, CA: Educational and Industrial Testing Services; 1971.
- Middleton EL, Schwartz MF. Errorless learning in cognitive rehabilitation: A critical review. *Neuropsychological Rehabilitation*. 2012; 22(2):138–168. DOI: 10.1080/09602011.2011.639619 [PubMed: 22247957]
- Murji S, Rourke SB, Donders J, Carter SL, Shore D, Rourke BP. Theoretically derived CVLT subtypes in HIV-1 infection: Internal and external validation. *Journal of International Neuropsychology Society*. 2003; 9(01)doi: 10.1017/s1355617703910010
- O'brien A, Chiaravalloti N, Carlos Arango-Lasprilla J, Lengenfelder J, Deluca J. An investigation of the differential effect of self-generation to improve learning and memory in multiple sclerosis and traumatic brain injury. *Neuropsychological Rehabilitation*. 2007; 17(3):273–292. DOI: 10.1080/09602010600751160 [PubMed: 17474057]
- Pyc MA, Rawson KA. Testing the retrieval effort hypothesis: Does greater difficulty correctly recalling information lead to higher levels of memory? *Journal of Memory and Language*. 2009; 60:437–447. DOI: 10.1016/j.jml.2009.01.004

- Rippeth JD, Heaton RK, Carey CL, Marcotte TD, Moore DJ, Gonzalez R, et al. Methamphetamine dependence increases risk of neuropsychological impairment in HIV infected persons. *Journal of the International Neuropsychological Society*. 2004; 10:1–14. DOI: 10.1017/s1355617704101021 [PubMed: 14751002]
- Roediger HL, Butler AC. The critical role of retrieval practice in long-term retention. *Trends in Cognitive Sciences*. 2011; 15(1):20–27. DOI: 10.1016/j.tics.2010.09.003 [PubMed: 20951630]
- Roediger HL, Karpicke JD. Test-Enhanced Learning: Taking Memory Tests Improves Long-Term Retention. *Psychological Science*. 2006; 17(3):249–255. DOI: 10.1111/j.1467-9280.2006.01693.x [PubMed: 16507066]
- Rohling ML, Faust ME, Beverly B, Demakis G. Effectiveness of cognitive rehabilitation following acquired brain injury: a meta-analytic re-examination of Cicerone et al.'s (2000, 2005) systematic reviews. *Neuropsychology*. 2009; 23(1):20–39. doi:/10.1037/a0013659. [PubMed: 19210030]
- Smith DJ, Muir WJ, Blackwood DH. Neurocognitive impairment in euthymic young adults with bipolar spectrum disorder and recurrent major depressive disorder. *Bipolar Disorders*. 2006; 8(1): 40–46. DOI: 10.1111/j.1399-5618.2006.00275.x [PubMed: 16411979]
- Sumowski JF, Chiaravalloti N, DeLuca J. Retrieval practice improves memory in multiple sclerosis: Clinical application of the testing effect. *Neuropsychology*. 2010a; 24(2):267–272. DOI: 10.1037/a0017533 [PubMed: 20230121]
- Sumowski JF, Wood HG, Chiaravalloti N, Wylie GR, Lengenfelder J, Deluca J. Retrieval practice: A simple strategy for improving memory after traumatic brain injury. *Journal of International Neuropsychology Society*. 2010b; 16(06):1147–1150. DOI: 10.1017/s1355617710001128
- Sumowski JF, Leavitt VM, Cohen A, Paxton J, Chiaravalloti ND, DeLuca J. Retrieval practice is a robust memory aid for memory-impaired patients with MS. *Multiple Sclerosis Journal*. 2013; 19(14):1943–1946. DOI: 10.1177/1352458513485980 [PubMed: 23674564]
- van Gorp WG, Rabkin JG, Ferrando SJ, Mintz J, Ryan E, Borkowski T, McElhiney M. Neuropsychiatric predictors of return to work in HIV/AIDS. *Journal of International Neuropsychology Society*. 2006; 13(01)doi: 10.1017/s1355617707070117
- Wang CE, Halvorsen M, Sundet K, Steffensen AL, Holte A, Waterloo K. Verbal memory performance of mildly to moderately depressed outpatient younger adults. *Journal of Affective Disorders*. 2006; 92(2-3):283–286. DOI: 10.1016/j.jad.2006.02.008 [PubMed: 16533537]
- Weber E, Blackstone K, Woods SP. Cognitive neurorehabilitation of HIV-associated neurocognitive disorders: A qualitative review and call to action. *Neuropsychology Review*. 2013; 23(1):81–98. DOI: 10.1007/s11065-013-9225-6 [PubMed: 23417497]
- Weber E, Woods SP, Kellogg E, Grant I, Basso MR. Self-generation enhances verbal recall in individuals infected with HIV. *Journal of International Neuropsychology Society*. 2011; 18(01): 128–133. DOI: 10.1017/s135561771100124x
- Weintraub S, Dikmen SS, Heaton RK, Tulsky DS, Zelazo PD, Bauer PJ, et al. Gershon RC. Cognition assessment using the NIH Toolbox. *Neurology*. 2013; 80(Issue 11, Supplement 3):S54–S64. DOI: 10.1212/wnl.0b013e3182872ded [PubMed: 23479546]
- Woods SP, Scott JC, Dawson MS, Morgan EE, Carey CL, Heaton RK, et al. Construct validity of Hopkins verbal learning test-revised component process measures in an HIV-1 sample. *Archives of Clinical Neuropsychology*. 2005; 20:1061–1071. DOI: 10.1016/j.acn.2005.06.007 [PubMed: 16198529]
- Woods PS, Morgan EE, Dawson M, Cobb Scott J, Grant I. The HIV Neurobehavioral Research Center. Action (verb) fluency predicts dependence in instrumental activities of daily living in persons infected with HIV-1. *Journal of Clinical and Experimental Neuropsychology*. 2006; 28(6):1030–1042. DOI: 10.1080/13803390500350985 [PubMed: 16822741]
- World Health Organization. Composite International Diagnostic Interview (CIDI, Version 2.1). Geneva, Switzerland: World Health Organization; 1998.

Figure 1.
The retrieval practice paradigm adopted from Sumowski et al. (2010a; 2010b).

Figure 2. Short-delay cued-recall memory performance across all three study conditions. Error bars represent standard errors.

Table 1

Cohort's demographic, psychiatric, medical, and HIV disease characteristics.

	HIV - (n = 21)	HIV + (n = 52)	p	Group Differences
Age (years)	61.66 (1.34)	55.86 (0.85)	< .001	HIV+ > HIV-
Education (years)	14.23 (0.61)	13.63 (0.39)	0.41	--
Ethnicity (% White)	57.14	63.46	0.61	--
Gender (% men)	71.43	90.38	0.04	HIV+ > HIV-
NIH Toolbox Premorbid IQ	109.56 (1.73)	109.74 (1.73)	0.95	--
Memory Impairment (% impaired)	4.76	26.92	0.01	HIV+ > HIV-
Generalized Anxiety Disorder (%)	4.76	17.31	0.12	--
Major Depressive Disorder (%)	38.1	73.08	0.006	HIV+ > HIV-
Substance Abuse (%)	47.62	69.23	0.08	--
Diabetes (%)	15	17	0.99	--
Hyperlipidemia (%)	35	55	0.1	--
Hepatitis C Virus (%)	14.29	26.92	0.23	--
HIV Duration (years)	-	19.79 (1.64)	--	--
AIDS (%)	-	47.22	--	--
CD4 Count (cells/ μ L)	-	663.28 (41.71)	--	--
Nadir CD4 (cells/ μ L)	-	174.73 (25.09)	--	--
Prescribed cART Status (%)	-	98.08	--	--
RNA in Plasma Detectable (%)	-	8	--	--
Among Subjects on cART	-	7	--	--

Note. Data represents M (SE) or %. AIDS = Acquired Immune Deficiency Syndrome; CD4 = Cluster of Differentiation 4; cART = combination antiretroviral therapy