Published in final edited form as:

J Card Fail. 2017 May ; 23(5): 427–431. doi:10.1016/j.cardfail.2017.02.003.

Participation in Cardiac Rehabilitation among Patients with Heart Failure

Linda G. Park, PhD, NP^{a,b}, David W. Schopfer, MD, MAS^{a,c}, Ning Zhang, MS^a, Hui Shen, MS^a, and Mary A. Whooley, MD^{a,c,d}

^aSan Francisco Veterans Affairs Medical Center, San Francisco, CA

^bSchool of Nursing, Department of Community Health Systems, University of California, San Francisco, CA

^cDepartment of Medicine, General Internal Medicine, University of California, San Francisco, CA

^dDepartment of Medicine, Epidemiology, and Biostatistics, University of California, San Francisco, CA

Abstract

Objectives—We sought to determine: a) the proportion of patients with heart failure (HF) who participated in cardiac rehabilitation (CR); and b) patient characteristics associated with participation.

Background—CR is linked to reduced mortality and morbidity including improvements in cardiorespiratory fitness, psychosocial state, and quality of life in patients with HF. However, little is known about CR utilization among patients with HF.

Methods—A retrospective study was conducted using national data from the Centers for Medicare & Medicaid Services and the Veterans Health Administration (VA). We used primary discharge ICD-9 codes to identify patients hospitalized for HF from 2007–2011 then identified CR participation using CPT codes from claims data. Multivariate logistic regression was used to identify patient characteristics associated with CR participation.

Results—There were 66,710 Veterans and 243,208 Medicare beneficiaries hospitalized for HF and 1,554 (2.3%) and 6,280 (2.6%), respectively, who attended one or more sessions of outpatient CR. Among Medicare beneficiaries, men were more likely than women to participate in CR (3.7% vs. 1.8%; $p < 0.001$), but there was no gender difference among Veterans (2.3% vs. 2.8%; $p = 0.40$). Characteristics associated with participation in CR among both groups included younger age, white race, and history of ischemic heart disease.

Corresponding Author: Linda G. Park, PhD, MS, NP, Assistant Professor, Department of Community Health Systems, University of California, San Francisco, 2 Koret Way, Box 0608, San Francisco, CA 94143-0608, Phone: 415-502-6616, Fax: 415-476-6042, linda.park@ucsf.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conclusions—Very few HF patients participated in CR with lower rates among those who were older, non-white, and female with a history of depression or other chronic medical conditions. Since Medicare has recently introduced coverage for CR in patients with systolic HF, we must increase efforts to improve CR participation, especially among these vulnerable groups.

Keywords

cardiovascular disease; Medicare; Veterans; exercise

A focus of heart failure (HF) management is to reduce symptoms and improve exercise capacity. Exercise-based cardiac rehabilitation (CR) is a comprehensive secondary prevention program involving exercise training, risk factor modification, education, and counseling to reduce the burden of the effects of cardiovascular illness.^{1,2} CR participation is associated with reduced morbidity and mortality as well as improvement of quality of life, symptoms and functional capacity.^{3–6} Clinical guidelines on HF management include exercise training and CR as a Class I recommendation.⁷

Despite these recommendations, CR is still vastly underutilized in the United States.^{3,7–11} One major barrier to referral of CR included lack of insurance coverage for HF.¹² However, in February 2014, Centers for Medicare & Medicaid Services (CMS) introduced coverage of CR for patients with HF.¹³ As a result of these recent policy changes, more patients with HF are expected to enroll in CR.

Few studies have evaluated the proportion of patients with HF who participate in CR. A recent study of U.S. hospitals in the Get With the Guidelines database showed a 10.4% *referral* rate to CR for patients with HF prior to the CMS approval,¹⁴ but the *participation* rate in CR is unknown. The Veterans Health Administration (VA) has provided CR services for Veterans with HF independent of CMS approval, which makes it a unique population to study CR use without the potential barrier of insurance coverage. By analyzing both VA and Medicare data in this retrospective study, we sought to determine: 1) the proportion of patients with HF who participated in CR during a 5-year period; and 2) significant patient factors associated with participation vs. non-participation.

Methods

Study Design and Data Collection

A retrospective study was conducted using national data from two separate data sources: VA Healthcare System and the Medicare claim files. Patients were identified using a random sample of Medicare beneficiaries from a 5% sample of Medicare claims and VA hospital discharge records through CMS and VA datasets. We identified unique patients who were discharged from inpatient medical facilities with a diagnosis of HF from 1/1/2007 – 12/31/11. Primary ICD-9 codes were used to define HF. Patients were excluded if death occurred within 30 days of hospitalization. We extracted data on comorbidities that appeared once in inpatient data or twice in outpatient data.

We defined CR participation as one or more CR sessions within 12 months after hospitalization for HF. The number of unique patients who participated in CR programs was obtained using outpatient files with CPT codes for CR.

Data Analysis

We compared sociodemographic characteristics and comorbidities of CR participants and non-participants for both VA and CMS data. Characteristics of patients who participated in CR were compared to those who did not participate in CR using t-tests or Wilcoxon rank-sum tests for continuous variables and chi-square analyses for dichotomous or categorical variables. A multivariable logistic regression was used to determine factors independently associated with CR participation. Variables that were significantly associated with CR participation at an *a priori* significance level of $p < 0.05$ were entered into a multivariable model. SAS version 9.3 (SAS Institute Inc., Cary, NC) was used for all analyses.

Results

During the five-year period, 66,710 Veterans were hospitalized for HF. Only 1,554 (2.3%) attended at least one session of outpatient CR. Among Medicare beneficiaries, 243,208 were hospitalized for HF with 6,208 (2.6%) participating in at least one session of outpatient CR (Figure 1). For Medicare beneficiaries, CR participation increased from 1.3 to 3.1%, while participation rates remained relatively flat among Veterans in the same time period (2.0 to 2.1%).

After adjusting for sociodemographic and comorbid differences, men were more likely than women to participate in CR among Medicare beneficiaries [odds ratio (OR) 1.57; CI 1.49–1.65; $P < 0.0001$], but there was no difference among Veterans (Table 1). Younger age and white race were both associated with greater participation. Among Veterans, being married and higher income were associated with greater participation. Ischemic heart disease and hyperlipidemia were significant predictors of CR participation in both groups (OR 7.51, CI 6.53–8.62; $P < 0.0001$). Chronic conditions including diabetes, cerebrovascular disease, chronic obstructive pulmonary disease, chronic kidney disease, and depression were also associated with less participation in CR.

Discussion

Participation in one or more sessions of outpatient CR was extremely low at 2.3% among Veterans and 2.6% among Medicare beneficiaries who were hospitalized for HF between 2007–2011. Although these rates were expected to be low, new strategies must be developed to increase CR participation in the HF population since coverage for CR for HF has been introduced. A major reason for underutilization of CR is likely the requirement for HF with reduced ejection fraction (HFrEF), although more than half of patients with HF have preserved ejection fraction (HFpEF) with comparable morbidity and mortality outcomes.¹⁵ It is informative that participation was extremely low for Veterans with HF where reimbursement was not a factor. This result supports the fact that there are other significant contributing factors to lack of CR referral and participation.

Compared with the recent findings of a 10.4% referral rate for patients after hospitalization for HF to CR,¹⁴ our study included three times as many patients and more importantly, the former study investigated *referral* to CR whereas our study evaluated CR participation. Clearly both efforts to increase referral and participation need improvement and require further investigation after the CMS approval in 2014.

In addition to the major historical barrier of insurance coverage for patients with HF, the reasons for low utilization of CR are multifactorial such as transportation issues, lack of physician referral, fear of exercise and injury, etc.^{16–18} Since CR has traditionally been for patients with IHD and cardiac surgery, patients with HF are less likely to be prescribed CR. Many misconceptions of the potential harm of exercise causing ventricular remodeling hindered the CR referral.¹⁸ Access to transportation and geographic distance to CR programs is a major barrier as nearly 24% of VA enrollees are from rural areas¹⁹ and 76% of VA enrolled Veterans live over one hour from a VA CR center. Also, only stable patients with HFrEF are currently considered eligible for CR.

The gender difference we found in Medicare beneficiaries with HF is consistent with trends that have been established over the past decade about CR referral, enrollment, and adherence.^{20–22} Although women may have greater morbidity and mortality in the short term period following HF hospitalization,²³ they are significantly less likely than men to access CR.²⁴ Barriers to CR participation that women have reported include perceiving exercise as tiring or painful, transportation, family responsibilities, and comorbidities (e.g., musculoskeletal issues).²⁵

We found less CR participation in Medicare patients with comorbidities (e.g., COPD, chronic kidney disease, diabetes, etc.). We speculate the reasons for less participation are multifactorial such as frailty, intolerance to exercise, depression, etc. More attention is needed in these high risk patients who would benefit from cardiopulmonary and cardiovascular improvement as well as self-care education and social support.

This study has several strengths and limitations that should be considered when interpreting the results. First, the most significant strength of this study is our presentation of CR participation rates, not referral rates. Second, this study includes a national sample of both Medicare and Veteran patients. All of the VA facilities were included; therefore, the potential variation in CR participation across medical facilities was accounted for with its own unique characteristics such as different patient populations and delivery of health care services. The most important limitation to consider is that these analyses reflect patients who received CR after being hospitalized with HF and not from an outpatient database. It is more common for patients to be referred for CR from an inpatient hospitalization¹⁴; therefore, we do not believe the lack of outpatient sources was significant. Also, since these patients were referred prior to HF approval, it is unclear if ejection fraction would have been a factor. Veterans with dual coverage who may have attended a non-VA CR program may not have been identified. However, this number is small and probably would not significantly change our results. Finally, we did not include musculoskeletal conditions as comorbidities in our analysis although they have been known to be a barrier for CR among women.²⁵

Conclusion

We found that CR was vastly underutilized among patients hospitalized for HF over a 5-year period among both VA and CMS beneficiaries. Because these data were prior to the CMS amendment to include CR as a benefit for patients with HF, we expected to find low participation rates. However, the similar rates between Medicare and VA patients where CR for patients with HF has been a covered benefit was surprising. This suggests that insurance coverage of CR for patients with HF is not the major factor for low participation. Instead, we found that less CR participation is associated with having comorbidities. Data from this study provide a basis for establishing goals to improve clinical care of patients with HF that include addressing the burden of comorbidities.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We would like to thank Chris Vercammen, MD, MS, MPH for his statistical support.

Conflicts of Interest and Source of Funding: Dr. Park was supported by the Veterans Affairs Quality Scholars Fellowship Program to complete this project. This project was supported by the Department of Veterans Affairs Health Services Research and Development Measurement Science QUERI (IP1 HX002002), the National Center for Advancing Translational Sciences of the NIH (Award Number KL2TR001870), and by the UCSF Clinical Translational Sciences Institute (UL1 TR000004 RR024131). Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the NIH.

References

1. Balady GJ, Williams MA, Ades PA, Bittner V, Comoss P, Foody JM, et al. Core components of cardiac rehabilitation/secondary prevention programs: 2007 update: a scientific statement from the American Heart Association Exercise, Cardiac Rehabilitation, and Prevention Committee, the Council on Clinical Cardiology; the Councils on Cardiovascular Nursing, Epidemiology and Prevention, and Nutrition, Physical Activity, and Metabolism; and the American Association of Cardiovascular and Pulmonary Rehabilitation. *Circulation*. May-Jun;2007 27(3):121–9. DOI: 10.1161/circulationaha.106.180945
2. O'Connor CM, Whellan DJ, Lee KL, Keteyian SJ, Cooper LS, Ellis SJ, et al. Efficacy and safety of exercise training in patients with chronic heart failure: HF-ACTION randomized controlled trial. *JAMA*. 2009; 301(14):1439–50. DOI: 10.1001/jama.2009.454 [PubMed: 19351941]
3. Thomas RJ, King M, Lui K, Oldridge N, Piña IL, Spertus J. AACVPR/ACC/AHA 2007 performance measures on cardiac rehabilitation for referral to and delivery of cardiac rehabilitation/secondary prevention services. *Circulation*. 2007; 116(14):1611–42. DOI: 10.1016/j.jacc.2007.04.033 [PubMed: 17885210]
4. Downing J, Balady GJ. The role of exercise training in heart failure. *J Am Coll Cardiol*. 2011; 58(6): 561–9. DOI: 10.1016/j.jacc.2011.04.020 [PubMed: 21798416]
5. Flynn KE, Piña IL, Whellan DJ, Blumenthal JA, Ellis SJ, Fine LJ, et al. Effects of exercise training on health status in patients with chronic heart failure: HF-ACTION randomized controlled trial. *JAMA*. 2009; 301(14):1451–9. DOI: 10.1001/jama.2009.457 [PubMed: 19351942]
6. Sagar VA, Davies EJ, Briscoe S, Coats AJ, Dalal HM, Lough F, et al. Exercise-based rehabilitation for heart failure: systematic review and meta-analysis. *Open Heart*. 2015; 2(1):e000163.doi: 10.1136/openhrt-2014-000163 [PubMed: 25685361]
7. Ades PA, Waldmann ML, McCann WJ, Weaver SO. Predictors of cardiac rehabilitation participation in older coronary patients. *Arch Intern Med*. 1992; 152(5):1033–35. [PubMed: 1580707]

8. Suaya JA, Shepard DS, Normand SL, Ades PA, Prottas J, Stason WB. Use of cardiac rehabilitation by Medicare beneficiaries after myocardial infarction or coronary bypass surgery. *Circulation*. 2007; 116(15):1653–62. DOI: 10.1161/CIRCULATIONAHA.107.701466 [PubMed: 17893274]
9. Dolansky MA, Moore SM. Older adults' use of postacute and cardiac rehabilitation services after hospitalization for a cardiac event. *Rehabil Nurs*. 2008; 33(2):73–81. DOI: 10.1002/j.2048-7940.2008.tb00207.x [PubMed: 18330386]
10. Arena R, Williams M, Forman DE, Cahalin LP, Coke L, Myers J, et al. Increasing referral and participation rates to outpatient cardiac rehabilitation: the valuable role of healthcare professionals in the inpatient and home health settings: a science advisory from the American Heart Association. *Circulation*. 2012; 125(10):1321–9. DOI: 10.1161/CIR.0b013e318246b1e5 [PubMed: 22291128]
11. Balady GJ, Ades PA, Bittner VA, Franklin BA, Gordon NF, Thomas RJ, et al. Referral, enrollment, and delivery of cardiac rehabilitation/secondary prevention programs at clinical centers and beyond: a presidential advisory from the American Heart Association. *Circulation*. 2011; 124(25):2951–60. DOI: 10.1161/CIR.0b013e31823b21e2 [PubMed: 22082676]
12. Goel K, Lennon RJ, Tilbury RT, Squires RW, Thomas RJ. Impact of cardiac rehabilitation on mortality and cardiovascular events after percutaneous coronary intervention in the community. *Circulation*. 2011; 123(21):2344–52. DOI: 10.1161/CIRCULATIONAHA.110.983536 [PubMed: 21576654]
13. Centers for Medicare & Medicaid. [Accessed June 16, 2014] Decision memo for cardiac rehabilitation (CR) programs - Chronic heart failure (CAG-00437N). <https://www.cms.gov/medicare-coverage-database/details/nca-decision-memo.aspx?NCAId=270>
14. Golwala H, Pandey A, Ju C, Butler J, Yancy C, Bhatt DL, et al. Temporal trends and factors associated with cardiac rehabilitation referral among patients hospitalized with heart failure. *J Am Coll Cardiol*. 2015; 66:917–26. DOI: 10.1016/j.jacc.2015.06.1089 [PubMed: 26293762]
15. Quiroz R, Doros G, Shaw P, Liang CS, Gauthier DF, Sam F. Comparison of characteristics and outcomes of patients with heart failure preserved ejection fraction versus reduced left ventricular ejection fraction in an urban cohort. *Am J Cardiol*. 2014; 113(4):691–6. DOI: 10.1016/j.amjcard.2013.11.014 [PubMed: 24484862]
16. Neubeck L, Freedman SB, Clark AM, Briffa T, Bauman A, Redfern J. Participating in cardiac rehabilitation: a systematic review and meta-synthesis of qualitative data. *Eur J Prev Cardiol*. 2012; 19(3):494–503. DOI: 10.1177/1741826711409326 [PubMed: 22779092]
17. Schopfer DW, Priano S, Allsup K, Helfrich CD, Ho PM, Rumsfeld JS, et al. Factors associated with utilization of cardiac rehabilitation among patients with ischemic heart disease in the Veterans Health Administration: A qualitative study. *J Cardiopulm Rehabil*. 2016; 36(3):167–73. DOI: 10.1097/HCR.0000000000000166
18. Schopfer DW, Forman DE. Growing relevance of cardiac rehabilitation for an older population with heart failure. *J Card Fail*. 2016; 22(12):1015–22. DOI: 10.1016/j.cardfail.2016.10.010 [PubMed: 27769907]
19. Veterans Health Administration Office of Rural Health. [Accessed on June 1, 2016] Fact Sheet: Information about the VHA Office of Rural Health and rural veterans. May. 2014 http://www.ruralhealth.va.gov/docs/factsheets/ORH_General_FactSheet_2014.pdf
20. Scott LA, Ben-Or K, Allen JK. Why are women missing from outpatient cardiac rehabilitation programs? A review of multilevel factors affecting referral, enrollment, and completion. *J Womens Health*. 2002; 11(9):773–91. DOI: 10.1089/15409990260430927
21. Samayoa L, Grace SL, Gravely S, Scott LB, Marzolini S, Colella TJ. Sex differences in cardiac rehabilitation enrollment: a meta-analysis. *Can J Cardiol*. 2014; 30(7):793–800. DOI: 10.1016/j.cjca.2013.11.007 [PubMed: 24726052]
22. Colella TJ, Gravely S, Marzolini S, Grace SL, Francis JA, Oh P, et al. Sex bias in referral of women to outpatient cardiac rehabilitation? A meta-analysis. *Eur J Prev Cardiol*. 2015; 22(4):423–41. DOI: 10.1177/2047487314520783 [PubMed: 24474091]
23. Klempfner R, Koifman E, Goldenberg I, Hamdan A, Tofler GH, Kopel E. The Israel nationwide heart failure survey: Sex differences in early and late mortality for hospitalized heart failure patients. *J Card Fail*. 2014; 20(3):193–8. DOI: 10.1016/j.cardfail.2013.12.016 [PubMed: 24374113]

24. Oosenbrug E, Marinho RP, Zhang J, Marzolini S, Colella TJ, Pakosh M, et al. Sex differences in cardiac rehabilitation adherence: A meta-analysis. *Can J Cardiol*. 2016; 32(11):1316–24. DOI: 10.1016/j.cjca.2016.01.036 [PubMed: 27129618]
25. Grace SL, Gravelly-Witte S, Kayaniyl S, Brual J, Suskin N, Stewart DE. A multisite examination of sex differences in cardiac rehabilitation barriers by participation status. *J Womens Health*. 2009; 18(2):209–16. DOI: 10.1089/jwh.2007.0753

Figure 1. Percent of participation in VA (N=66,710) vs. CMS (N=243,208) cardiac rehabilitation from 2007–2011

*Of the total unique number of VA patients discharged with HF over 5 years, 2.3% participated in CR. In Medicare recipients, 2.6% participated in CR.

Table 1

Factors independently associated with participation in cardiac rehabilitation among Veterans and patients with Medicare with heart failure

		VA (N=66,710)		CMS (N=243,208)	
		Odds Ratio (95% CI)	P-value	Odds Ratio (95% CI)	P-value
Age at discharge (per 10-year increase)		0.73 (0.70–0.78)	<0.0001	0.71 (0.69 – 0.72)	<0.0001
Gender	Female	1.0	--	1.0 (reference)	--
	Male	0.83 (0.58–1.20)	0.32	1.57 (1.49–1.65)	<0.0001
Race	White	1.0 (reference)	--	1.0 (reference)	--
	Black	0.88 (0.77–1.01)	0.07	0.36 (0.33–0.40)	<0.0001
	Asian/PI/AI*	1.10 (0.73–1.66)	0.65	0.45 (0.37–0.54)	<0.0001
	Hispanic	0.66 (0.49–0.89)	0.01	0.20 (0.15–0.28)	<0.0001
	Unknown	0.61 (0.48–0.77)	0.00	0.40 (0.17–0.99)	0.05
	Other	1.0 (reference)	--	N/A	--
Marital status	Unknown/Missing	1.21 (0.85–1.72)	0.28	N/A	--
	Married	1.17 (1.05–1.31)	0.00	N/A	--
Income	0–9,000	1.0 (reference)	--	N/A	--
	9,000–33,000	0.88 (0.78–1.01)	0.06	N/A	--
	>33,000	1.22 (1.04–1.43)	0.02	N/A	--
Hypertension		1.14 (0.95–1.37)	0.15	1.15 (0.97–1.35)	0.10
Ischemic heart disease		2.72 (2.35–3.15)	<0.0001	7.51 (6.53–8.62)	<0.0001
Hyperlipidemia		1.65 (1.45–1.89)	<0.0001	5.05 (4.52–5.63)	<0.0001
Diabetes		0.94 (0.84–1.05)	0.29	0.78 (0.74–0.82)	<0.0001
Cerebrovascular disease		0.87 (0.75–1.00)	0.05	0.74 (0.69–0.79)	<0.0001
Peripheral vascular disease		1.00 (0.88–1.12)	0.93	1.02 (0.97–1.08)	0.37
Chronic obstructive pulmonary disease		0.88 (0.79–1.98)	0.02	0.63 (0.60–0.67)	<0.0001
Chronic kidney disease		0.96 (0.86–1.07)	0.42	0.89 (0.84–0.93)	<0.0001
Depression		1.10 (0.98–1.24)	0.18	0.72 (0.68–0.77)	<0.0001

* Asian/Pacific Islander/American Indian