Published in final edited form as:

Indian Pediatr. 2017 March 15; 54(3): 185–192.

Intergenerational change in anthropometric indices of children and their predictors in New Delhi Birth Cohort

Sikha Sinha, M.Sc^{1,2} [Senior Research Fellow (Indian Council of Medical Research)], Abha Rani Aggarwal, Ph.D³ [Scientist-F], Clive Osmond, Ph.D⁴ [Senior Scientist & Professor of Biostatistics], Caroline H.D. Fall, DM⁴ [Professor of International Pediatric Epidemiology & Consultant in Child health], Santosh K. Bhargava, MD⁵ [Founder, New Delhi Birth Cohort], and Harshpal Singh Sachdev, MD¹ [Senior Consultant, Pediatrics and Clinical Epidemiology]

¹Sitaram Bhartia Institute of Science and Research, New Delhi, India

²University School of Medicine and Para-medical Health Sciences, Guru Gobind Singh Indraprastha University, Delhi, India

³National Institute of Medical Statistics, Indian Council of Medical Research, New Delhi, India

⁴MRC Lifecourse Epidemiology Unit, University of Southampton, Southampton, UK

⁵Sunder Lal Jain Hospital, New Delhi, India

Abstract

Objective—Evaluate intergenerational change in anthropometric indices of children and their predictors.

Design—Prospective cohort

Participants—New Delhi Birth Cohort subjects (F1), born between 1969 and 1972, were followed-up for anthropometry at birth and 6 monthly intervals until 21 years. Their children (F2) below 10 years were evaluated anthropometrically.

Outcome measure—Intergenerational change (F2-F1) in height, weight and body mass index (BMI) of children in comparison to their parents at corresponding ages.

Results—432 F2-F1 pairs were analysed in 0-5 (26.9%) and 5-10 (73.1%) years age groups. Children were considerably taller (0-5 years 0.99 SD; 5-10 years 1.17 SD) and heavier (0-5 years 0.77 SD; 5-10 years 1.52 SD) while only 5-10 years old were broader (had a higher BMI; 1.03 SD), than their parents. These increases for 0-5 and 5-10 years, respectively corresponded to 3.9 and 6.4 cm for height, 1.3 and 5.4 kg for weight and 0.2 and 1.9 kg/m² for BMI. Lower parents' anthropometric indices and poorer water supply and sanitation facilities; higher age of parents at

Correspondence to: Prof. Harshpal Singh Sachdev, Senior Consultant Pediatrics and Clinical Epidemiology, Sitaram Bhartia Institute of Science and Research, B-16 Qutab Institutional Area, New Delhi 110016, India. hpssachdev@gmail.com.

Author contributions

SS, ARA, HPS, CHDF and SKB conceptualised the study. SS, ARA, CO and HPS analyzed the data. SS drafted the initial manuscript. All authors contributed to the critical revision of the article.

Competing interests

The authors declare that they have no competing interests.

child birth and of children when measured (for height and weight); and more parental education (for weight and BMI), were associated with greater intergenerational gains in children.

Conclusion—Over one generation in an urban middle-class population, whose general living conditions had improved, under-five children have become considerably taller and heavier and 5-10 year old children have additionally become broader, than their parents at corresponding ages. Child populations probably “grow up” before “growing out”.

Keywords

Anthropometry; Body mass index; Intergenerational effect; Secular trend

Introduction

There is a perception that despite considerable economic growth, India has not made commensurate progress in addressing anthropometric undernutrition¹. The current national prevalence of undernutrition in children below five years is still high; 39% are stunted, 29% are underweight and 15% are wasted². However, this common belief appears somewhat flawed; there has been substantial improvement in stunting and underweight over the years^{3,4}. A comparison of the two latest national surveys, National Family Health Survey-3 (2005-06) and Rapid Survey on Children (2013-14), reveals a decline in stunting (48% to 39%) and underweight (43% to 29%) but only a marginal reduction in wasting (20% to 15%)^{2,5}. Similarly, available data from 14 states in the recent National Family Health Survey-4 (2015-16) confirms a decline in stunting and underweight. However, wasting decreased in only 6 states (by 2% to 15%) and paradoxically increased in 8 states (by 1% to 9%)⁶.

National surveys and regional cross sectional studies do not provide a robust indication of secular change and its quantum, especially in those reaping the benefits of development, for example, the middle socio-economic strata in urban settings. Information of this nature would provide a reasonable yardstick for improvement goals in childhood anthropometry, especially if novel nutrition interventions are being considered. Further, such data would identify the potential burden of over-nutrition in children born to relatively undernourished parents; a nutrition transition pattern that could explain the emerging epidemic of non-communicable diseases (NCDs) in the country.⁷

A robust estimate could be obtained through an intergenerational cohort design. This would offer the advantages of partially controlling for genetic variations while comparing anthropometry of children with their parents at similar ages. To the best of our knowledge, no such data are available in the Indian context to inform policy. This communication reports the intergenerational change in anthropometric indices of children below ten years of age and their predictors in the New Delhi Birth Cohort (NDBC).

Methods

The NDBC was drawn from a population of 119,799 living in 12 km² area of south Delhi during 1969-72^{8,9}. 20,755 married women of reproductive age were recruited and followed

regularly every other month to record menstrual dates. Information on the socio-demographic profile of the family were collected during recruitment by a social worker. These included maternal (*F0 generation*) schooling, paternal (*F0 generation*) occupation, and household socio-economic characteristics (type of family and house, and water supply and sanitation facilities). Women who became pregnant were visited every two months initially and on alternate days from the 37th week of gestation. There were 9169 pregnancies resulting in 8181 live births of cohort subjects (*F1 generation*). Trained personnel recorded the length and weight of the babies within 72 hours of birth, at the ages of 3, 6, 9 and 12 months (± 7 days) and every 6 months (± 15 days) until 14-21 years using standardised techniques⁹.

These F1 subjects were again followed-up in different adult ages namely 26-33, 33-39, 36-42 and 42-46 years for anthropometry and cardio-metabolic risk factors. Socio-demographic profile recorded during these visits included education and occupation of the the F1 subject, occupation of F1 spouse, type of housing, material possessions, family size, toilet, drinking water source and supply, and general water source and supply. Simultaneously, their children (*F2 generation*) below ten years of age were invited to attend the clinics for anthropometry. The study was approved by the respective institutional ethics committees. After obtaining informed consent from parents (F1) and assent from children (F2, >6 years of age), the length/height and weight of the F2 generation were recorded using standardised techniques¹⁰. Recumbent length in children below 2 years of age was measured using an infantometer, and height in children ≥ 2 years of age using a portable stadiometer to the nearest 0.1 cm. Weight was measured using calibrated digital weighing scales with a sensitivity of 0.1 kilogram. Body Mass Index (BMI) was calculated as weight in kilograms divided by height in meters squared.

Statistical Analysis

Data analysis was performed using SPSS 20.0. The intergenerational change in anthropometry was computed using two types of standardized scores (Z-scores); the World Health Organisation reference (WHO) and internally within the cohort. For the former, F2 and F1 generation anthropometry was converted to WHO length/height-for-age, weight-for-age and BMI-for-age Z-scores at the date of measurement, using an SPSS macro for age groups 0-5 years¹¹ and 5-19 years¹². For the internal cohort Z-scores, the longitudinal height, weight and BMI measurements for the F1 generation were modelled into growth charts using Cole's LMS method^{13,14}. These charts were used to compute age and sex specific Z-scores for F2 children at the date of measurement. To compare intergenerational anthropometry at similar ages, the F1 Z-score was interpolated at the exact age of the F2 measurement, using neighbouring F1 measurements. The interpolation was acceptable if the F1 observation was within 6 months for ages up to 1 year; within 1 year for ages 1 to 2 years; within 1.5 years for ages 2 to 3 years; and within 2 years thereafter. To estimate the comparison in absolute units, the interpolated F1 Z-score at the exact date of F2 measurement was back-transformed using the L, M and S values for the cohort Z-scores¹⁵.

The intergenerational change in size was computed as F2-F1. This intergenerational model had a 3-level data structure comprising the F0 generation having one or more F1 children

(F1-F1 siblings) and the F1 generation having one or more F2 children (F2-F2 siblings). Thus, a mixed model¹⁶ approach was used to allow for this structure. There were 13 sibships among the F1 generation. We fitted models to allow for this third level of variation, but there were too few sibships to disturb the intergenerational effect size estimates and their standard errors, so subsequently we ignored this variance component.

The change was quantified in two age groups: 0-5 years and 5-10 years, adjusting for all combinations of the sex of the child and parent. This intergenerational change was further adjusted for socio-demographic characteristics. These included maternal (F0) schooling, wealth and water supply, sanitation and hygiene (WASH) score at F1 birth; and education of the F1 parent (cohort subject), wealth and WASH score at F2 measurement. We used the 1st principal component score¹⁷ to derive the wealth and WASH scores at F1 birth and F2 measurement from the individual variables (listed in the footnotes of the relevant tables). These covariates were used in multivariate models as standardized units (mean 0), so that uncentred variables do not cause shifts in computed intergenerational changes.

Results

Among the 337 F1 subjects there were 13 pairs of F1-F1 siblings and among the 432 F2 children, 245 were single children, 89 were paired siblings and 3 were three siblings (Web Figure 1). We thus analysed 432 F2-F1 pairs, which included 178 (41.2%) father-son, 138 (31.9%) father-daughter, 71 (16.4%) mother-son and 45 (10.4%) mother-daughter comparisons. Among them, 116 (72 boys) were aged 0-5 years and 316 (177 boys) were 5-10 years old, with mean (SD) ages of 3.5 (1.2) and 7.9 (1.3) years, respectively. The mean (SD) height-for-age, weight-for-age and BMI-for-age (WHO Z-scores) were -0.97 (1.39), -0.68 (1.19) and -0.08 (1.37), respectively for younger children and -0.42 (1.22), -0.31 (1.55) and -0.12 (1.49), respectively for older children.

The socio-demographic characteristics at the time of the parent's (F1 generation) birth and at the time of the child's (F2 generation) measurement are summarized in Web Table 1. Comparatively, parents (F1) had a poorer socio-demographic profile; one-third were residing in a flat or bungalow, the mean (SD) members sharing a room were 3.6 (1.7), four-fifths had shared toilet facilities, 60% had a common water supply and only one-fourth of their F0 mothers had completed 10 or more years of education. In contrast, at the time of the child's measurement, almost all of them were living in either flat or independent house, mean (SD) members sharing a room were 1.9 (0.9), most had separate toilet facilities (96%) and water supply (81%), and 62% of their F1 mothers had 15 or more years of education (graduate).

Table 1 summarizes the inter-generational change (F2-F1) in anthropometry. Children were significantly ($P < 0.001$) taller and heavier than their parents at corresponding ages; the increase was similar (~ 1 SD) for height in both age groups but was higher for weight in older children (1.52 vs 0.77 WHO Z-score; non overlapping confidence intervals). However, the BMI-for-age increase (1 SD) was significant ($P < 0.001$) only in the 5-10 years age group. Considering the entire age range (0-10 years), the intergenerational increases (SD) in anthropometric indices (mean; 95% confidence intervals) were: Length/Height-for-age: 1.13 (1.01, 1.26), Weight-for-age: 1.32 (1.17, 1.47), and BMI-for-age: 0.79 (0.63, 0.95) ($P < 0.001$).

for all). These average Z-score increases for the 0-5 and 5-10 year age groups respectively corresponded to 3.9 and 6.4 cm for height, 1.3 and 5.4 kg for weight, and 0.2 and 1.9 kg/m² for BMI (Figure 1). The intergenerational change among boys and girls was comparable. Sensitivity analyses, using internal cohort Z scores and narrower window ranges for F1 Z-score interpolation at the exact age of F2 measurement revealed similar findings (data available on request).

Using the WHO definition¹⁸, in comparison to their parents, under-five children had a substantially lower prevalence of stunting (44.7% vs. 18.4%; n=114) and underweight (25.9% vs. 12.5%; n=112). However, wasting remained unchanged (4.6% vs. 5.5%; n= 109).

Table 2 depicts the multivariate associations of intergenerational change. Parents' (F1) anthropometry had the most consistent and strongest associations ($P < 0.001$), with effect sizes ranging from 0.47 to 0.65 SD decreases per SD. Thus children of shorter, lighter and thinner parents had gained more in height, weight and BMI, respectively. The age of the parent (F1) at child birth and the age of the child (F2) at measurement were significant ($P < 0.001$) positive predictors for F2 height-for-age and weight-for-age but not for BMI-for-age. Amongst the socio-demographics characteristics, poorer WASH status at the F1 parents' birth predicted greater increase in all three F2 indices. In contrast, higher parental (F1) education predicted greater F2 gain in weight and BMI, and had no association with F2 height gain. Web Table 2 quantifies the adjusted intergenerational changes across sub-groups of various predictors.

The scatterplot in Figure 2 illustrates the transition in BMI categories among parents and children at the same age. The cut-offs were slightly modified from WHO recommendations¹⁹ to ensure sufficient numbers in each category: Thin < -1.5 Z, Normal -1.5 to 0.5 Z, and Overweight > 0.5 Z. In comparison to 127 children, only 29 parents were overweight at the same age. While a higher proportion of children of overweight parents were overweight, the greatest increase in BMI categories occurred in thin parents. A quarter of their children were overweight while half were normal. In contrast, among all children born to overweight parents, only 41% were overweight and 59% were normal.

Discussion

Children below ten years were considerably taller (~ 1 SD) and heavier (0.8-1.5 SD), while only 5-10 years old were broader (~ 1 SD), than their parents at corresponding ages. Independent predictors of greater gains in children included parents with lower anthropometric indices and poorer WASH facilities; higher age of parents at child birth and of children at measurement (height and weight); and more parental education (weight and BMI). These data from an urban middle-class population provide robust confirmation of substantial increases in the body size of Indian children over one recent generation. The findings are in broad agreement with the observed decline in anthropometric undernutrition from national surveys²⁻⁶. Interestingly, the 26% reduction in stunting over 30 years in this small sample roughly corresponds to the 1% annual decline documented nationally²⁻⁶.

A scarcity of similar data precludes emphatic international comparisons. A recent analysis of a century of trends in adult human height (1896 to 1996 births) concluded that South Asians (including Indians) were among the shortest and had experienced little increase (<5 cm) during this era²⁰. We could locate only one directly comparable estimate from Britain, pertaining to a 15 years earlier period, with similar or better development index status than our cohort^{21,22}. At 7 years, these offspring were only slightly taller (0.19 SD or 1 cm) and broader (boys: 0.16 SD or 0.23 kg/m² and girls: 0.25 SD or 0.46 kg/m²) than their parents. The substantially larger intergenerational increases (~1 SD) seen in our cohort children will probably translate into commensurate or greater absolute (cm, kg, kg/m²) gains in adulthood. These findings should inspire optimism that with rapid improvements in living conditions, anthropometric gaps from high income countries will reduce, particularly because a plateauing trend for height is evident in some developed nations²⁰.

In the absence of targeted food or nutrient supplementation, it is not unreasonable to ascribe these intergenerational gains to improvements in general living conditions. However, from a policy perspective, filtering out important predictors is desirable. Poorer water supply and sanitation facilities of the parents (F1) at birth predicted greater F2 gains. A Cochrane systematic review of cluster randomized trials, intervening for only 9-12 months also suggests that WASH interventions confer a small benefit on linear growth (~0.1 SD) in under-five children²³. The current national impetus on “Swach Bharat Abhiyaan” is therefore timely and appropriate. In conformity with earlier experiences^{24,25}, higher parental literacy was a positive predictor, thereby re-emphasising the importance of improving education. Higher ages of parents at child birth and of children at measurement were important positive predictors of intergenerational gains in height and weight. A quadratic relation of childhood anthropometry with age of child birth has been documented in pooled analyses of cohorts from Low and Middle Income countries (LMICs)²⁶; our sample was probably constituted by the linear component of this association. However, both these variables are also proxies for exposure duration, thereby suggesting that sustained improvement in living conditions resulted in greater benefit.

Poorer parental anthropometry had the strongest and most consistent predictive ability for greater intergenerational gains. Potential explanations for this include statistical regression to the mean, narrowing of socio-economic inequalities and greater biological response among deprived strata. This observation augurs well for attempting equity for secular increases in the height of populations. However, excessive BMI gain in children of thin parents (Figure 2), if primarily due to increased adiposity, could be providing the backdrop for the current escalation of cardio-metabolic risk factors in Indian children and adolescents^{7,27,28}. This hypothesis is in concordance with our earlier observation of increased risk of diabetes mellitus in adults who were relatively thin as children but continued to become obese relative to themselves⁹.

Under-five children were considerably taller and heavier but only older subjects were additionally broader than their parents. This provides more direct evidence of the earlier postulate, based on cross-sectional comparisons²⁹, that children “grow up” (get taller) before “growing out” (get broader). This observation is also consistent with the increased

prevalence of obesity in older children from India²⁸ and abroad^{29–31}. The underlying mechanisms for this phenomenon are unclear to us and merit further exploration.

The main strengths of our study include the comparison of children and their parents at corresponding ages from carefully collected prospective data for the latter, an urban LMIC setting of relatively rapid socio-economic development, and appropriate multi-level modelling with available confounder adjustment. Important limitations include a somewhat small sample size based on a proportion of the currently available cohort. Among the parents, there was a considerably higher representation of fathers because outmigration after marriage was common in female cohort subjects (F1).

Implications for Research and Policy

Future research priorities could consider: (i) Confirmation in a larger sample, including adolescents; (ii) Validation of our findings in similar cohorts from other LMICs, if available; (iii) Mathematical modelling to predict expected anthropometric gains for shorter periods directly comparable to the periodicity of national surveys; and (iv) Evaluating relationships of intergenerational changes in anthropometric indices with children's cardio-metabolic risk factors and body composition.

Sustained improvement in general living conditions leads to considerable (~1 SD) increases in height, weight and BMI within one generation. The current governance focus on inclusive development is therefore apt, especially if the benefits percolate preferentially to the underprivileged. Isolated vertical interventions (for example, nutrient supplementation) should only be entertained if there is convincing evidence of substantial benefit above that expected from developmental transition. Findings from the NFHS-4 survey of notable reductions in stunting and underweight with nearly stagnant or even increased wasting prevalence in some states should not fuel exaggerated concerns and action to screen for and treat for severe acute malnutrition, as this phenomenon is expected for a population undergoing development-related anthropometric transition. Vigilance may be required to address the potential of greater cardio-metabolic risk in families showing large intergenerational increases in BMI.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Funding

We wish to thank the Indian Council of Medical Research for supporting Ms. Sikha Sinha through the Senior Research Fellowship Scheme. The original cohort studies were supported by the National Center for Health Statistics, USA and the Indian Council of Medical Research, India. Later phases have been supported by Department of Biotechnology, India; British Heart Foundation, UK; Wellcome Trust, UK; Medical Research Council; UK; Department for International Development, UK; and Bill and Melinda Gates Foundation, USA.

References

1. Paul VK, Sachdev HS, Mavalankar D, Ramachandran P, Sankar MJ, Bhandari N, et al. Reproductive health, and child health and nutrition in India: meeting the challenge. *Lancet*. 2011 Jan 22; 377(9762):332–49. [PubMed: 21227494]

2. [Accessed September 6th, 2016] Rapid Survey on Children 2013-2014. Available from: http://wcd.nic.in/issnip/National_Fact%20sheet_RSOC%20_02-07-2015.pdf
3. Sachdev HP. Overcoming challenges to accelerating linear growth in Indian children. *Indian Pediatr.* 2012 Apr; 49(4):271–5. [PubMed: 22565072]
4. Sachdev, HPS., Shah, D. Epidemiology of maternal and fetal malnutrition in South Asia. *Perinatal and Newborn Care in South Asia.* Bhutta, ZA., editor. Oxford: Oxford University Press; 2007. p. 75-105.
5. [Accessed September 6th, 2016] NFHS-3 Vol- I Summary of findings. [http://www.rchiips.org/nfhs/NFHS-3%20Data/VOL-1/Summary%20of%20Findings%20\(6868K\).pdf](http://www.rchiips.org/nfhs/NFHS-3%20Data/VOL-1/Summary%20of%20Findings%20(6868K).pdf)
6. [Accessed September 6th, 2016] NFHS 4 Factsheet. Available from: http://rchiips.org/NFHS/factsheet_NFHS-4.shtm
7. Sachdev, HPS. Exploring agricultural levers for mitigating the overnutrition burden in India. International Food Policy Research Institute Discussion Paper 01183; 2012 May. p. 1-29. <http://www.ifpri.org/publication/exploring-agricultural-levers-mitigating-overnutrition-burden-india> [Accessed September 6th, 2016]
8. Richter LM, Victora CG, Hallal PC, Adair LS, Bhargava SK, Fall CH, et al. COHORTS Group. Cohort profile: the Consortium of Health-Orientated Research in Transitioning Societies. *Int J Epidemiol.* 2012; 41:621–26. [PubMed: 21224276]
9. Bhargava SK, Sachdev HS, Fall CH, Osmond C, Lakshmy R, Barker DJ, et al. Relation of serial changes in childhood body-mass index to impaired glucose tolerance in young adulthood. *N Engl J Med.* 2004 Feb 26; 350(9):865–75. [PubMed: 14985484]
10. World Health Organization. Physical status: the use and interpretation of anthropometry. Report of a WHO Expert Committee Technical Report. 1995854
11. [Accessed September 6th, 2016] The WHO Child Growth Standards. <http://www.who.int/childgrowth/standards/en/>
12. [Accessed September 6th, 2016] WHO Growth Reference data for 5-19 years. <http://www.who.int/growthref/en/>
13. Cole TJ. The LMS method for constructing normalized growth standards. *Eur J Clin Nutr.* 1990 Jan; 44(1):45–60.
14. Pan H, Cole T. LMS chart maker Light version 2.54 A program for calculating age-related reference centiles using the LMS method.
15. Construction of LMS Parameters for the Centers for Disease Control and Prevention 2000 Growth Charts. *National Health Statistics Reports.* 2013 Feb 11.(63)
16. Cheng J, Edwards LJ, Maldonado-Molina MM, Komro KA, Muller KE. Real longitudinal data analysis for real people: building a good enough mixed model. *Stat Med.* 2010 Feb 20; 29(4):504–20. [PubMed: 20013937]
17. Vyas S, Kumaranayake L. Constructing socio-economic status indices: how to use principal components analysis. *Health Policy Plan.* 2006 Nov; 21(6):459–68. [PubMed: 17030551]
18. [Accessed September 6th, 2016] Global database on Child Growth and Malnutrition. <http://www.who.int/nutgrowthdb/about/introduction/en/index5.html>
19. [Accessed September 6th, 2016] WHO BMI-for-age criteria. http://www.who.int/growthref/who2007_bmi_for_age/en/
20. NCD Risk Factor Collaboration (NCD-RisC). A century of trends in adult human height. *Elife.* 2016 Jul 26.5 pii: e13410.
21. Li L, Power C. Influences on childhood height: comparing two generations in the 1958 British birth cohort. *Int J Epidemiol.* 2004 Dec; 33(6):1320–8. [PubMed: 15358746]
22. Pinot de Moira A, Power C, Li L. Changing influences on childhood obesity: a study of 2 generations of the 1958 British birth cohort. *Am J Epidemiol.* 2010 Jun 15; 171(12):1289–98. [PubMed: 20488872]
23. Dangour AD, Watson L, Cumming O, Boisson S, Che Y, Velleman Y, et al. Interventions to improve water quality and supply, sanitation and hygiene practices, and their effects on the nutritional status of children. *Cochrane Database Syst Rev.* 2013 Aug 1.(8):CD009382. [PubMed: 23904195]

24. Galobardes B, McCormack VA, McCarron P, Howe LD, Lynch J, Lawlor DA, Smith GD. Social inequalities in height: persisting differences today depend upon height of the parents. *PLoS One*. 2012; 7(1):e29118. [PubMed: 22238588]
25. [Accessed September 6th, 2016] UNICEF conceptual framework. <http://www.unicef.org/nutrition/training/2.5/4.html>
26. Fall CH, Sachdev HS, Osmond C, Restrepo-Mendez MC, Victora C, Martorell R, et al. COHORTS investigators. Association between maternal age at childbirth and child and adult outcomes in the offspring: a prospective study in five low-income and middle-income countries (COHORTS collaboration). *Lancet Glob Health*. 2015 Jul; 3(7):e366–77. [PubMed: 25999096]
27. Garg P, Kaur S, Gupta D, Osmond C, Lakshmy R, Sinha S, et al. Variability of thinness and its relation to cardio-metabolic risk factors using four body mass index references in school-children from Delhi, India. *Indian Pediatr*. 2013 Nov 8; 50(11):1025–32. [PubMed: 23798637]
28. Hoque ME, Doi SA, Mannan M, Long K, Niessen LW, Mamun AA. Prevalence of overweight and obesity among children and adolescents of the Indian subcontinent: a meta-analysis. *Nutr Rev*. 2014 Aug; 72(8):541–50. [PubMed: 25040110]
29. Smith SM, Craig LC, Raja AE, McNeill G, Turner SW. Growing up before growing out: secular trends in height, weight and obesity in 5–6- year- old children born between 1970 and 2006. *Arch Dis Child*. 2013 Apr; 98(4):269–73. [PubMed: 23413314]
30. Moon JS. Secular trends of body sizes in Korean children and adolescents: from 1965 to 2010. *Korean J Pediatr*. 2011 Nov; 54(11):436–42. [PubMed: 22253640]
31. Zong XN, Li H. Physical growth of children and adolescents in China over the past 35 years. *Bull World Health Organ*. 2014 Aug 1; 92(8):555–64. [PubMed: 25177070]

What is already known?

National surveys suggest a decline in anthropometric undernutrition but there is no robust quantification of secular increases in body size of children over a recent generation.

What this study adds?

Children below ten years were considerably taller (~1 SD) and heavier (0.8-1.5 SD), while only 5-10 year old children were broader (had a higher BMI; ~1 SD), than their parents at corresponding ages.

Figure 1. Intergenerational changes in anthropometric indices (WHO Z-scores)

Mixed model adjusted for the sex of the children and parents

Error bars represent 95% confidence intervals.

Figure 2. Scatterplot of transition in Body Mass Index (WHO Z Scores) in parents and their children at corresponding ages.

These categories were defined using WHO BMI Z-score as follows: Thin <-1.5, Normal -1.5 to 0.5, and Overweight >0.5.

Table 1
Intergenerational changes in anthropometry

Anthropometric change (F2-F1)	0-5 years			5-10 years		
	N	Mean (95% CI)	P value	N	Mean (95% CI)	P value
WHO Z-score						
Height for age	114	0.99 (0.73; 1.25)	<0.001	315	1.17 (1.03; 1.31)	<0.001
Weight for age	112	0.77 (0.53; 1.00)	<0.001	305 *	1.52 (1.34; 1.70)	<0.001
BMI for age	110	0.14 (-0.14; 0.42)	0.310	310	1.03 (0.85; 1.21)	<0.001
Absolute units						
Height (cm)	114	3.9 (2.9; 4.9)	<0.001	315	6.4 (5.6; 7.2)	<0.001
Weight (kg)	112	1.3 (0.8; 1.7)	<0.001	311	5.4 (4.7; 6.1)	<0.001
BMI (kg/m ²)	110	0.2 (-0.2; 0.6)	0.255	310	1.9 (1.5; 2.2)	<0.001

Mixed model adjusted for the sex of the children and parents

* WHO weight-for-age could not be interpolated for 6 parents because the reference stops at 10 years of age.

Table 2
Predictors of intergenerational change in anthropometric indices (multivariate model)

Predictors (Standardized units)	Height for age (N=331)	Weight for age (N=323)	BMI for age (N=325)
	Effect size (95% CI) ; P value	Effect size (95% CI) ; P value	Effect size (95% CI) ; P value
Parent anthropometry (WHO reference) @	-0.47 (-0.62, -0.31); <0.001	-0.49 (-0.67, -0.32); <0.001	-0.65 (-0.82, -0.49); <0.001
Age of parent at child birth	0.34 (0.18, 0.51); <0.001	0.37 (0.19, 0.56); <0.001	0.17 (-0.02, 0.36); 0.080
Age of child	0.25 (0.09, 0.40); 0.002	0.32 (0.14, 0.50); <0.001	0.12 (-0.08, 0.31); 0.246
At parent birth			
Maternal (F0) Schooling	0.01 (-0.17, 0.18); 0.954	-0.02 (-0.22, 0.19); 0.874	-0.02 (-0.23, 0.18); 0.816
Wealth, Housing, paternal (F0) occupation *	0.08 (-0.10, 0.25); 0.393	0.15 (-0.07, 0.36); 0.176	0.16 (-0.05, 0.37); 0.125
WASH (Sanitation & water supply) #	-0.17 (-0.32, -0.03); 0.021	-0.25 (-0.43, -0.08); 0.004	-0.22 (-0.39, -0.04); 0.014
At child measurement			
Education of parent (F1)	0.11 (-0.07, 0.29); 0.221	0.26 (0.05, 0.47); 0.015	0.29 (0.08, 0.50); 0.007
Wealth, Housing, maternal & paternal (F1) occupation §	0.08 (-0.11, 0.26); 0.421	0.10 (-0.12, 0.32); 0.374	0.05 (-0.17, 0.26); 0.672
WASH (Sanitation & water supply) ∇	0.00 (-0.14, 0.14); 0.969	-0.07 (-0.23, 0.09); 0.375	-0.10 (-0.26, 0.06); 0.236

Mixed model analysis adjusted for the sex of the children and parents. All other predictors were also inserted simultaneously in the multivariate model as standardized units.

The intraclass correlation coefficients (ICC) were 0.14 for height-for-age, 0.27 for weight-for-age and 0.20 for BMI-for-age.

@ Parental measurement and intergenerational change were for the same anthropometric index (for example, parent height-for-age for intergenerational change in height-for-age).

* Wealth score at F1 birth was generated from type of family, combination of type of housing and ownership, paternal (F0) occupation, per capita household annual income and crowding

WASH score at F1 birth was generated from type of toilet, water supply and facilities for sanitation and water supply

§ Wealth score at F2 measurement was generated using maternal and paternal (F1) occupation, material possession score and crowding

∇ WASH score at F2 measurement was generated from toilet and water supply (general and drinking water) facilities