Published in final edited form as:

J Pediatr Surg. 2017 June ; 52(6): 1026–1030. doi:10.1016/j.jpedsurg.2017.03.030.

Epithelial IGF1R is dispensable for IGF2 mediated enhanced intestinal adaptation in retinoblastoma deficient mice

Raphael C. Sun, Pamela M. Choi, Jose L. Diaz-Miron, Joshua Sommovilla, Jun Guo, Christopher R. Erwin, and Brad W. Warner

Division of Pediatric Surgery; St Louis Children's Hospital, Department of Surgery; Washington University School of Medicine, St. Louis, MO 63110, USA

Abstract

Purpose—Previously, we demonstrated enhanced adaptation after small bowel resection (SBR) in intestinal-specific Retinoblastoma (Rb) deficient mice along with elevated levels of insulin-like growth factor 2 (IGF2) expression within the villi. The purpose of this study was to verify that the insulin-like growth factor 1 receptor (IGF1R) plays a role in this phenomenon.

Methods—Inducible and intestinal specific Rb and IGF1R double knockout mice (iRb/IGF1R-IKO) (n=4) and Rb single knockout mice (iRb-IKO) (n=5) underwent 50% mid SBR. On post-operative day 28, mice were harvested, and structural adaptation was measured as changes in crypt depth and villus height. Rates of enterocyte proliferation were recorded. IGF2 expression within the remnant villi was measured via RT-PCR.

Results—Both iRb-IKO and iRb/IGF1R-IKO mice demonstrated enhanced adaptation with at least a 45% increase in both crypt depth and villus height in the proximal and distal remnant bowel. Both groups showed elevation of IGF2 expression in the remnant villi, but there were no differences between the two groups.

Conclusion—Epithelial IGF1R is dispensable for IGF2-mediated enhanced intestinal adaptation in retinoblastoma deficient mice. Our findings suggest that IGF2 signals for enhanced adaptation in cells outside of the epithelium. Further investigation is needed to study the IGF2/IGF1R signaling interaction within the mesenchyme.

Keywords

short gut syndrome; intestinal adaptation; insulin-like growth factor 1 receptor; insulin-like growth factor-2; retinoblastoma deletion

Corresponding Author: Brad W. Warner, M.D., St. Louis Children's Hospital, One Children's Place; Suite 5S40, St. Louis MO 63110, (314) 454-6022 – Phone, (314) 454-2442 –Fax, brad.warner@wustl.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Disclosures: Dr. Warner is on the Data Safety Monitoring Board of Shire Pharmaceuticals and a Scientific Advisory Board Member for Prolacta Bioscience.

INTRODUCTION

Short gut syndrome is a morbid clinical condition that follows a massive small bowel resection (SBR) required to treat several pediatric conditions such as necrotizing enterocolitis, midgut volvulus, and intestinal atresias (1). Intestinal adaptation is the response to massive SBR and is characterized by structural increases in crypt depth and villus height resulting in increased absorptive and digestive mucosal surface area (2, 3). The exact mechanisms for adaptation are incompletely understood and continues to be an area of research with the intent to achieve enteral autonomy in short gut patients.

Insulin-like growth factor 2 (IGF2) is a recognized intestinotrophic factor which binds and signals through insulin-like growth factor 1 receptor (IGF1R) (4). Studies have suggested that infusion of such intestinotrophic factors have an exaggerated adaptation response (5–10). Previously, our lab demonstrated that disrupting the expression of retinoblastoma protein (Rb) within the enterocytes of mice resulted in taller villi (11). Furthermore, we discovered that IGF2 expression was significantly elevated within the villi of the Rb-null mice (12, 13). The deletion of enterocyte Rb in the setting of SBR resulted in even taller villi and deeper crypts thereby enhancing adaptation. This phenomenon of enhanced adaptation was prevented when IGF2 expression was deleted. Therefore, IGF2 is considered to be a necessary intestinotrophic factor for Rb mediated enhanced adaptation after SBR (13).

We have previously demonstrated that disrupting the expression of growth factor receptors such as IGF1R within the intestinal epithelium has no effect on the degree of intestinal adaptation after SBR (14) (15). We also learned that IGF2 by itself is not required for normal adaptation to occur. These findings suggest that IGF1R may play a more important role in the sub-epithelial compartments such as muscularis or perhaps that there may be other compensatory mechanisms which cause other pathways to be upregulated during the process of intestinal adaptation when the expression of an epithelial-targeted growth factor receptor is knocked out.

Given our previous findings, we sought to determine whether IGF1R within the enterocytes is necessary for enhanced adaptation to occur after SBR in the setting of Rb deletion knowing that IGF2 is elevated and the driving force behind Rb-mediated adaptive growth. To investigate this, we crossed Rb (flox/flox) and IGF1R (flox/flox) mice with the Villin-Cre ER mouse line to create an Rb/IGF1R double knockout mouse line.

MATERIALS AND METHODS

All of our protocols and experiments were approved by the Washington University Animals Studies Committee (Protocol #20130038) and strictly followed National Institutes of Health (NIH) animal care guidelines.

Animals

Previously, we obtained mice with a tamoxifen-inducible Cre-fusion protein with a villin promoter (Sylvie Robine, Curie institute, Paris, France). We obtained mice with floxed Rb gene and IGF1R gene (Jackson Laboratories, Bar Harbor, ME). We crossed (Villin Cr-

ER(+); Rb(f/f)) with (Villin Cr-ER(+); IGF1R(f/f)) mice to create an inducible, intestinal epithelial-specific Rb and IGF1R double knockout mice (iRb/IGF1R-IKO). Double knockout iRb/IGF1R-IKO mice (n=4) were induced by injecting tamoxifen (0.5 mg/day; Sigma, St. Louis, Missouri) intraperitoneally for 3 days after small bowel resection. Intestinal specific single knockout mice (iRb-IKO) (n=5) served as controls. Tamoxifen was prepared as previously described (16). Mice were kept in the approved animal holding area with a 12 hour light-dark schedule and given rodent chow ad lib after weaning.

Small bowel resection

All mice underwent 50% mid small bowel resection (SBR) in a manner similar to our previously described proximal SBR (2). A mid-SBR was performed in these mice to determine whether there was an aboral gradient of IGF2 expression along the bowel. The intestinal resections were performed by transecting the bowel approximately 8 cm distal from the ligament of Treitz and at 8 cm proximal from the cecum. The resected segment was approximately an average of 12cm in length. After this segment of bowel was removed, intestinal continuity was reestablished by an end-to-end primary anastomosis using interrupted 9-0 monofilament sutures. A 2 cm proximal and distal segment of the resected ileum was fixed in 10% neutral-buffered formalin for baseline histology. All mice were fed a liquid rodent diet post operatively. (Microstabilized Rodent Liquid Diet LD101, Purina Mills, St Louis, MO)

Experimental design

Both proximal and distal portions of the resected bowel was collected at the time of SBR as a baseline control. On post-operative day (POD) 28, the remnant bowel was removed. Histology was collected to compare adaptive structural changes within the crypts and villi against preoperative baseline measurements. Enterocytes were isolated and used for protein analysis to confirm gene deletion and IGF2 mRNA levels.

After 28 days, structural adaptation was assessed by measuring changes in crypt depth and villus height on both the proximal and distal remnant bowel. Intestinal tissue was cut into two sections each, 50µm apart followed by H&E staining. Paraffin-embedded tissue sections were also immunohistochemically stained with p-histone 3. Rates of enterocyte proliferation were measured and calculated to a proliferation index by counting the number of p-histone 3 positive cells divided by total number of cells per crypt. A minimum of 20 crypts were counted. All histology slides were measured using MetaMorph Computer program (Molecular Devices, Downingtown PA). Western blots were used to confirm Rb and IGF1R deletion within the enterocytes. RT-PCR was used to measure the relative level of IGF2 within the villi.

Harvest and enterocyte isolation

The remnant intestine was harvested on POD 28 as previously reported (17). Briefly, a midline laparotomy was performed and the entire small intestine was flushed with ice-cold phosphate buffered saline containing protease inhibitors (0.2nM phenylmethanesulfonyl fluoride, 5µg/mL aprotinin, 1µM benzamidine, 1mM sodium orthovanadate, and 2µM cantharidin; EMD, Gibbstown, NJ). A 2 cm segment of bowel proximal and distal to the

anastomosis was fixed in 10% neutral-buffered formalin to compare with intraoperative histology collected at the time of resection. The remainder of the remnant bowel was used to isolate crypt and villus enterocytes as we have previously described (17). Protein was isolated for further analysis to verify double knockout. RT-PCR was used to analyze the degree of IGF2 expression present within the villus enterocytes.

Western blotting

Villus and crypt samples were isolated and lysate was prepared as previously described (15). Protein concentration was determined using RC DC kit (Bio-Rad; Hercules, CA). Proteins were loaded into a gel in equal concentration amounts. Antibodies used in this study included Rb, IGF1R, and lamin B1 (Cell Signaling Technology; Danvers MA). The proteins were detected using Bio-Rad ChemiDoc™ XRS+ system with image Lab™ software (Bio-Rad, Hercules CA).

Real-time quantitative PCR

RNA was prepared from harvested crypts and villi as previously described (17) and were homogenized in lysis buffer (RNAqueous kit, Ambion, Austin TX). The RNA was extracted according to kit instructions and stored at -80°C . Total RNA concentration was determined using a NanoDrop Spectrophotometer (ND-1000, NanoDrop Technologies, Wilmington DE). IGF-2 gene expression was determined using primers and reagents from Applied Biosystems and β -Actin was used as the endogenous control. Applied Biosystems 7500 Fast Real-Time PCR system (Applied Biosystems, Foster City CA) was used to obtain relative RNA expression.

Statistics

Statistical analysis was performed using Student's t-test to compare the two experimental groups. A p-value < 0.05 was considered significant.

RESULTS

Confirmation of enterocyte IGF1R and Rb knockout

The successful deletion of IGF1R and Rb protein in the small intestine was confirmed by Western blot (Figure 1). Rb knockout was also confirmed using RT-PCR to evaluate for mRNA relative levels (data not shown).

Structural adaptation

When subjected to 50% mid SBR, both iRb-IKO and iRb/IGF1R-IKO mice demonstrated enhanced adaptation with a minimum 45% increase in both crypt depth and villus height in both proximal and distal segments of the remnant bowel (Figure 2, 3a,b).

There was no statistical difference in the degree of adaptation between the iRB-IKO and iRB/IGF1R-IKO mice in either proximal or distal remnant intestine. When looking at the proximal bowel, the iRB-IKO mice had an average of $82.3 \pm 6.6\%$ increase in the crypt depth and an average of $69.6 \pm 12.6\%$ increase in the villus height after SBR while the iRb/IGF1R-

IKO had an average of $90.8 \pm 17.0\%$ and $71.2 \pm 10.3\%$ in the crypt depth and villus height respectively.

When examining the distal bowel, the iRB-IKO mice had an average of $59.5 \pm 8.9\%$ increase in the crypt depth and an average of $49.2 \pm 5.4\%$ increase in the villus height after SBR while the iRb/IGF1R-IKO had an average of $67.4 \pm 8.6\%$ and $51.1 \pm 10.3\%$ in the crypt depth and villus height respectively.

Enterocyte proliferation rates

Rates of proliferation in both iRb/IGF1R-IKO and iRB-IKO mice were similar and statistically insignificant following SBR. iRb/IGF1R-IKO had an average proliferative index of 0.34 ± 0.02 and the iRB-IKO mice had an average of 0.31 ± 0.04 ($p=0.1$).

IGF2 mRNA expression

Total RNA was isolated from villus enterocytes on POD28. Previously our lab demonstrated a two fold increase of IGF2 mRNA expression in the remnant bowel between iRb-IKO and WT littermates after proximal SBR. Here we demonstrate that IGF1R deficiency had no effects on IGF2 mRNA expression in either the proximal and distal remnant bowel after mid (Figure 4).

DISCUSSION

In our mouse SBR model, we successfully deleted both Rb and IGF1R within the intestinal epithelium. By doing this, we induced elevated IGF2 mRNA expression levels in both the proximal and distal remnant bowels after a 50% mid SBR. Using Rb mediated enhanced adaptation in the single knockout iRb-IKO mice as a control, we found that the double knockout (iRb/IGF1R-IKO) had an equivalent increase in adaptation within the remnant proximal and distal remnant bowel. The magnitude of adaptation in either group of mice was enhanced compared with normal resection-induced adaptation (2). Although IGF1R was deleted within the enterocytes in this mouse model, proliferation and structural adaptation was not affected and was not required for Rb-mediated enhanced adaptation.

IGF2 is a known intestinotrophic factor and binds to IGF1R (18). Systemic administration of IGF1, another growth factor which binds to IGF1R has previously been shown to amplify adaptive intestinal mucosal growth after SBR (19) (10). Normal adaptation after SBR still occurs when IGF1R expression was disrupted within the enterocytes (14). Further, blocking expression of both IGF1R or epidermal growth factor receptor within the epithelial compartment does not appear to affect structural adaptation or cell proliferation (15). In this experiment, not only were we able to delete IGF1R but we demonstrated compensatory increases in IGF2 mRNA levels in the context of Rb deletion. This would lead us to believe IGF2 has an effect outside of the epithelium and can bind to IGF1R or other receptors within the subepithelial cells of the bowel wall to mediate downstream signaling and resection-induced intestinal growth.

Independent of IGF1R, stimulation of the insulin receptor with insulin has been demonstrated to enhance resection-induced intestinal adaptation (20–22). Previous reports

have shown that IGF2 is able to bind to insulin receptors and may therefore provide compensatory activity for the lack of IGF1R downstream signaling within enterocytes (23).

The reason we performed a 50% mid SBR was to see the deleterious effects of Rb knockout in both the proximal and distal bowel. At baseline, both unoperated iRB-IKO and iRB/IGF1R-IKO mice had a gradual decrease of IGF2 mRNA expression levels from the proximal to the distal end with the highest expression at the jejunum. After mid SBR, relative mRNA levels of IGF2 increased compared to their baseline IGF2 levels within the villi.

We observed a difference between the proximal and distal bowel in both IGF2 levels and degree of structural adaptation after mid SBR. In our previous findings, we showed a significant positive linear relationship between the IGF2 mRNA expression and the percent increase of villus height after SBR (13). Here, we reveal a similar relationship with the proximal remnant bowel having higher relative IGF2 mRNA expression levels compared to the distal remnant bowel. The degree of adaptation for the proximal villus height was nearly 70% increase in both the iRB-IKO and iRB/IGF1R-IKO mice compared to the distal remnant bowel which was approximately a 50% increase.

It must also be considered that since IGF2 is normally higher in the jejunum compared to the ileum may contribute to why villus height is greater in the jejunum compared with the ileum. The degree of IGF2 expression may have a correlation with intestinal growth at baseline and this mechanism should also be further investigated to understand the relationship between growth factors and structural development.

We have previously demonstrated Rb deletion within the intestinal epithelium prior to SBR does not produce enhanced adaptation (24). This supports the observation that timing of the deletion of Rb plays a role in the additive degree of adaptation. Future experiments would be required to understand the temporal profile and effects of RB deletion and SBR.

In summary, IGF2 expression in both the proximal and distal remnant bowel is necessary for Rb-mediated enhanced adaptation after 50 % mid SBR. This phenotype was consistent despite absent IGF1R within the intestinal epithelium. These findings suggest that Rb-mediated enhanced adaptation and elevated IGF2 expression signals either via the epithelial insulin receptor, or IGF1R in other cells of the intestinal wall. This is an important physiologic response and our new findings provide new mechanistic insights into the important process of intestinal adaptation. Further investigation of this intestinotrophic signaling pathways in other cell compartments outside of the epithelium will be important areas of further investigation designed to better therapeutic interventions for patients with short gut syndrome patients.

Acknowledgments

This work was supported by National Institutes of Health Grants R01 DK 059288 (Warner), P30DK52574–Morphology and Murine Models Cores of the Digestive Diseases Research Core Center of the Washington University School of Medicine, and The Children's Surgical Sciences Research Institute of the St. Louis Children's Hospital Foundation. Dr. Sun was also supported by a Research Fellowship Award through the Association for Academic Surgery Foundation.

References

1. Goulet O, Ruemmele F. Causes and management of intestinal failure in children. *Gastroenterology*. 2006; 130(2)(Suppl 1):S16–28. [PubMed: 16473066]
2. Helmrath MA, VanderKolk WE, Can G, Erwin CR, Warner BW. Intestinal adaptation following massive small bowel resection in the mouse. *Journal of the American College of Surgeons*. 1996; 183(5):441–9. [PubMed: 8912612]
3. Rubin DC, Levin MS. Mechanisms of intestinal adaptation. *Best practice & research Clinical gastroenterology*. 2016; 30(2):237–48. [PubMed: 27086888]
4. Conlon MA, Francis GL, Tomas FM, Wallace JC, Howarth GS, Ballard FJ. Continuous 14 day infusion of IGF-II increases the growth of normal female rats, but exhibits a lower potency than IGF-I. *The Journal of endocrinology*. 1995; 144(1):91–8. [PubMed: 7534330]
5. Shin CE, Helmrath MA, Falcone RA Jr, Fox JW, Duane KR, Erwin CR, et al. Epidermal growth factor augments adaptation following small bowel resection: optimal dosage, route, and timing of administration. *The Journal of surgical research*. 1998; 77(1):11–6. [PubMed: 9698525]
6. Chaet MS, Arya G, Ziegler MM, Warner BW. Epidermal growth factor enhances intestinal adaptation after massive small bowel resection. *Journal of pediatric surgery*. 1994; 29(8):1035–8. discussion 8–9. [PubMed: 7965502]
7. Burrin DG, Wester TJ, Davis TA, Amick S, Heath JP. Orally administered IGF-I increases intestinal mucosal growth in formula-fed neonatal pigs. *The American journal of physiology*. 1996; 270(5 Pt 2):R1085–91. [PubMed: 8928910]
8. Lemmey AB, Ballard FJ, Martin AA, Tomas FM, Howarth GS, Read LC. Treatment with IGF-I peptides improves function of the remnant gut following small bowel resection in rats. *Growth factors*. 1994; 10(4):243–52. [PubMed: 7803042]
9. Lemmey AB, Martin AA, Read LC, Tomas FM, Owens PC, Ballard FJ. IGF-I and the truncated analogue des-(1-3)IGF-I enhance growth in rats after gut resection. *The American journal of physiology*. 1991; 260(2 Pt 1):E213–9. [PubMed: 1996625]
10. Dahly EM, Guo Z, Ney DM. IGF-I augments resection-induced mucosal hyperplasia by altering enterocyte kinetics. *American journal of physiology Regulatory, integrative and comparative physiology*. 2003; 285(4):R800–8.
11. Choi PM, Guo J, Erwin CR, Wandu WS, Leinicke JA, Xie Y, et al. Disruption of retinoblastoma protein expression in the intestinal epithelium impairs lipid absorption. *American journal of physiology Gastrointestinal and liver physiology*. 2014; 306(10):G909–15. [PubMed: 24742992]
12. Choi P, Guo J, Erwin CR, Warner BW. IGF-2 mediates intestinal mucosal hyperplasia in retinoblastoma protein (Rb)-deficient mice. *Journal of pediatric surgery*. 2013; 48(6):1340–7. [PubMed: 23845628]
13. Choi PM, Sun RC, Sommovilla J, Diaz-Miron J, Guo J, Erwin CR, et al. IGF-2 Is Necessary for Retinoblastoma-Mediated Enhanced Adaptation after Small-Bowel Resection. *Journal of gastrointestinal surgery : official journal of the Society for Surgery of the Alimentary Tract*. 2014
14. Sun RC, Choi PM, Guo J, Erwin CR, Warner BW. Insulin-like growth factor 2 and its enterocyte receptor are not required for adaptation in response to massive small bowel resection. *Journal of pediatric surgery*. 2014; 49(6):966–70. ;discussion 70. [PubMed: 24888844]
15. Sun RC, Diaz-Miron JL, Choi PM, Sommovilla J, Guo J, Erwin CR, et al. Both epidermal growth factor and insulin-like growth factor receptors are dispensable for structural intestinal adaptation. *Journal of pediatric surgery*. 2015; 50(6):943–7. [PubMed: 25818318]
16. Metzger D, Chambon P. Site- and time-specific gene targeting in the mouse. *Methods*. 2001; 24(1): 71–80. [PubMed: 11327805]
17. Guo J, Longshore S, Nair R, Warner BW. Retinoblastoma protein (pRb), but not p107 or p130, is required for maintenance of enterocyte quiescence and differentiation in small intestine. *The Journal of biological chemistry*. 2009; 284(1):134–40. [PubMed: 18981186]
18. Lund PK. Molecular basis of intestinal adaptation: the role of the insulin-like growth factor system. *Annals of the New York Academy of Sciences*. 1998; 859:18–36. [PubMed: 9928367]

19. Bortvedt SF, Lund PK. Insulin-like growth factor 1: common mediator of multiple enterotrophic hormones and growth factors. *Current opinion in gastroenterology*. 2012; 28(2):89–98. [PubMed: 22241077]
20. Shamir R, Kolacek S, Koletzko S, Tavori I, Bader D, Litmanovitz I, et al. Oral insulin supplementation in paediatric short bowel disease: a pilot observational study. *Journal of pediatric gastroenterology and nutrition*. 2009; 49(1):108–11. [PubMed: 19503001]
21. Ben Lulu S, Coran AG, Mogilner JG, Shaoul R, Shamir R, Shehadeh N, et al. Oral insulin stimulates intestinal epithelial cell turnover in correlation with insulin-receptor expression along the villus-crypt axis in a rat model of short bowel syndrome. *Pediatric surgery international*. 2010; 26(1):37–44. [PubMed: 19847442]
22. Ben Lulu S, Coran AG, Shehadeh N, Shamir R, Mogilner JG, Sukhotnik I. Oral insulin stimulates intestinal epithelial cell turnover following massive small bowel resection in a rat and a cell culture model. *Pediatric surgery international*. 2012; 28(2):179–87. [PubMed: 21986618]
23. Blakesley VA, Scrimgeour A, Esposito D, Le Roith D. Signaling via the insulin-like growth factor-I receptor: does it differ from insulin receptor signaling? *Cytokine & growth factor reviews*. 1996; 7(2):153–9. [PubMed: 8899293]
24. Leinicke JA, Longshore S, Wakeman D, Guo J, Warner BW. Regulation of retinoblastoma protein (Rb) by p21 is critical for adaptation to massive small bowel resection. *Journal of gastrointestinal surgery : official journal of the Society for Surgery of the Alimentary Tract*. 2012; 16(1):148–55. discussion 55. [PubMed: 22042567]

Figure 1.

Western Blot confirming absent retinoblastoma (Rb) and insulin-like growth factor-1 receptor (IGF1R) within enterocytes in induced Rb/IGF1R- intestinal knockout (iRb/IGF1R-IKO) mice. Lamin B1 was used as loading control.

Figure 2.

Percent increase in crypt depth and villus height at 28 days following a 50% mid-small bowel resection of both proximal and distal remnant bowel in induced retinoblastoma-intestinal knockout (iRb-IKO) and iRb/insulin-like growth factor-1 receptor double knockout (iRb/IGF1R-IKO) mice. There were no significant differences between the two groups of mice at any location.

Figure 3.

a: Photomicrograph of resected proximal small bowel on the left compared to proximal small bowel at 28 days following a 50% mid-small bowel resection demonstrating enhanced intestinal adaptation (taller villi, deeper crypts).

b. Photomicrograph of resected distal small bowel on the left compared to distal small bowel at 28 days following a 50% mid-small bowel resection demonstrating enhanced intestinal adaptation (taller villi, deeper crypts).

Figure 4.

Relative IGF2 mRNA expression in induced retinoblastoma – intestinal knockout (iRb-IKO) and iRb/insulin-like growth factor-1 receptor double knockout (iRB/IGF1R-IKO) mice at 28 days following a 50% mid-small bowel resection. Both proximal and distal remnant intestine for iRb-IKO and iRB/IGF1R-IKO mice are shown. Beta-actin was used as endogenous control.