Published in final edited form as:

Org Lett. 2008 October 02; 10(19): 4383–4386. doi:10.1021/ol801860s.

Titanocene(III)-Catalyzed Formation of Indolines and Azaindolines

Peter Wipf and John P. Maciejewski

Department of Chemistry, University of Pittsburgh, Pittsburgh, Pennsylvania 15260

Abstract

Reductive cyclization of epoxides tethered to substituted anilines and aminopyridines in the presence of 3 mol % of titanocene(III) chloride and stoichiometric manganese metal promotes a radical annulation to form 3,3-disubstituted indolines and azaindolines.

Graphical abstract

The continued development of new methodology for the construction of indolines^{1,2} and azaindolines³ is well justified by the tremendous therapeutic potential associated with these heterocyclic building blocks.⁴ Natural products containing the 3,3-disubstituted indoline motif include the clinically used antitumor agent vinblastine (**1**),⁵ the novel marine metabolite diazonamide A (**2**),⁶ and the cage-like structures of kopsidarine (**3**),⁷ and scholarisine A (**4**)⁸ (Figure 1).

Alkyl radical cyclization onto pyridine rings to prepare azaindolines is common;⁹ in contrast, methods to construct indolines by alkyl radical cyclization onto the aromatic nucleus without using xanthate transfer are scarce.^{2b,c} While the vast majority of known protocols for indoline synthesis utilize nucleophilic, Lewis acid and transition metal cross-coupling techniques, we envisioned a radical process that installs a quaternary carbon simultaneous with pyrrolidine formation from readily available epoxidized allylic amines.¹⁰ The reductive opening of epoxides using either stoichiometric¹¹ or catalytic¹² titanocene(III) chloride was particularly attractive toward this goal. Nearly two decades after the first reports of using *in situ* prepared titanocene(III) chloride to generate reactive radical intermediates from epoxides, novel applications of this reagent continue to emerge.¹³

To test the potential of this methodology in the synthesis of 3,3-disubstituted indoline and azaindoline heterocycles, epoxide **4a**¹⁴ was subjected to 10 mol % of Cp_2TiCl_2 at room

temperature in degassed THF (0.1 M) in the presence of 2 equiv of zinc dust. However, these conditions gave only traces of the desired indoline and mainly undesired side products. Fortunately, upon lowering the concentration of the substrate to 0.03 M and using 10 mol % of precatalyst in the presence of 0.80 equiv of 20–50 mesh manganese powder under sonication¹⁵ at room temperature, a 3:1 ratio of indoline **5a** and tetrahydroquinoline **6a** was detected by GC analysis of the crude reaction mixture (Table 1, entry 1). Sub-stoichiometric amounts of manganese metal increased the ratio in favor of indoline **5a**; a possible indication of reversible radical pathways.^{12c} Addition of a methyl group to the epoxide in substrate **4b** ($R^2=CH_3$) improved the efficiency of the titanocene(III) chloride catalyzed process and afforded **5b** as a single product in 82% yield (entry 2). With 1,1-disubstituted epoxides, sonication was not essential and conventional magnetic stirring also afforded the product in good yield (entry 3).

Model substrates **4a** and **4b** were designed to facilitate the cyclization by the presence of two symmetrical *N*-phenyl substituents. However, in order to broaden the scope of this reaction, we intended to replace one of them with a suitable nitrogen protective group. For reasons that are not completely clear, the secondary amine ($R^1=H$) mostly decomposed in the reaction mixture, and only a minor amount of reduced amino alcohol was formed. Alternatively, among protecting groups at this position, including the *p*-toluenesulfonyl, benzyl, trifluoroacetyl, and *t*-butoxycarbonyl functions, the benzylcarbamate (Cbz) group proved to be the most versatile substituent after we re-optimized the reaction parameters. At room temperature, anilide **4c** afforded a ~2:1 mixture of **5c**:**7c** at 0.1 M concentration in the presence of 10 mol % of titanocene (Table 1, entry 4). The undesired reduced epoxide **7c** could be suppressed by lowering the precatalyst loading to 3 mol % of Cp_2TiCl_2 while increasing the reaction temperature to THF at reflux.¹⁶ Under these conditions, indoline **8c** was isolated in 63% yield over 2 steps (entry 5). When these optimized conditions were applied to the earlier model system **5a**, indoline **5b** was formed in 89% yield (entry 6). High yields were also obtained for epoxide **4a**, which provided a 3.8:1 mixture of **5a** and **6a** in 87% yield. We also explored the sensitivity of the conversion to air. When the epoxide-opening rearrangement with substrate **4c** was performed in a flask kept open to the atmosphere in non-degassed THF, the reactivity was greatly diminished (entry 8). Additional control experiments using either solely the precatalyst (entry 9) or the manganese metal (entry 10) under otherwise identical reaction conditions failed to afford indoline products.

The general scope of indoline formation was further illustrated by the conversions summarized in Table 2. Alkyl-substituted substrates, including tetrahydroquinoline **15** afforded the corresponding indolines in modest to good yields (entries 2–4). The electron-rich 5-methoxyindoline **18** was isolated in low yield (21% over 2 steps). Electron-deficient substrates underwent the epoxide-opening rearrangement to afford substituted indolines in good yields (entries 6 and 7). However, an attempt to prepare the tricyclic pyrroloindole **24** from epoxide **23** did not yield any of the cyclized product (entry 8). *Meta*-substituted anilines generally led to 1:1 mixtures of regioisomeric indolines (data not shown).

Azaindoles are attractive bioisosteres of indoles in pharmaceutical research.¹⁷ We therefore investigated the possibility of applying this methodology toward the formation of azaindolines, using aminopyridines in place of anilines. Chemoselective epoxidation of

alkenes in the presence of a pyridine ring isprecedented, however, attempts to efficiently prepare the epoxide on an unsubstituted aminopyridine substrate were unsuccessful. This was primarily due to the reactivity of the nucleophilic pyridine nitrogen toward either *m*-CPBA or DMDO.¹⁸ In contrast, an *ortho*-chlorine substitution¹⁹ proved to be sufficient to attenuate the nucleophilic character of the pyridine nitrogen. Curtius rearrangement of the known carboxylic acid **25**²⁰ and trapping of the intermediate isocyanate with benzyl alcohol afforded aminopyridine **26** (Scheme 1). Subsequent methallylation of the Cbz-protected amine and epoxidation with *m*-CPBA led to epoxide **27**. Under the optimized titanocene(III) chloride conditions, cyclization provided an intermediate 4,6-dichloro-5-azaindoline, which was concurrently deprotected and dechlorinated with Pd/C under an atmosphere of H₂ to give azaindoline **28** in 52% yield over 2 steps.

As a mechanism for the titanocene(III) chloride catalyzed epoxide-opening annulation, we propose that the formation of the β -titanoxy radical **30**^{11,12} is followed by a reversible cyclization onto the aromatic ring, forming the cyclohexadienyl radical intermediate **31** (Figure 2).²¹ Oxidation of the dienyl radical by trace amounts of dioxygen and proton loss affords the indoline **32**.²² Furthermore, protodemetalation by collidinium hydrochloride leads to product **33**, and presumably regenerates the precatalyst Cp₂TiCl₂.²³

The titanocene(III) chloride catalyst or the manganese byproducts may also serve as a Lewis acid to promote an epoxide-opening Friedel-Crafts alkylation. However, 5-membered ring benzene annulations by epoxide openings under Friedel-Crafts conditions are quite inefficient,²⁴ and a stoichiometric amount of reducing agent is required to drive the titanocene(III) chloride catalyzed process to completion.²³ Furthermore, it has previously been shown that the MnCl₂ produced in the reaction is not Lewis-acidic enough to promote epoxide-opening reactions.^{12a}

In summary, we have developed a novel titanocene(III) chloride catalyzed epoxide-opening arene annulation that affords 3,3-disubstituted indolines and tolerates a range of substituents on the aromatic ring. This methodology can be extended toward other five-membered heterocycles, as demonstrated by the preparation of a 3,3-disubstituted 5-azaindoline.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work has been supported by the NIH/NIGMS CMLD program (GM067082), and, in part, by R01-GM55433.

References

1. For a review on indoline synthesis, see: Gribble GW. J Chem Soc, Perkin Trans I. 2000:1045.
2. (a) Clive DLJ, Sunasee R. Org Lett. 2007; 9:2677. [PubMed: 17559217] For indoline preparation using xanthate transfer, see: (b) Zard, SZ. Radical Reactions in Organic Synthesis. Oxford University Press Inc; New York: 2003. p. 153-163. (c) Ly TM, Quiclet-Sire B, Sortais B, Zard SZ. Tetrahedron Lett. 1999; 40:2533.

3. For a review on azaindoline synthesis see: Song JJ, Reeves JT, Gallou F, Tan Z, Yee NK, Senanayake CH. *Chem Soc Rev.* 2007; 36:1120. [PubMed: 17576479]
4. Sun H, Ehlhardt WJ, Kulanthaivel P, Lanza DL, Reilly CA, Yost GS. *J Pharmacol Exp Ther.* 2007; 322:843. [PubMed: 17502430]
5. Gigant B, Wang C, Ravelli RBG, Roussi F, Steinmetz MO, Curmi PA, Sobel A, Knossow M. *Nature.* 2005; 435:519. [PubMed: 15917812]
6. Lachia M, Moody CJ. *Nat Prod Rep.* 2008; 25:227. [PubMed: 18389137]
7. Subramaniam G, Kam TS. *Helv Chim Acta.* 2008; 91:930.
8. Cai XH, Tan QG, Liu YP, Feng T, Du ZZ, Li WQ, Luo XD. *Org Lett.* 2008; 10:577. [PubMed: 18205369]
9. (a) Bertini F, Galli R, Perchinunno M, Minisci F. *Tetrahedron.* 1971; 27:3575. (b) Studer, A., Bossart, M. *Radicals in Organic Synthesis.* Renaud, P., Sibi, MP., editors. Vol. 2. Wiley-VCH; Weinheim: 2001. p. 62-76. (c) Togo, H. *Advanced Free Radical Reactions for Organic Synthesis.* Elsevier; New York: 2004. p. 57-121. (d) Storey JMD, Ladwa MM. *Tetrahedron Lett.* 2006; 47:381.
10. For our previous studies on indoline and isoindolinone synthesis using nucleophilic addition strategies, see: (a) Wipf P, Kim Y. *Tetrahedron Lett.* 1992; 33:5477. (b) Pierce JG, Waller DL, Wipf P. *J Organomet Chem.* 2007; 692:4618. [PubMed: 19684878]
11. (a) Nugent WA, RajanBabu TV. *J Am Chem Soc.* 1988; 110:8561. (b) Rajanbabu TV, Nugent WA. *J Am Chem Soc.* 1994; 116:986.
12. (a) Gansäuer A, Bluhm H, Pierobon M. *J Am Chem Soc.* 1998; 120:12849. (b) Gansäuer A, Pierobon M, Bluhm H. *Angew Chem Int Ed.* 1998; 37:101. (c) Gansäuer A, Barchuk A, Keller F, Schmitt M, Grimme S, Gerenkamp M, Mück-Lichtenfeld C, Daasbjerg K, Svith H. *J Am Chem Soc.* 2007; 129:1359. and references cited therein. [PubMed: 17263420]
13. For reviews, see: (a) Gansäuer A, Pierobon M, Bluhm H. *Synthesis.* 2001; 16:2500. (b) Barrero AF, Quilez del Moral JF, Sanchez EM, Arteaga JF. *Eur J Org Chem.* 2006:1627. For additional applications see: (c) Fernandez-Mateos A, Herrero Teijon P, Mateos Buron L, Rabanedo Clemente R, Rubio Gonzalez R. *J Org Chem.* 2007; 72:9973. [PubMed: 18044915] (d) Monleon LM, Grande M, Anaya J. *Tetrahedron.* 2007; 63:3017.
14. Barvainiene B, Stanisauskaite A, Getautis V. *Chem Het Comp.* 2006; 42:123.
15. Sonication was used to increase reagent solubility.
16. **Typical Procedure:** To a flame-dried 2-neck flask was added 129 mg (0.43 mmol) of **4c**, 3.2 mg (0.01 mmol) of Cp_2TiCl_2 , 102 mg (0.64 mmol) of collidine hydrochloride, and 35.6 mg (0.64 mmol) of Mn^0 . The vessel was fitted with a reflux condenser and purged 3 times with Ar. After addition of 4.3 mL of deoxygenated THF (0.1 M), the reaction mixture was placed in a pre-heated oil bath and heated at reflux for 3 h. The color of the solution gradually changed from light pink to a dark violet. The mixture was cooled to room temperature, quenched with satd. NH_4Cl , extracted with 3×10 mL of Et_2O , washed with brine, dried (MgSO_4), and concentrated *in vacuo*. The residue was redissolved in 5 mL of MeOH and treated with 30 mg (25% w/w, 0.01 mmol) of Pd/C. The mixture was stirred at room temperature under 1 atm of H_2 and the disappearance of starting material was monitored by TLC (hexanes/EtOAc, 1:1). The solution was then quenched with Celite[®], filtered, and purified by chromatography on SiO_2 (hexanes/EtOAc, 1:2) to afford 44 mg (0.26 mmol, 63%, 2 steps) of **8c** as a yellow oil.
17. Popowycz F, Merour JY, Joseph B. *Tetrahedron.* 2007; 63:8689.
18. Scott ME, Schwarz CA, Lautens M. *Org Lett.* 2006; 8:5521. [PubMed: 17107062]
19. Bacque E, Qacemi ME, Zard SZ. *Org Lett.* 2004; 6:3671. [PubMed: 15469320]
20. Mello JV, Finney NS. *Org Lett.* 2001; 3:4263. [PubMed: 11784193]
21. (a) Curran DP, Liu H, Josien H, Ko SB. *Tetrahedron.* 1996; 52:11385. (b) Clyne MA, Aldabbagh F. *Org Biomol Chem.* 2006; 4:268. [PubMed: 16391769] (c) Clive DLJ, Peng J, Fletcher SP, Ziffle VE, Wingert D. *J Org Chem.* 2008; 73:2330. [PubMed: 18278944]
22. (a) Bowman WR, Heaney H, Jordan BM. *Tetrahedron.* 1991; 47:10119. (b) Beckwith ALJ, Bowry VW, Bowman WR, Mann E, Parr J, Storey JMD. *Angew Chem Int Ed.* 2004; 43:95. (c) Curran DP, Keller AI. *J Am Chem Soc.* 2006; 128:13706. [PubMed: 17044696]
23. Performing the reaction with epoxide **4c** using 3 mol % of Cp_2TiCl_2 and 15 mol % of Mn^0 , the desired Cbz-protected indoline was isolated in only 35% yield.

24. For Fridel-Crafts cycloalkylation with arlyalkyl oxiranes, see:(a) Taylor SK, Hockerman GH, Karrick GL, Lyle SB, Schramm SB. J Org Chem. 1983; 48:2449.(b) Taylor SK, May SA, Stansby ES. J Org Chem. 1996; 61:2075.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 1.
Representative natural products with 3,3-disubstituted indoline scaffolds (outlined in bold).

Figure 2.
Proposed catalytic cycle for the titanocene(III) chloride catalyzed epoxide-opening annulation.

Scheme 1.
Preparation of 5-azaindoline **28**

Table 1

Optimization of the titanocene(III) chloride catalyzed reductive cyclization of anilines.^{a,b}

entry	epoxide	Cp ₂ TiCl ₂ (mol %)	Mn (equiv)	concentration [M]	R ¹	R ²	product(s), yield
1	4a	10	0.80	0.03	Ph	H	5a:6a (3:1) ^{c,d}
2	4b	10	0.65	0.03	Ph	CH ₃	5b , 82% ^d
3	4b	10	0.65	0.03	Ph	CH ₃	5b , 84% ^e
4	4c	10	1.5	0.1	Cbz	CH ₃	5c:7c (2:1) ^f
5	4c	3	1.5	0.1	Cbz	CH ₃	8c , 63% ^g
6	4b	3	1.5	0.1	Ph	CH ₃	5b , 89%
7	4a	3	1.5	0.1	Ph	H	5a:6a (3.8:1), 87%
8	4c	3	1.5	0.1	Cbz	CH ₃	5c , 14% ^h
9	4c	3	0	0.1	Cbz	CH ₃	5c , 0%
10	4c	0	1.5	0.1	Cbz	CH ₃	5c , 0%

^aFor epoxide preparation, see Supporting Information.^bAll reactions were performed in degassed THF heated at reflux unless otherwise noted.^cYield not determined; product ratio was determined by GC analysis of crude reaction mixtures.^dReaction was performed in degassed THF at room temperature using sonication.^eReaction was performed in degassed THF at room temperature using magnetic stirring.^fYield not determined; product ratios were determined by ¹H NMR analysis of crude reaction mixtures.^gYield was determined over 2 steps.^hStarting material **4c** was recovered in 43% yield.

Table 2

Indolines prepared using titanocene(III) catalysis

entry	epoxide	indoline	yield
1			65%
2			62% ^a
3			35% ^a
4			69% ^a
5			21% ^a
6			56% ^a
7			41%
8			—

^aYields determined over 2 steps.