Published in final edited form as:

Soc Sci Med. 2017 June ; 182: 136–141. doi:10.1016/j.socscimed.2017.03.052.

Social identity and support for counteracting tobacco company marketing that targets vulnerable populations

Sabeeh A. Baig^a, Jessica K. Pepper^{a,c}, Jennifer C. Morgan^a, and Noel T. Brewer^{a,b}

^aDepartment of Health Behavior, Gillings School of Global Public Health, University of North Carolina, 325B Rosenau Hall, Campus Box 7440, Chapel Hill, NC, 27599 USA

^bLineberger Comprehensive Cancer Center, University of North Carolina, 101 Manning Dr, Chapel Hill, NC, 27514 USA

^cRTI International, 3040 E. Cornwallis Rd., Research Triangle Park, NC 27709 USA

Abstract

Rationale—Tobacco companies use advertising to target vulnerable populations, including youth, racial/ethnic minorities, and sexual minorities.

Objective—We sought to examine how personal identity affects support for population-specific anti-smoking advertisements that could serve as countermeasures to industry practices.

Methods—In 2014–2015, we surveyed probability phone samples of adults and adolescents ($n = 6,139$) and an online convenience sample of adults ($n = 4,137$) in the United States. We experimentally varied the description of tobacco industry marketing practices (no description, general, or specific to a target group). The four prevention target groups were teens; African Americans; Latinos; and gays, lesbians, and bisexuals (GLBs). Participants were either members or non-members of their prevention target group.

Results—Support was highest for anti-smoking advertisements targeting teens, moderate for Latinos and African Americans, and lowest for GLBs. In-group members expressed higher support than out-group members when anti-smoking advertisements targeted African Americans, Latinos, and GLBs (all $p < .05$). However, when teens were the target prevention group, in-group members expressed lower support than out-group members ($p < .05$). The description of industry marketing practices did not have an effect. Results were similar across the phone and online studies.

Conclusions—Our findings suggest that the public strongly supports advertisements to prevent smoking among teens, but support for similar efforts among other vulnerable populations is comparatively low. Anti-smoking campaigns for vulnerable populations may benefit from a greater understanding of the role of social identity in shaping public support for such campaigns.

Corresponding author: Noel T. Brewer, PhD, Department of Health Behavior, Gillings School of Global Public Health, University of North Carolina, 325 Rosenau Hall CB 7440, Chapel Hill, NC 27516, ntb@unc.edu +1 919-966-3282.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Keywords

tobacco control; cigarette smoking; health disparities; vulnerable populations; social identity

Introduction

The leading preventable cause of death in the United States is tobacco use (CDC, 2014). However, smoking rates and consequences are not distributed evenly across the population. Youth are by far the most likely to initiate smoking (Chen & Jacques-Tiura, 2014). Among adults, the highest rates of smoking are among low-income individuals, people with mental health problems, and gays, lesbians, and bisexuals (GLBs; Annamalai, Singh, & O'Malley, 2015; Jamal et al., 2015; Lee, Griffin, & Melvin, 2009). The incidence of lung cancer is substantially higher among African Americans than whites, with the majority of cases being attributable to cigarette smoking (Haiman et al., 2006). Among males over age 50, African Americans have a higher rate of smoking-attributable mortality than whites (Ho & Elo, 2013). Given that Hispanics are a fast-growing segment of the U.S. population, smoking and its related health consequences in this group are of particular interest (Johnson & Lichter, 2008).

The tobacco industry has historically targeted vulnerable populations with marketing that may have contributed to disparities in smoking and subsequent health problems. For example, R.J. Reynolds Tobacco Company developed Project SCUM (Sub-Culture Urban Marketing) in 1995 as a strategy for marketing Camel and Red Kamel cigarettes to gay residents and homeless individuals in the San Francisco area (Stevens, Carlson, & Hinman, 2004; Washington, 2002). Philip Morris has targeted the Hispanic community in the U.S. since 1988 by sponsoring and distributing cigarettes at cultural and sporting events (Washington, 2002). Tobacco companies have also targeted African Americans by advertising in major African American publications such as *Ebony* and making large donations to advocacy organizations such as the National Association for the Advancement of Colored People (Anderson, 2011; Washington, 2002).

The Internet has provided tobacco companies with new opportunities to reach vulnerable populations beyond traditional marketing strategies and media. Online e-cigarette marketing to youth exemplifies the utility of this platform. As of 2014, online vendors were selling 7,764 flavors of e-liquid, including candy and other sweet flavors that may be particularly appealing to children and adolescents (Zhu et al., 2014). The unregulated nature of the Internet could enable further targeted marketing of tobacco products to minority populations (Liang et al., 2015; Ribisl, 2003).

Anti-smoking campaigns that target vulnerable groups can be effective countermeasures to industry marketing practices. Public support allows agencies to implement programs that target diverse groups, and the lack of that support can imperil new programs (Sun, 2014). Furthermore, public support for anti-smoking advertising campaigns may increase their impact (Samu & Bhatnagar, 2008). Targeted anti-smoking campaigns that have public support may also encourage interpersonal communication among community members about smoking behaviors, potentially increasing campaign effectiveness (Hall et al., 2015;

Sun, 2014). Finally, within targeted communities, support by key opinion leaders may be especially important to the viability and longevity of anti-smoking campaigns (Howard et al., 2000; Valente & Pumpuang, 2006).

Support for anti-smoking advertisements may vary by social identity. According to social identity theory, social interactions make people aware of similarities and differences with others when social categories, such as race and sexual orientation, become salient (Hornsey, 2008). When social categories are salient, people enhance similarities within their own group and differences between their group and other groups (Hogg & Reid, 2006). They also try to protect their own group and improve its status in order to maintain a positive social identity (Major, Mendes, & Dovidio, 2013). Social identity theory can provide insights about health and well-being (Haslam, Jetten, Postmes, & Haslam, 2009). For example, if group identity is made salient in health-related campaign materials, in-group members' desire to improve the health of their peers may enhance support for that health campaign.

Drawing on some aspects of social identity theory, we hypothesized that support for targeted anti-smoking campaigns is higher among in-group members of a target population compared to out-group members. We also hypothesized that anti-smoking advertisements targeting teens would elicit the greatest support because people view youth as a vulnerable group that society should protect, and many are already aware that the tobacco industry targets youth (Henriksen, Feighery, Wang, & Fortmann, 2004; Landman, Ling, & Glantz, 2002; Slater, Chaloupka, Wakefield, Johnston, & O'Malley, 2007). We examined these hypotheses in two national experiments.

Experiment 1

Methods

Sample—From September 2014 to June 2015, the Carolina Survey Research Laboratory (CSRL) conducted phone surveys with a probability sample of 5,014 U.S. adults ages 18 or older identified using random digit dial landline and cell phone frames. Additional information on sampling and methodology is available elsewhere (Boynton et al., 2016). Interviewers obtained verbal consent from adult participants. CSRL also conducted a phone survey with a probability sample of 1,125 adolescents aged 13 to 17 from November 2014 to June 2015. Interviewers obtained verbal parental consent and adolescent assent. The survey included smokers and non-smokers in order to capture support for targeted anti-smoking advertisements among a variety of stakeholders. The Institutional Review Board at the University of North Carolina approved both two studies.

Procedures—We randomly assigned participants to one of 24 conditions in a $3 \times 4 \times 2$ between-subjects factorial experiment. The first factor was *industry marketing practices*: no statement about these practices, a general statement (e.g. "Cigarette companies target certain groups with advertising"), or a population-specific statement (e.g. "Cigarette companies specifically target [teens; gays, lesbians, and bisexuals; African Americans; or Latinos] with advertising"). The second factor was the *prevention target group* that anti-smoking advertisements would focus on: teens, GLBs, African Americans, or Latinos. The third factor was *in-group membership* (i.e., whether the respondents themselves were teens,

GLBs, African Americans, or Latinos). We assigned participants, in the order above, to respond about their own group. Participants who were not members of any of the four prevention target groups responded about a randomly-assigned group. Assignment to the industry marketing practice condition was also random.

Measures—The survey assessed support for anti-smoking advertisements with the question, “How much would you support organizations like the CDC running anti-smoking advertisements for [prevention target group]?” The response options were “not at all”, “a little”, “somewhat”, or “a lot” (coded as 1–4). The survey also measured sex, age, participant education (adults), maternal education (adolescents), income (adults), race, ethnicity, sexual orientation (adults), numeracy, and smoking status. We defined adult current smokers as those who had smoked at least 100 cigarettes in their lifetime and currently smoked every day or some days and adolescent current smokers as those who had smoked at least one cigarette in the past 30 days (Arrozola et al., 2015; Davis et al., 2009).

Data Analysis—We analyzed data from the phone and online surveys separately. We excluded 202 participants who did not respond to the primary outcome measure, resulting in an analytic sample of 5,937 participants. Missing data were more common for people in the GLB than teen prevention target group conditions and for in-group than out-group members (both $p < .001$). To check whether random assignment created demographically equivalent groups by industry marketing practices (no statement, general, or specific), we used chi-square tests and t -tests. Among “out-group” members (i.e., adults who were not GLBs, African-Americans, or Latinos), we used the same procedures for checking successful randomization into prevention target group. Five demographic characteristics of adult participants (sex, race, ethnicity, education, and sexual orientation) and four demographic characteristics of adolescent participants (sex, race, ethnicity, and maternal education) were equivalent across experimental conditions (11 of 11 tests, all $p > .05$).

We used a $3 \times 4 \times 2$ between-subjects ANOVA to examine the effects of industry marketing practices (no statement, general, or specific), prevention target group (teens, GLBs, African Americans, or Latinos), and in-group membership on support for anti-smoking advertisements. Statistical analyses used SAS (v 9.4) and a critical alpha of .05, except for post-hoc pairwise t -tests that used Holm-Bonferroni adjustments to critical alpha.

Results

About half of the adult participants were male (47%), had no college degree (56%), and had household incomes less than \$50,000 per year (54%) (Table 1). A minority of adult participants were current smokers (23%). Among adolescents, about half were male (50%), a quarter had low numeracy (27%), and few were current smokers (4%).

Prevention Target Group—Prevention target group affected support for anti-smoking advertisements ($p < .001$) (Table 2). Participants expressed higher support for anti-smoking advertisements aimed at teens ($M = 3.48$, $SD = 0.85$) than Latinos ($M = 3.06$, $SD = 1.14$; $p < .001$), similar support for Latinos and African Americans ($p = .31$), and higher support for African Americans ($M = 3.01$, $SD = 1.14$) than GLBs ($M = 2.50$, $SD = 1.27$; $p < .001$).

In-group Membership—Support varied by group membership so that in-group members ($M = 3.27$, $SD = 1.01$) expressed more support for anti-smoking advertisements than out-group members ($M = 2.97$, $SD = 1.19$; $p < .001$) (Table 2). The effect of group membership was moderated by prevention target group ($p < .001$). Post-hoc analyses showed that support for anti-smoking advertisements was higher among in-group members than out-group members for African Americans ($p < .001$), Latinos ($p < .001$), and GLBs ($p < .01$) (Figure 1). In contrast, support was lower among in-group than out-group members when teens were the prevention target group ($p = .04$).

Industry Marketing Practices—Support did not differ by the description of industry marketing practices ($p = .78$) (Table 2). None of the two-way or three-way interactions with industry marketing practices were statistically significant.

Experiment 2

We sought to replicate the findings of Experiment 1 in another national study conducted online among adults.

Methods

We conducted an online survey in December 2014 with a convenience sample of 4,137 adults ages 18 or older, recruited using Amazon Mechanical Turk. The online survey announcement encouraged smokers to participate but did not prevent non-smokers from participating. Because Experiment 2 did not include adolescents, it did not have an in-group condition for teens. Experiment 2 used identical measures as those in the adult phone survey in Experiment 1. Eight participants were missing on the primary outcome, leaving an analytic sample of 4,129 participants.

Five demographic characteristics of participants (sex, race, ethnicity, education, and sexual orientation) were equivalent across experimental conditions (6 of 7 tests, all $p > .05$); the exception was that Latinos were more commonly assigned to the “no statement” condition than the other two industry marketing practice conditions in the online survey ($p < .05$). Analyses for Experiment 2 were similar to those for Experiment 1 except for one difference. Because one cell of the factorial design (teen in-group) was missing for the online survey of adults, we dropped responses about teens in the ANOVA. Thus, we used a $3 \times 3 \times 2$ between-subjects ANOVA. We used an additional one-way ANOVA among out-group participants to examine differences in support for anti-smoking advertisements for all four prevention target groups including teens.

Results

Sample characteristics were similar to those of adult participants in Experiment 1 (Table 1).

Prevention Target Group—In the online survey, prevention target again affected support for anti-smoking advertisements ($p < .001$) (Table 3). Support was similar for anti-smoking advertisements targeting Latinos ($M = 2.90$, $SD = 1.04$) and African Americans ($p = .24$) and higher for African Americans ($M = 2.85$, $SD = 1.07$) than GLBs ($M = 2.50$, $SD = 1.16$; $p < .001$). A separate analysis of out-group participants only found that they expressed

higher support for teens ($M = 3.38$, $SD = 0.88$) than Latinos ($M = 2.81$, $SD = 1.04$; $p < .001$).

In-group Membership—Support varied by group membership so that in-group members ($M = 2.93$, $SD = 1.09$) expressed more support than out-group members ($M = 2.65$, $SD = 1.10$; $p < .001$) (Table 3). The effect of group membership was moderated by prevention target group ($p < .001$). Support for anti-smoking advertisements was again higher among in-group members than out-group members for African Americans ($p < .001$), Latinos ($p < .001$), and GLBs ($p = .008$) (Figure 2). However, the in-group/out-group difference was larger for African Americans ($M = .58$) than for Latinos or GLBs (M s = .33 and .18, both $p < .05$). Because no teens participated in the online survey, it was not possible to examine the effect of membership on support for this target prevention group.

Industry Marketing Practices—Support did not differ by the description of industry marketing practices ($p = .13$) (Table 3). None of the two-way or three-way interactions with this variable were statistically significant.

Discussion

The findings of Experiment 2 were very similar to Experiment 1. Targeted anti-tobacco marketing is an important strategy for reducing tobacco use among vulnerable groups (e.g., the teen-targeted Truth campaign successfully reduced smoking among adolescents) (Hershey et al., 2005; Thrasher, 2004). Our findings indicated strong public support for anti-smoking campaigns for teens. However, public support for similar efforts for GLBs, African Americans, and Latinos was lower. Tobacco countermeasures for a vulnerable population may wish to actively and visibly involve members of that population in order to activate in-group identity and consequent support.

As hypothesized, in-group members were generally more supportive of anti-smoking advertisements than out-group members. As described by social identity theory, group membership allows individuals to achieve a positive social identity (Haslam et al., 2009). As such, social identity may affect how individuals react to health-related norms and engage in health-related behaviors. For three prevention target groups (African-Americans, Latinos, and GLBs) in Experiment 1, in-group members expressed higher support for anti-smoking campaigns than out-group members, but this finding was reversed for teens. This reversal may be because adults view teens as a group that needs protection, but teens do not view themselves that way. Adolescents minimize or do not fully understand the negative consequences of smoking (Mantler, 2013; Slovic, 1998, 2000). If teens view themselves and their peers as unaffected by the harms of smoking, they may not see a need to support anti-smoking campaigns for teens. Alternatively, teens may be conceptualized as an inherently heterogeneous group because it includes GLBs, African-Americans, and Latinos. If being a teen is not a primary social identity for adolescents, then making this identity salient through targeting could have a smaller effect on support for targeted anti-smoking campaigns among teens than adults.

Our hypothesis that support would be higher for teens than for other groups was confirmed. In both experiments, support for anti-smoking campaigns for teens was higher than for all other groups, possibly reflecting broad agreement among the public that teens should be prevented from engaging in risky health behaviors. The finding may also reflect widespread public knowledge that special efforts to support teens are necessary because the tobacco industry specifically targets youth, a message widely delivered by the Truth campaign. The emphasis on teens by current anti-smoking efforts is both practical in terms of targeting the group at highest risk for initiating (youth) and building on broad public support.

Our descriptions of industry marketing practices had no effect on support, perhaps because it was a low-dose manipulation (only briefly mentioned or not) in comparison to the other two manipulations. Any effect of the industry marketing practices manipulation may have been hampered by poor knowledge that certain groups are a priority for tobacco prevention and control efforts. For example, if participants did not know that GLBs have some of the highest rates of smoking in the U.S., urgency to act against tobacco industry marketing specifically to GLBs may have been low (Lee et al., 2009). Alternatively, because people understand that many industries use targeted marketing, participants may not have been surprised to learn that tobacco companies do the same. Thus, the presence or absence of a brief statement about targeted tobacco marketing may have had limited importance to participants.

Strengths and Limitations

Experiment 1 relied on large probability samples with strong representation of minority populations relevant to tobacco use prevention and control. Although Experiment 2 used a convenience sample, its results were similar to those from Experiment 1. A limitation is that individuals who were members of prevention target groups were assigned in a specific order (teens, GLBs, African Americans, and Latinos) to an experimental condition, but they may be members of more than one of these groups. Thus, they may have responded about a group that was not their primary identity. We also did not examine psychosocial factors that explain the rank order of prevention target groups with respect to support for targeted advertisements. We speculate that respondents may have felt that youth are especially vulnerable to manipulation by tobacco companies and were largely unaware of the high rates of smoking among GLBs. Another limitation is that we could not examine the extent to which in-group members supported anti-smoking campaigns for out-group members. These are important areas for future research.

Other limitations concern the primary outcome measure. Because the primary outcome measure referenced the CDC, differential knowledge of what the CDC is or the extent to which they view the CDC as a credible source may have influenced participants' responses (Schmidt, Ranney, Pepper, & Goldstein, 2016). The primary outcome measure also used a four-point response scale, which could have limited variability in the participants' responses. However, research indicates that four-point scales still have good psychometric properties, and the large sample sizes afforded substantial power (Lozano, García-Cueto, & Muñiz, 2008). Finally, the experiment presented hypothetical scenarios, a useful strategy for early exploration of topics like acceptability of new products, medical services, and advertising

approaches (Brewer & Fazekas, 2007; Brewer, Richman, DeFrank, Reyna, & Carey, 2012; Pepper, Reiter, McRee, & Brewer, 2012).

Research and Practice Implications

People expressed higher support for tobacco countermeasures that were tied to the social identities specified in our experiment than countermeasures that were not. By extension, incorporating identity processes in the study of health and well-being might be helpful in the development of novel, sustainable social marketing and health communication campaigns. In order to do so, future studies should examine whether our findings hold for other health issues such as infectious diseases and obesity and other social identities. Replication of our findings about group membership across multiple health issues may support the claim that members of vulnerable groups are more motivated to improve the health of their own vulnerable groups than individuals who are not members. Future studies should further examine the impact of specifying both the in-group and out-group on support for targeted anti-smoking campaigns. It is possible that doing so will activate positive and negative stereotypes about the out-groups, which may have unexpected effects on campaign support.

Our findings also have implications for tobacco control funding. Future studies should examine what “ingredients” of messages work the best to bolster support for targeted anti-smoking campaigns for vulnerable groups. For example, providing in-depth descriptions of specific tobacco industry marketing campaigns that target vulnerable populations, like Project SCUM, might enhance support. Including information about smoking disparities or other population-specific threats could do the same by providing meaningful context (Niederdeppe, Bu, Borah, Kindig, & Robert, 2008). Future studies should also explore the psychological processes that salient group membership activates and how anti-smoking campaigns can take advantage of social identity processes to increase impact and reach. Finally, resources for tobacco control should target GLBs in particular given that this group has some of the highest smoking rates in the US and the lowest public support for targeted campaigns. This research will hopefully result in theory-driven, effective campaigns to prevent and reduce smoking among vulnerable groups.

References

- Anderson SJ. Marketing of menthol cigarettes and consumer perceptions: a review of tobacco industry documents. *Tobacco Control*. 2011; 20(Supplement 2):ii20–ii28. <https://doi.org/10.1136/tc.2010.041939>. [PubMed: 21504928]
- Annamalai A, Singh N, O'Malley SS. Smoking Use and Cessation Among People with Serious Mental Illness. *The Yale Journal of Biology and Medicine*. 2015; 88(3):271–7. [PubMed: 26339210]
- Boynton MH, Agans RP, Bowling JM, Brewer NT, Sutfin EL, Goldstein AO, ... Ribisl KM. Understanding how perceptions of tobacco constituents and the FDA relate to effective and credible tobacco risk messaging: A national phone survey of U.S. adults, 2014–2015. *BMC Public Health*. 2016; 16(1) <https://doi.org/10.1186/s12889-016-3151-5>.
- Brewer NT, Fazekas KI. Predictors of HPV vaccine acceptability: A theory-informed, systematic review. *Preventive Medicine*. 2007; 45(2–3):107–114. <https://doi.org/10.1016/j.ypmed.2007.05.013>. [PubMed: 17628649]
- Brewer NT, Richman AR, DeFrank JT, Reyna VF, Carey LA. Improving communication of breast cancer recurrence risk. *Breast Cancer Research and Treatment*. 2012; 133(2):553–561. <https://doi.org/10.1007/s10549-011-1791-9>. [PubMed: 21964579]

- Chen X, Jacques-Tiura AJ. Smoking initiation associated with specific periods in the life course from birth to young adulthood: Data from the National Longitudinal Survey of youth 1997. *American Journal of Public Health*. 2014; 104(2):e119–e126. <https://doi.org/10.2105/AJPH.2013.301530>.
- Haiman CA, Stram DO, Wilkens LR, Pike MC, Kolonel LN, Henderson BE, Le Marchand L. Ethnic and racial differences in the smoking-related risk of lung cancer. *New England Journal of Medicine*. 2006; 354(4):333–342. [PubMed: 16436765]
- Hall M, Peebles K, Bach L, Noar S, Ribisl K, Brewer N. Social Interactions Sparked by Pictorial Warnings on Cigarette Packs. *International Journal of Environmental Research and Public Health*. 2015; 12(10):13195–13208. <https://doi.org/10.3390/ijerph121013195>. [PubMed: 26506363]
- Haslam SA, Jetten J, Postmes T, Haslam C. Social identity, health and well-being: An emerging agenda for applied psychology. *Applied Psychology*. 2009; 58(1):1–23. <https://doi.org/10.1111/j.1464-0597.2008.00379.x>.
- Henriksen L, Feighery EC, Wang Y, Fortmann SP. Association of retail tobacco marketing with adolescent smoking. *American Journal of Public Health*. 2004; 94(12):2081–2083. <https://doi.org/10.2105/AJPH.94.12.2081>. [PubMed: 15569957]
- Hershey JC, Niederdeppe J, Evans WD, Nonnemaker J, Blahut S, Holden D, ... Haviland ML. The Theory of “truth”: How Counterindustry Campaigns Affect Smoking Behavior Among Teens. *Health Psychology*. 2005; 24(1):22–31. <https://doi.org/10.1037/0278-6133.24.1.22>. [PubMed: 15631559]
- Ho JY, Elo IT. The Contribution of Smoking to Black-White Differences in U.S. Mortality. *Demography*. 2013; 50(2):545–568. <https://doi.org/10.1007/s13524-012-0159-z>. [PubMed: 23086667]
- Hogg MA, Reid SA. Social Identity, Self-Categorization, and the Communication of Group Norms. *Communication Theory*. 2006; 16(1):7–30. <https://doi.org/10.1111/j.1468-2885.2006.00003.x>.
- Hornsey MJ. Social Identity Theory and Self-categorization Theory: A Historical Review. *Social and Personality Psychology Compass*. 2008; 2(1):204–222. <https://doi.org/10.1111/j.1751-9004.2007.00066.x>.
- Howard KA, Rogers T, Howard-Pitney B, Flora JA, Norman GJ, Ribisl KM. Opinion leaders’ support for tobacco control policies and participation in tobacco control activities. *American Journal of Public Health*. 2000; 90(8):1283. [PubMed: 10937010]
- Jamal A, Homa DM, O’Connor E, Babb SD, Caraballo RS, Singh T, ... King BA. Current Cigarette Smoking Among Adults - United States, 2005–2014. *MMWR. Morbidity and Mortality Weekly Report*. 2015; 64(44):1233–40. <https://doi.org/10.15585/mmwr.mm6444a2>. [PubMed: 26562061]
- Johnson KM, Lichter DT. Natural Increase: A New Source of Population Growth in Emerging Hispanic Destinations in the United States. *Population and Development Review*. 2008; 34(2): 327–346. <https://doi.org/10.1111/j.1728-4457.2008.00222.x>.
- Landman A, Ling PM, Glantz Sa. Tobacco industry youth smoking prevention programs: protecting the industry and hurting tobacco control. *American Journal of Public Health*. 2002; 92(6):917–30. <https://doi.org/10.2105/AJPH.92.6.917>. [PubMed: 12036777]
- Lee JGL, Griffin GK, Melvin CL. Tobacco use among sexual minorities in the USA, 1987 to May 2007: a systematic review. *Tobacco Control*. 2009; 18(4):275–282. <https://doi.org/10.1136/tc.2008.028241>. [PubMed: 19208668]
- Liang Y, Zheng X, Zeng DD, Zhou X, Leischow SJ, Chung W. Exploring How the Tobacco Industry Presents and Promotes Itself in Social Media. *Journal of Medical Internet Research*. 2015; 17(1):e24. <https://doi.org/10.2196/jmir.3665>. [PubMed: 25608524]
- Lozano LM, García-Cueto E, Muñoz J. Effect of the Number of Response Categories on the Reliability and Validity of Rating Scales. *Methodology*. 2008; 4(2):73–79. <https://doi.org/10.1027/1614-2241.4.2.73>.
- Major B, Mendes WB, Dovidio JF. Intergroup relations and health disparities: a social psychological perspective. *Health Psychology*. 2013; 32(5):514–24. <https://doi.org/10.1037/a0030358>. [PubMed: 23646834]
- Mantler T. A systematic review of smoking Youths’ perceptions of addiction and health risks associated with smoking: Utilizing the framework of the health belief model. *Addiction Research & Theory*. 2013; 21(4):306–317. <https://doi.org/10.3109/16066359.2012.727505>.

- Max W, Sung HY, Tucker LY, Stark B. The disproportionate cost of smoking for African Americans in California. *American Journal of Public Health*. 2010; 100(1):152–158. [PubMed: 19965569]
- National Center for Chronic Disease Prevention and Health Promotion (US) Office on Smoking and Health. The Health Consequences of Smoking—50 Years of Progress: A Report of the Surgeon General. Atlanta (GA): Centers for Disease Control and Prevention (US); 2014. Retrieved from <http://www.ncbi.nlm.nih.gov/books/NBK179276/>
- Niederdeppe J, Bu QL, Borah P, Kindig DA, Robert SA. Message design strategies to raise public awareness of social determinants of health and population health disparities. *The Milbank Quarterly*. 2008; 86(3):481–513. <https://doi.org/10.1111/j.1468-0009.2008.00530.x>. [PubMed: 18798887]
- Pepper JK, Reiter PL, McRee AL, Brewer NT. Advertisements promoting human papillomavirus vaccine for adolescent boys: Does source matter? *Sexually Transmitted Infections*. 2012; 88(4): 264–265. <https://doi.org/10.1136/sextrans-2011-050197>. [PubMed: 22223814]
- Ribisl KM. The potential of the internet as a medium to encourage and discourage youth tobacco use. *Tobacco Control*. 2003; 12(Suppl 1):i48–59. [PubMed: 12773785]
- Samu S, Bhatnagar N. The efficacy of anti-smoking advertisements: the role of source, message, and individual characteristics. *International Journal of Nonprofit and Voluntary Sector Marketing*. 2008; 13(3):237–250. <https://doi.org/10.1002/nvsm.326>.
- Schmidt AM, Ranney LM, Pepper JK, Goldstein AO. Source Credibility in Tobacco Control Messaging. *Tobacco Regulatory Science*. 2016; 2(1):31–37. <https://doi.org/10.18001/TRS.2.1.3>. [PubMed: 27525298]
- Slater SJ, Chaloupka FJ, Wakefield M, Johnston LD, O'Malley PM. The impact of retail cigarette marketing practices on youth smoking uptake. *Archives of Pediatrics & Adolescent Medicine*. 2007; 161(5):440–445. [PubMed: 17485618]
- Slovic P. Do Adolescent Smokers Know the Risks? *Duke Law Journal*. 1998; 47(6):1133–1141. [PubMed: 10557547]
- Slovic P. What does it mean to know a cumulative risk? Adolescents' perceptions of short-term and long-term consequences of smoking. *Journal of Behavioral Decision Making*. 2000; 13(2):259.
- Stevens P, Carlson LM, Hinman JM. An Analysis of Tobacco Industry Marketing to Lesbian, Gay, Bisexual, and Transgender (LGBT) Populations: Strategies for Mainstream Tobacco Control and Prevention. *Health Promotion Practice*. 2004; 5(3):129–134. <https://doi.org/10.1177/1524839904264617>.
- Sun Y. Rethinking public health: Promoting public engagement through a new discursive environment. *American Journal of Public Health*. 2014; 104(1):e6–e13.
- Thrasher JF. The impact of anti-tobacco industry prevention messages in tobacco producing regions: evidence from the US truth(R) campaign. *Tobacco Control*. 2004; 13(3):283–288. <https://doi.org/10.1136/tc.2003.006403>. [PubMed: 15333885]
- Valente TW, Pumpuang P. Identifying Opinion Leaders to Promote Behavior Change. *Health Education & Behavior*. 2006; 34(6):881–896. <https://doi.org/10.1177/1090198106297855>.
- Washington HA. Burning Love: big tobacco takes aim at LGBT youths. *American Journal of Public Health*. 2002; 92(7):1086–95. [PubMed: 12084686]
- Zhu S-H, Sun JY, Bonnevie E, Cummins SE, Gamst A, Yin L, Lee M. Four hundred and sixty brands of e-cigarettes and counting: implications for product regulation. *Tobacco Control*. 2014; 23(Suppl 3):iii3–9. <https://doi.org/10.1136/tobaccocontrol-2014-051670>. [PubMed: 24935895]

Highlights

- Support for anti-smoking advertisements for teens is higher than for other vulnerable groups.
- Consistent with theory, in-group members had higher support than non-members.
- Understanding reasons for support may improve future anti-smoking advertising campaigns.

Figure 1.
Support for anti-smoking advertisements, Experiment 1

Figure 2.
Support for anti-smoking advertisements, Experiment 2

Table 1

Participant characteristics.

	Adult Phone (<i>n</i> = 5,014) %	Adolescent Phone (<i>n</i> = 1,125) %	Adult Online (<i>n</i> = 4,137) %
Male	47.3	49.9	49.4
Age, years			
13–17	-	100	-
18–25	16.2	-	24.5
26–34	14.9	-	39.8
35–44	15.4	-	19.2
45–54	19.8	-	9.9
55–64	17.9	-	5.6
65+	15.8	-	1.0
Education			
< High school	10.5	4.3	1.0
High school	24.7	16.3	12.5
Some college	20.7	16.9	30.9
Associate's degree	9.9	10.2	12.4
College degree	21.2	31.9	33.7
Master's degree	10.1	16.8	7.5
Professional or doctoral degree	2.9	3.6	2.0
Household income, per year			
\$0 – \$24,999	28.2	-	22.7
\$25,000 – \$49,999	26.2	-	34.5
\$50,000 – \$74,999	19.0	-	23.0
\$75,000 – \$100,000	10.8	-	12.0
> \$100,000 per year	15.8	-	7.8
Race			
White	69.6	80.2	83.0
Black or African American	19.6	10.6	7.6
Native American	2.7	1.6	0.9
Asian	2.1	2.0	5.0
Other	6.0	5.7	3.5
Hispanic	8.6	7.6	8.2
Sexual orientation			
Straight	94.4	-	88.4
Gay, lesbian, or bisexual	3.8	-	11.5
Other	1.8	-	0.1
Smoker	23.0	3.7	37.8

Note. Maternal education is reported for adolescents. High school includes equivalency certificate (general educational development). Current smoking is defined for adults as having ever smoked at least 100 cigarettes and currently smoking every day or some days and for adolescents as smoking at least once during the last 30 days.

Table 2

Effects of manipulations on support for anti-smoking ads, Experiment 1.

	<i>df</i>	<i>F</i>
Prevention target group	3	105.36**
In-group membership	1	30.25**
Prevention target group × In-group membership	3	11.17**
Industry marketing practices	2	0.25
Industry marketing practices × Target group	6	1.98
Industry marketing practices × In-group membership	2	2.06
Industry marketing practices × Target group × In-group membership	6	0.61

Note. $n = 5,937$. Prevention target groups were teens; African Americans; Latinos; or gays, lesbians, and bisexuals. In-group membership was whether the study participant was a member of the target group. Industry marketing practices was no description of the industry practice; general description of it; or specific to a target group.

*
 $p < .05$;

**
 $p < .001$.

Table 3

Effects of manipulations on support for anti-smoking ads, Experiment 2.

	<i>df</i>	<i>F</i>
Prevention target group ^a	2	61.62 **
In-group membership ^b	1	76.72 **
Prevention target group × In-group membership	2	8.25 **
Industry marketing practices ^c	2	2.02
Industry marketing practices × Target group	4	1.73
Industry marketing practices × In-group membership	2	2.91
Industry marketing practices × Target group × In-group membership	4	0.25

Note. $n = 4,129$. Prevention target groups were African Americans; Latinos; or gays, lesbians, and bisexuals. In-group membership was whether the study participant was a member of the target group. Industry marketing practices was no description of the industry practice; general description of it; or specific to a target group.

*
 $p < .05$;

**
 $p < .001$.