Published in final edited form as:

Res Sports Med. 2017 ; 25(2): 219–230. doi:10.1080/15438627.2017.1282361.

Prevalence and proposed mechanisms of chronic low back pain in baseball: part i

Joseph G. Wasser, Jason L. Zaremski, Daniel C. Herman, and Heather K. Vincent

Department of Orthopaedics and Rehabilitation, Division of Research; Interdisciplinary Center for Musculoskeletal Training and Research, University of Florida, Gainesville, FL, United States of America

Abstract

The prevalence of low back pain (LBP) among active baseball players ranges between 3 and 15%. The execution of baseball-specific manoeuvres, such as pitching or batting, may be related to the onset of LBP. These baseball motions are complex and require appropriate activation of the core musculature to produce a well-timed motion with forces minimized at the extremities. The spine, core and back musculature are involved with acceleration and deceleration of rotational motions. This narrative review synthesizes the available evidence of the prevalence of and mechanical factors underlying LBP in the baseball population. Possible mechanical mechanisms linking baseball play to LBP include aberrant motion, improper timing, high lumbar stress due to mechanical loading and lumbopelvic strength deficits. Potential clinical implications relating to these possible mechanical mechanisms will also be highlighted. The state of the evidence suggests that there are deficits in understanding the role of baseball motion and playing history in the development of spine conditions.

Keywords

Baseball; low back pain; kinematics; stress; injury

Introduction

The preponderance of research relating to baseball injuries has focused on the incidence and prevention of throwing arm injuries, as these are the most prevalent injury type in baseball players of all ages and skill levels (Dick et al., 2007; Hangai et al., 2009, 2010; Li et al., 2013; Posner, Cameron, Wolf, Belmont, & Owens, 2011; Schmidt et al., 2014). However, the lumbar spine is an understudied but pivotal link to the kinematic chain where the energy transfer occurs from the lower body to upper body during throwing and hitting (Young, Herring, Press, & Casazza, 1996). The execution of baseball-specific manoeuvres, such as

CONTACT: Joseph G. Wasser, wassejg@ortho.ufl.edu, Department of Orthopaedics and Rehabilitation, Division of Research, UF Orthopaedics and Sports Medicine Institute (OSMI), PO Box 112727, Gainesville, FL, 32611, USA.

ORCID

Joseph G. Wasser <http://orcid.org/0000-0002-1634-5658>

Disclosure statement

No potential conflict of interest was reported by the authors.

pitching or batting, is complex and requires appropriate activation of the core musculature to produce a well-timed motion with forces minimized at the extremities. The spine, core and back musculature are involved with acceleration and deceleration of rotational motions. The movement of the pelvis and torso about the spine contribute 50% of the energy into throwing (Oliver & Keeley, 2010). In healthy players with adequate dynamic strength and appropriate execution of movement, the risk for low back pain (LBP) and non-contact injury is relatively low compared with reported non-contact upper and lower extremity injuries (Dick et al., 2007; Posner et al., 2011). LBP and/or injury may develop as a consequence of deficiencies in core muscle strength, uncoupling of the movement of the kinematic chain and biomechanical loading (Seroyer et al., 2010).

Currently there are approximately 19 million student-athletes and adults participating in organized baseball play (Dick et al. 2007b). Among collegiate and professional baseball players alone, low back injury rates range from 8.3% to 15%.; Dick et al., 2007 (Posner et al., 2011, 2011) Chronic LBP prevalence ranges from 1% to 40% of baseball players of all experience levels (Bono, 2004; P. A.; d'Hemecourt, Gerbino, & Micheli, 2000). LBP is a significant issue for the baseball player, as it contributes to missed participation time, disability and early career termination (Petering & Webb, 2011). Moreover, because baseball motions of throwing and batting involve whole body neuromuscular coordination, pain or injury to the low back can alter neuromuscular patterns of the upper body and subsequently contribute to secondary injuries elsewhere in the kinematic chain (Chaudhari, McKenzie, Pan, & Oñate, 2014). At present, there is not yet a synopsis of the potential mechanical factors that contribute to LBP and methods to address these factors. This focused, narrative review integrates the available evidence of the potential contributors to and prevalence of chronic LBP and acute on chronic low back injuries in baseball. Specific history questions during the clinical assessment are provided to guide the clinician on identification of mechanical issues underlying LBP.

Literature search

Relevant studies in English were identified through a literature search of MEDLINE and the Cochrane Database from 1990 to 2015. Reference lists of identified sources were also manually reviewed to obtain additional relevant sources. A literature search was conducted with the keywords including “baseball”, “low back”, “lumbar”, “injury”, “back injury”, “epidemiology”, “surveillance”, “collegiate”, “youth”, “elite”, “batting” and “pitcher”. To supplement the information necessary to capture factors that may contribute to injury, the terms “biomechanics”, “motion”, “swing”, “kinematics”, “muscle strength”, “kinematic chain”, “rotation” and “core musculature” were used. A total of **1,270** articles were initially identified. These references were screened to determine whether or not evidence of LBP risk related to baseball play, potential injury mechanisms and methods to reduce LBP in this population was included. Papers were excluded if the content was not related to American-style baseball, did not include low back injuries, were in females, or described acute injuries. Using the levels of evidence from the Agency for Healthcare Research and Quality, U.S. Department of Health and Human Services, the papers presented in the epidemiology section next were categorized into Level 1 (randomized controlled trials), Level 2 (non-randomized controlled trials including prospective cohorts), Level 3 (observational studies

with controls, case control studies), Level 4 (observational studies without controls) and opinion papers. Studies of cervical and thoracic spine and acute injuries without chronic onset were not included in this review.

Association of LBP in individuals with baseball playing history

A total of seven studies included specific LBP data in populations with baseball playing history (Dick et al., 2007; Earhart, Roberts, Roc, Gryzlo, & Hsu, 2012; Hangai et al., 2009, 2010; Ladenhauf, Fabricant, Grossman, Widmann, & Green, 2013; Li et al., 2013; Posner et al., 2011). Among these, three were Level III (Dick et al., 2007; Hangai et al., 2009; Posner et al., 2011) and three were Level IV (Earhart et al., 2012; Hangai et al., 2010; Li et al., 2013) evidence.

Adolescent players

Patterns of back pain are different in adolescent and high school athletes compared with adults (Purcell & Micheli, 2009). Skeletal immaturity coupled with exercise stressors can contribute to LBP and back injuries. High quality evidence in youth baseball players is limited. However, a retrospective study of a cohort of adolescents (mean age 13.9 ± 2.2 years) with symptomatic baseball participation was identified as the fourth most common sport identified, 8.9% of the patients being identified as baseball players, within the cohort of patients with spondylolysis (Ladenhauf et al., 2013). In other studies referenced by Ladenhauf et al., LBP frequency occurred in between 8.5% and 14.6% of youth baseball players (ages 12–15.5; Anderson, Sarwark, Conway, Logue, & Schafer, 2000; Blanda, Bethem, Moats, & Lew, 1993; Miller, Congeni, & Swanson, 2004).

Collegiate student-athletes

Hangai et al. (2009) conducted a cross-sectional study involving trained collegiate athletes ($n = 308$) and focused on the incidence of disc degeneration among these athletes. Baseball players have up to a 3.23 greater likelihood of being diagnosed with disc degeneration over non-athletes. This odds ratio ranks higher than athletes from other sports (not including football), such as volleyball, basketball and soccer (Hangai et al., 2009). In another cross-sectional study by Hangai et al. (2010), 4,667 new, non-varsity athlete university students were surveyed for their experience with LBP and the duration of which they may have played competitive sports when they were younger. The findings of the self-report survey revealed that those students who participated in competitive sports had higher prevalence of LBP than non-active students. Moreover, those students who had a long-duration sports experience (participated in sport from elementary school through high school) had the highest rates of LBP compared with students who had less sport experience. These students had a LBP prevalence rate of 71.7%. When examining LBP rates based on sport type, baseball was associated with the second highest LBP prevalence. Students who played baseball were 3.2 times more likely to experience LBP in their lifetime only behind volleyball which had an odds ratio of 3.8. The findings suggest that the asymmetric baseball postures and motions lead to asymmetric spondylosis in right and left sides and cause mechanical degeneration in intervertebral discs (Hangai et al., 2010).

Epidemiological data from the National Collegiate Athletic Association (NCAA) Surveillance System were derived from baseball players aged 16 years or older from the time frame of 1988–2004 (Dick et al., 2007). Among the three NCAA Divisions, the highest injury rate existed in Division I men's baseball, which may be due to a more accurate collection of injuries compared with Divisions II and III. Players were three times more likely to suffer an injury during a game than during practice. The trunk/back comprised 15% of the total injuries in collegiate baseball. The player activity/positions with the highest reported incidence of trunk/back injuries were the pitcher and batter (Dick et al., 2007).

Professional baseball players

Few studies have focused on the injuries incurred by Major League Baseball (MLB) players. One retrospective study of publically available data on MLB players found that disc herniation's occurred at a higher frequency during pitching and batting (27.6% and 38%) compared with players playing their position on the infield and in the outfield (20% and 14.4%; Earhart et al., 2012). Two additional retrospective studies gathered their data from publically available disabled lists and injury lists (Li et al., 2013; Posner et al., 2011). Posner et al. found that among the injured players, 7.8% and 7.4% of the fielders and pitchers, respectively, developed spine and back injuries. Other injuries in the structures that support the pelvis, hip and spine also occurred. For example, groin, hamstring and core injuries occur in 3.8–7.6% of players playing their position in the field and pitchers (Posner et al., 2011). Rates of back injury were similar between players that had been injured participating in games or practice (8.3% and 11.5%; Dick et al., 2007).

Injury types relevant to LBP in baseball players

Baseball throwing involves asymmetric spine loading, lateral bending and spinal flexion away from the throwing arm, and transverse plane motion to the opposite side with high rotational velocity. (Laudner et al., 2013) Evidence shows that sport motions that combine biomechanical stressors place significant loading on the vertebrae, including the pars interarticularis (Chosa, Totoribe, & Tajima, 2004; Crewe, Campbell, Elliott, & Alderson, 2013). Injury types that have a mechanical etiology include stress reactions and fractures, intervertebral disc degeneration and herniation, as well as strain of the lumbar region muscle/soft tissues. Injury severity is often dependent as to the chronicity of the current injury the player had sustained. Untreated chronic conditions have the potential to become a more significant injury, including stress fractures and spondylolysis.

Stress fractures

Stress reactions may occur as a consequence of overuse with repetitive mechanical stress that weakens the pars interarticularis and pedicles of the vertebrae. If untreated, stress reactions can lead to stress fractures and spondylolysis. Stress fractures (e.g. pars defects) occur in approximately 3.3% of high school baseball players (Changstrom, Brou, Khodae, Braund, & Comstock, 2015). There is general agreement that repetitive motions involving lumbar rotation and extension promote spondylolysis. (Chosa et al., 2004; Pierre A. d'Hemecourt, Zurakowski, Kriemler, & Micheli, 2002; Laudner et al., 2013) There are case reports of pedicle stress reactions and stress fractures in adolescent pitchers. (Button &

Petron, 2007; Sairyo et al., 2003) Case series have reported development of painful pars defects in baseball pitchers serious enough to warrant surgical repair (Higashino et al., 2007). In some instances, LBP and spondylosis on one side may cause the contralateral side to absorb more stress during baseball motions, thereby leading to spondylolysis in the contralateral pars interarticularis (Tezuka, Sairyo, Sakai, & Dezawa, 2014).

Vertebral disc degeneration

Lumbar disc degeneration can be associated with chronic and recurrent LBP (Hancock et al., 2015). Among Japanese baseball players (19.8 ± 0.9 years of age), 59.7% of the sample had radiographic disc degeneration at one or more levels in the lumbar spine (Hangai et al., 2009). The majority of the degenerative processes occurred at levels L4/L5 and L5/S1 (57.9% of those with degeneration). The incidence of degeneration in baseball was highest of the sports studied, including swimming, basketball, kendo, soccer and running (Hangai et al., 2009).

Potential mechanical mechanisms underlying LBP

Baseball motions, such as throwing and hitting, involve high velocity, complex sequencing of body segment rotations that require substantial stability and motor control. The lumbar spine experiences large compression, shear and rotational moments (Stefanakis, Luo, Pollintine, Dolan, & Adams, 2014). Factors that interfere with this sequencing or stability and control include dynamic lumbar or core strength deficits, restricted range of motion (ROM) about the thoracolumbar spine, as well as improper timing of segment motions. Similar to other sports like golf, improper motion mechanics can magnify these forces and lead to LBP (Tsai et al., 2010). Figure 1 provides an overview of major mechanical risk factors that may lead to LBP if not properly addressed. It is likely that the development and persistence of LBP is bidirectional; as mechanical mechanisms contribute to pain onset, the presence of pain may also interfere with normal mechanics. These mechanical forces may accumulate gradually with long playing history and mechanical stress exposure such that the LBP becomes chronic or disabling.

Dynamic strength of core and hip

Dynamic stability of the low back is important for injury prevention. A stable low back and core permits the proximal musculature to contract against it to develop force to be transferred to the distal body segments during throwing and batting (Kibler, Press, & Sciascia, 2006). Core musculature can be described as a set of 29 pairs of muscles that stabilize the lumbopelvic region (spine and pelvis) as well as help assisting the transfer of energy up the kinetic chain (Faries & Greenwood, 2007). With LBP, the paraspinals and multifidus are involved and show signs of atrophy and fatty infiltration (Alaranta, Tallroth, Soukka, & Heliövaara, 1993; Kader, Wardlaw, & Smith, 2000; Pillastrini et al., 2015; Wallwork, Stanton, Freke, & Hides, 2009). Atrophied muscles have a delay in the contractile activity prior to spinal rotation (Hides, Richardson, & Jull, 1996; Hodges & Richardson, 1996; Kader et al., 2000). Maintenance of stability during normal movement and loading patterns of throwing or hitting would be challenged (Panjabi, 2003).

In the motion of a baseball swing, the trunk initially rotates posteriorly over the pelvis in preparation to swing, and then forward at ball contact and follow-through (shoulder-to-pelvis crossover). This crossover is believed to place high stress on the spinal visco-elastic structures (thoracolumbar fascia, ligaments, disc and capsule) and push the physiologic limits of these tissues (Gluck, Bendo, & Spivak, 2008). Players with inadequate dynamic strength to control rotation at high speeds may accrue microtrauma to these soft tissues, which leads to onset of LBP (Dennis, Finch, McIntosh, & Elliott, 2008).

Dynamic strength is particularly important during phases of the motions that involve a weight shift from 1 foot to the other, during one-legged support phases of movement and during rapid acceleration phases of segment rotations (MacWilliams, Choi, Perezous, Chao, & McFarland, 1998; Ohta et al., 2015). Core musculature needs to be appropriately activated to maintain lumbopelvic stability during the forward step in hitting or the transfer of body weight to the leading foot to follow-through during throwing (Tsai et al., 2010). In addition, trunk extension, which has been shown to be controlled by eccentric contraction of the hips and spine flexor muscles and thoracolumbar fascia, essentially prepares the muscles in expectancy of a pitch or throw (Watkins et al., 1989; Young, Casazza, Press, & Herring, 1998). Upon follow-through after release of the ball, the spine should be maximally flexed with the upper trunk rotated. In a similar fashion, when hitting there is pelvis and upper trunk rotation towards the pitcher immediately prior to ball contact, and after ball contact the trunk axial rotation increases towards the pitcher (G. S. Fleisig, Hsu, Fortenbaugh, Cordover, & Press, 2013).

Potential deficits in muscle activation in baseball players have been studied. Electromyographic studies of muscles that support the pelvis in baseball pitchers have revealed that the gluteus maximus and biceps femoris muscles of the trailing and stride legs during a pitch are activated at approximately 125–170% of the amplitude as during a maximal isometric contractile force (Campbell, Stodden, & Nixon, 2010). Among collegiate pitchers, approximately 50% of pitchers had poor lumbopelvic motor control as demonstrated by the inability to stabilize during hip external and internal rotation motion (Chaudhari et al., 2014).

Timing of kinematic events

The pitching motion involves several key segment rotational events that should occur within specific time frames of the pitching cycle (Seroyer et al., 2010). The throwing cycle can be defined from the point of lead foot contact (0%) with the ground to ball release (100%). Maximal pelvic angular velocity of ~400–700°/s occurs between 28% and 35% of the throwing motion. Upper torso maximal rotational velocity occurs at 47–53% of the throwing motion, with a separation between the two motions ranging from 18% to 22% of the throwing cycle (Fortenbaugh D., n.d.). Maximal trunk axial rotation should occur near foot contact; this point in the pitching cycle is also when maximal trunk translational acceleration peaks (G. S. Fleisig et al., 2013).

The bat swing cycle similarly has several key sequential segmental rotational events and may be defined as the point of lead foot ground contact (0%) to ball separation from the bat (100%). Between the swing cycle start and finish, other important timing points should be

noted such as peak pelvis, trunk, shoulder and bat rotational velocities. There are no normative ranges for which these events occur. However, these events should occur sequentially along the kinematic chain with peak pelvic rotational velocity occurring first, then trunk, shoulder and finally bat. Maximal trunk axial angular acceleration in pitching ($11,600^\circ/\text{s}$) and batting ($7,200^\circ/\text{s}$) occur at lead foot contact and right after ball contact, respectively (Fleisig et al., 2013). The most physically challenging instant for the trunk and spine was found to be approximately at lead foot contact for pitching and after ball contact for batting (Fleisig, Chu, Weber, & Andrews, 2009; Fleisig et al., 2013). Early or late timing in the respective kinematic phases of throwing or hitting forces body segments distal in the kinematic chain to compensate and generate higher muscular forces to maintain overall ball or bat speed.

ROM and flexibility

ROM of the lumbar spine in flexion, extension, rotation and lateral bending may reveal issues that can lead to back pain (Purcell & Micheli, 2009). Athletes are subject to moving in all directions, however, common movements in baseball such as spinal flexion, rotation and lateral bending can shift focus of injury prevalence to potential ROM issues. Supporting muscles and connective tissues may affect the ROM of the spine. Limited hip and lumbar rotational and lateral motion may hinder performance but also be a cause of core strains during the high velocity movements of players. It should be noted that deficits or excessive ROM can contribute to the onset of musculoskeletal pain, however high amounts of flexibility and ROM can aid in performance.

Thoracolumbar ROM measurements of collegiate and minor league baseball pitchers are different compared with position players as well as general populations (Laudner et al., 2013). Pitchers have a significantly higher thoracolumbar rotational ROM to the non-throwing side as compared with their dominant throwing side (throwing side: $48.8^\circ \pm 6.4^\circ$ vs. **non-throwing side:** $51.9^\circ \pm 6.6^\circ$; Laudner et al., 2013). Baseball players have larger passive trunk ROM for extension and rotation to both throwing and non-throwing sides compared with the normative numbers (American Medical Association, 2001).

Synopsis of the presented evidence

Existing evidence regarding LBP in baseball is derived from cross-sectional mixed samples of active people, previously active people and public sources. The actual prevalence of LBP and related spine injuries in collegiate and MLB populations may be underestimated due to the qualifications of the disabled list in professional baseball and the NCAA Surveillance System in collegiate baseball (Dick et al., 2007a; Changstrom et al., 2015). With respect to the qualifications of the MLB disabled list, there is not a requirement to place a player on the disabled list with an injury such as a muscle strain (potentially in the case of a player exhibiting LBP). Many times, players exhibiting symptoms such as LBP usually find that the issue resolves itself before a consideration of placing them on the disabled list presents itself. Hence, a player can miss time due to LBP symptoms and not have it reported because they did not get placed on the MLB disabled list. LBP symptoms are not characterized well within each study or are not consistently characterized among studies. Despite the relatively

low quality of the evidence, it can be gleaned that the risk for lifetime incidence LBP is higher among persons who have played baseball compared with most other sports (Hangai et al., 2010; Schmidt et al., 2014). Moreover, long-term exposure to playing baseball was related to greater risk of developing LBP. Similar to other competitive sports, the risk for LBP and injury was greater during game play compared with practice.

Additional qualitative and quantitative data of these injuries including the intensity, type (sharp, aching, throbbing, intermittent) and specific region of the low back (e.g. sacroiliac joint, lumbar spine) would provide insight about the symptom burden and the specific areas at higher risk for injury in baseball players. Prospective evidence of LBP in baseball players is currently lacking. Prospective Level II studies of the incidence of lumbar spine injuries from preseason to the end of season would provide the temporal patterns of greater injury susceptibility. Documentation of pain onset, symptom severity and variations on training volume, type and/or time during the season would capture patterns of activity or stimuli related to worsening LBP. The importance of controlling LBP has major implications for independent living and maintenance of mobility. Although the careers of baseball players are finite, the lives of these players will continue past retirement. While baseball players have long lifespans (Abel & Kruger, 2005, 2006), they will also have a longer poorer quality of life due to baseball-related musculoskeletal degeneration with knee and finger arthritis being the two most reported (Dick et al., 2007; Meir, Weatherby, & Rolfe, 2007). As such, screening and prevention programmes LBP would provide important information for this population.

Conclusion

The available quantity and quality of evidence focusing on LBP and related injuries in baseball is low, but indicates up to 15% of players experience pain. (Dick et al., 2007) More serious injuries have been reported in baseball players pertaining to LBP, such as stress fractures and degenerative disc disease. Moreover, further research on this topic is needed to identify baseball activities that may specifically contribute to the pathology of LBP and other chronic low back injuries. Independent or coupled mechanisms such as aberrant motions, muscular strength deficits or suboptimal sequence of motion events may contribute LBP and related pathologies in the baseball population. Long-term tracking of performance, physical function and sport **motion with LBP is essential to** understand the pain impact on the player. Additional evidence of the activities that exacerbate or lessen LBP is needed to help players and their care teams develop specific treatment and training protocols.

References

- Abel EL, Kruger ML. Longevity of major league baseball players. *Research in Sports Medicine* (Print). 2005; 13(1):1–5. DOI: 10.1080/15438620590922031 [PubMed: 16389882]
- Abel EL, Kruger ML. The healthy worker effect in major league baseball revisited. *Research in Sports Medicine* (Print). 2006; 14(1):83–87. DOI: 10.1080/15438620500528406 [PubMed: 16700406]
- Alaranta H, Tallroth K, Soukka A, Heliövaara M. Fat content of lumbar extensor muscles and low back disability: A radiographic and clinical comparison. *Journal of Spinal Disorders*. 1993; 6(2):137–140. [PubMed: 8504225]
- American Medical Association. *Guides to the evaluation of permanent impairment*. Atlanta, GA: AMA Press; 2001.

- Anderson K, Sarwark JF, Conway JJ, Logue ES, Schafer MF. Quantitative assessment with SPECT imaging of stress injuries of the pars interarticularis and response to bracing. *Journal of Pediatric Orthopedics*. 2000; 20(1):28–33. [PubMed: 10641684]
- Blanda J, Bethem D, Moats W, Lew M. Defects of pars interarticularis in athletes: A protocol for nonoperative treatment. *Journal of Spinal Disorders*. 1993; 6(5):406–411. [PubMed: 8274809]
- Bono CM. Low-back pain in athletes. *The Journal of Bone & Joint Surgery*. 2004; 86(2):382–396. [PubMed: 14960688]
- Button JH, Petron DJ. Pedicle stress reaction in a teenage baseball pitcher. *American Journal of Physical Medicine & Rehabilitation/Association of Academic Physiatrists*. 2007; 86(11):956.doi: 10.1097/PHM.0b013e3181583fcd
- Campbell BM, Stodden DF, Nixon MK. Lower extremity muscle activation during baseball pitching. *Journal of Strength and Conditioning Research/National Strength & Conditioning Association*. 2010; 24(4):964–971. DOI: 10.1519/JSC.0b013e3181cb241b
- Changstrom BG, Brou L, Khodae M, Braund C, Comstock RD. Epidemiology of stress fracture injuries among US high school athletes, 2005–2006 through 2012–2013. *The American Journal of Sports Medicine*. 2015; 43(1):26–33. DOI: 10.1177/0363546514562739 [PubMed: 25480834]
- Chaudhari AMW, McKenzie CS, Pan X, Oñate JA. Lumbopelvic control and days missed because of injury in professional baseball pitchers. *The American Journal of Sports Medicine*. 2014; 42(11):2734–2740. DOI: 10.1177/0363546514545861 [PubMed: 25159541]
- Chosa E, Totoribe K, Tajima N. A biomechanical study of lumbar spondylolysis based on a three-dimensional finite element method. *Journal of Orthopaedic Research: Official Publication of the Orthopaedic Research Society*. 2004; 22(1):158–163. DOI: 10.1016/S0736-0266(03)00160-8 [PubMed: 14656675]
- Crewe H, Campbell A, Elliott B, Alderson J. Lumbo-pelvic biomechanics and quadratus lumborum asymmetry in cricket fast bowlers. *Medicine and Science in Sports and Exercise*. 2013; 45(4):778–783. DOI: 10.1249/MSS.0b013e31827973d1 [PubMed: 23135366]
- d’Hemecourt PA, Gerbino PG, Micheli LJ. Back injuries in the young athlete. *Clinics in Sports Medicine*. 2000; 19(4):663–679. [PubMed: 11019734]
- d’Hemecourt PA, Zurakowski D, Kriemler S, Micheli LJ. Spondylolysis: Returning the athlete to sports participation with brace treatment. *Orthopedics*. 2002; 25(6):653–657. [PubMed: 12083575]
- Dennis RJ, Finch CF, McIntosh AS, Elliott BC. Use of field-based tests to identify risk factors for injury to fast bowlers in cricket. *British Journal of Sports Medicine*. 2008; 42(6):477–482. DOI: 10.1136/bjism.2008.046698 [PubMed: 18390922]
- Dick R, Agel J, Marshall SW. National collegiate athletic association injury surveillance system commentaries: Introduction and methods. *Journal of Athletic Training*. 2007; 42(2):173–182. [PubMed: 21714302]
- Dick R, Sauers EL, Agel J, Keuter G, Marshall SW, McCarty K, McFarland E. Descriptive epidemiology of collegiate men’s baseball injuries: National collegiate athletic association injury surveillance system, 1988–1989 through 2003–2004. *Journal of Athletic Training*. 2007; 42(2):183–193. [PubMed: 17710166]
- Earhart JS, Roberts D, Roc G, Gryzlo S, Hsu W. Effects of lumbar disk herniation on the careers of professional baseball players. *Orthopedics*. 2012; 35(1):43–49. DOI: 10.3928/01477447-20111122-40 [PubMed: 22229920]
- Faries MD, Greenwood M. Stabilizing the confusion. *Strength & Cond Journal*. 2007; 29(2):10–25.
- Fleisig G, Chu Y, Weber A, Andrews J. Variability in baseball pitching biomechanics among various levels of competition. *Sports Biomechanics/International Society of Biomechanics in Sports*. 2009; 8(1):10–21. DOI: 10.1080/14763140802629958
- Fleisig GS, Hsu WK, Fortenbaugh D, Cordover A, Press JM. Trunk axial rotation in baseball pitching and batting. *Sports Biomechanics/International Society of Biomechanics in Sports*. 2013; 12(4):324–333. DOI: 10.1080/14763141.2013.838693
- Fortenbaugh, D. The biomechanics of the baseball swing (dissertation). University of Miami Scholarly Repository; n.d.
- Gluck GS, Bendo JA, Spivak JM. The lumbar spine and low back pain in golf: A literature review of swing biomechanics and injury prevention. *The Spine Journal: Official Journal of the North*

- American Spine Society. 2008; 8(5):778–788. DOI: 10.1016/j.spinee.2007.07.388 [PubMed: 17938007]
- Hancock MJ, Maher CM, Petocz P, Lin C-WC, Steffens D, Luque-Suarez A, Magnussen JS. Risk factors for a recurrence of low back pain. *The Spine Journal: Official Journal of the North American Spine Society*. 2015; doi: 10.1016/j.spinee.2015.07.007
- Hangai M, Kaneoka K, Hinotsu S, Shimizu K, Okubo Y, Miyakawa S, ... Ochiai N. Lumbar intervertebral disk degeneration in athletes. *The American Journal of Sports Medicine*. 2009; 37(1):149–155. DOI: 10.1177/0363546508323252 [PubMed: 18799691]
- Hangai M, Kaneoka K, Okubo Y, Miyakawa S, Hinotsu S, Mukai N, ... Ochiai N. Relationship between low back pain and competitive sports activities during youth. *The American Journal of Sports Medicine*. 2010; 38(4):791–796. DOI: 10.1177/0363546509350297 [PubMed: 20051500]
- Hides JA, Richardson CA, Jull GA. Multifidus muscle recovery is not automatic after resolution of acute, first-episode low back pain. *Spine*. 1996; 21(23):2763–2769. [PubMed: 8979323]
- Higashino K, Sairyo K, Katoh S, Sakai T, Kosaka H, Yasui N. Minimally invasive technique for direct repair of the pars defects in young adults using a spinal endoscope: A technical note. *Minimally Invasive Neurosurgery*. 2007; 50(3):186–190. DOI: 10.1055/s-2007-982511
- Hodges PW, Richardson CA. Inefficient muscular stabilization of the lumbar spine associated with low back pain. A motor control evaluation of transversus abdominis. *Spine*. 1996; 21(22):2640–2650. [PubMed: 8961451]
- Kader DF, Wardlaw D, Smith FW. Correlation between the MRI changes in the lumbar multifidus muscles and leg pain. *Clinical Radiology*. 2000; 55(2):145–149. DOI: 10.1053/crad.1999.0340 [PubMed: 10657162]
- Kibler WB, Press J, Sciascia A. The role of core stability in athletic function. *Sports Medicine (Auckland, NZ)*. 2006; 36(3):189–198.
- Ladenhauf HN, Fabricant PD, Grossman E, Widmann RF, Green DW. Athletic participation in children with symptomatic spondylolysis in the New York area. *Medicine and Science in Sports and Exercise*. 2013; 45(10):1971–1974. DOI: 10.1249/MSS.0b013e318294b4ed [PubMed: 23559123]
- Laudner K, Lynall R, Williams JG, Wong R, Onuki T, Meister K. Thoracolumbar range of motion in baseball pitchers and position players. *International Journal of Sports Physical Therapy*. 2013; 8(6):777–783. [PubMed: 24377063]
- Li X, Zhou H, Williams P, Steele JJ, Nguyen J, Jäger M, Coleman S. The epidemiology of single season musculoskeletal injuries in professional baseball. *Orthopedic Reviews*. 2013; 5(1):e3.doi: 10.4081/or.2013.e3 [PubMed: 23705061]
- MacWilliams BA, Choi T, Perezous MK, Chao EYS, McFarland EG. Characteristic ground-reaction forces in baseball pitching. *The American Journal of Sports Medicine*. 1998; 26(1):66–71. [PubMed: 9474404]
- Meir RA, Weatherby RP, Rolfe MI. A preliminary investigation into the long-term injury consequences reported by retired baseball players. *Journal of Science and Medicine in Sport/Sports Medicine Australia*. 2007; 10(3):187–190. DOI: 10.1016/j.jsams.2006.06.002
- Miller SF, Congeni J, Swanson K. Long-term functional and anatomical follow-up of early detected spondylolysis in young athletes. *The American Journal of Sports Medicine*. 2004; 32(4):928–933. [PubMed: 15150039]
- Ohta Y, Nakamoto H, Ishii Y, Ikudome S, Takahashi K, Shima N, Ren L. Muscle activation characteristics of the front leg during baseball swings with timing correction for sudden velocity decrease. *Plos One*. 2015; 10(4):e0124113.doi: 10.1371/journal.pone.0124113 [PubMed: 25918848]
- Oliver GD, Keeley DW. Gluteal muscle group activation and its relationship with pelvis and torso kinematics in high-school baseball pitchers. *Journal of Strength and Conditioning Research/ National Strength & Conditioning Association*. 2010; 24(11):3015–3022. DOI: 10.1519/JSC.0b013e3181c865ce
- Panjabi MM. Clinical spinal instability and low back pain. *Journal of Electromyography and Kinesiology: Official Journal of the International Society of Electrophysiological Kinesiology*. 2003; 13(4):371–379. [PubMed: 12832167]

- Petering RC, Webb C. Treatment options for low back pain in athletes. *Sports Health*. 2011; 3(6):550–555. DOI: 10.1177/1941738111416446 [PubMed: 23016058]
- Pillastrini P, Ferrari S, Rattin S, Cupello A, Villafañe JH, Vanti C. Exercise and tropism of the multifidus muscle in low back pain: A short review. *Journal of Physical Therapy Science*. 2015; 27(3):943–945. DOI: 10.1589/jpts.27.943 [PubMed: 25931765]
- Posner M, Cameron KL, Wolf JM, Belmont PJ, Owens BD. Epidemiology of major league baseball injuries. *The American Journal of Sports Medicine*. 2011; 39(8):1676–1680. DOI: 10.1177/0363546511411700 [PubMed: 21709023]
- Purcell L, Micheli L. Low back pain in young athletes. *Sports Health*. 2009; 1(3):212–222. DOI: 10.1177/1941738109334212 [PubMed: 23015874]
- Sairyo K, Katoh S, Sakamaki T, Komatsubara S, Endo K, Yasui N. Three successive stress fractures at the same vertebral level in an adolescent baseball player. *The American Journal of Sports Medicine*. 2003; 31(4):606–610. [PubMed: 12860553]
- Schmidt CP, Zwillingenberger S, Walther A, Reuter U, Kasten P, Seifert J, ... Stiehler M. Prevalence of low back pain in adolescent athletes - an epidemiological investigation. *International Journal of Sports Medicine*. 2014; 35(8):684–689. DOI: 10.1055/s-0033-1358731 [PubMed: 24424960]
- Seroyer ST, Nho SJ, Bach BR, Bush-Joseph CA, Nicholson GP, Romeo AA. The kinetic chain in overhand pitching: Its potential role for performance enhancement and injury prevention. *Sports Health*. 2010; 2(2):135–146. DOI: 10.1177/1941738110362656 [PubMed: 23015931]
- Stefanakis M, Luo J, Pollintine P, Dolan P, Adams MA. ISSLS prize winner: Mechanical influences in progressive intervertebral disc degeneration. *Spine*. 2014; 39(17):1365–1372. DOI: 10.1097/BRS.0000000000000389 [PubMed: 24831499]
- Tezuka F, Sairyo K, Sakai T, Dezawa A. Etiology of adult-onset stress fracture in the lumbar spine. *Journal of Spinal Disorders & Techniques*. 2014; doi: 10.1097/BSD.0000000000000162
- Tsai YS, Sell TC, Smoliga JM, Myers JB, Learman KE, Lephart SM. A comparison of physical characteristics and swing mechanics between golfers with and without a history of low back pain. *The Journal of Orthopaedic and Sports Physical Therapy*. 2010; 40(7):430–438. DOI: 10.2519/jospt.2010.3152 [PubMed: 20592479]
- Wallwork TL, Stanton WR, Freke M, Hides JA. The effect of chronic low back pain on size and contraction of the lumbar multifidus muscle. *Manual Therapy*. 2009; 14(5):496–500. DOI: 10.1016/j.math.2008.09.006 [PubMed: 19027343]
- Watkins RG, Dennis S, Dillin WH, Schnebel B, Schneiderman G, Jobe F, ... Pink M. Dynamic EMG analysis of torque transfer in professional baseball pitchers. *Spine*. 1989; 14(4):404–408. [PubMed: 2524110]
- Young, J., Casazza, B., Press, J., Herring, S. Biomechanical aspects of the spine in pitching. In: Andrews, JR, Zarins, B., Wilk, KE., editors. *Injuries in baseball*. Philadelphia, PA: Lippincott-Raven; 1998. p. 23–35.
- Young JL, Herring SA, Press JM, Casazza BA. The influence of the spine on the shoulder in the throwing athlete. *Journal of Back and Musculoskeletal Rehabilitation*. 1996; 7(1):5–17. DOI: 10.3233/BMR-1996-7103 [PubMed: 24572551]

Figure 1.
Proposed key mechanisms of low back pain in baseball players.