Published in final edited form as:

Behav Brain Sci. 2016 January ; 39: e201. doi:10.1017/S0140525X15001727.

Glutamate and Norepinephrine Interaction: Relevance to Higher Cognitive Operations and Psychopathology

Chadi G. Abdallah, MD^{1,2}, Lynnette A. Averill, PhD^{1,2}, John H. Krystal, MD^{1,2}, Steven M. Southwick, MD^{1,2,*}, and Amy F.T. Arnsten, PhD^{1,3}

¹Clinical Neurosciences Division, VA National Center for PTSD, VA Connecticut Healthcare System, 950 Campbell Avenue, 151E, West Haven, CT, USA, 06516

²Department of Psychiatry, Yale University School of Medicine, 300 George Street, Suite 901, New Haven, CT, USA, 06511

³Department of Neurobiology, Yale University School of Medicine, 200 South Frontage Road, SHM C-303, New Haven, CT, USA, 06510

Abstract

Mather and colleagues present an impressive interdisciplinary model of arousal-induced norepinephrine (NE) release and its role in selectively enhancing/inhibiting perception, attention, and memory consolidation. This model will require empirical investigation to test its validity and generalizability beyond classic NE circuits as it simplifies extremely complex and heterogeneous actions including NE mechanisms related to higher cognitive circuits and psychopathology.

In the current issue, Mather and colleagues propose a molecular model, termed Glutamate Amplifies Noradrenergic Effects (GANE), through which arousal enhances or inhibits perception, attention, and memory consolidation (Mather, Clewett, Sakaki, & Harley, 2015). In this model, arousal precipitates phasic release of norepinephrine (NE) throughout the brain, but “hot spots” of NE release are generated near activated glutamate circuits, sufficient to engage low affinity β adrenoceptors, which further increase glutamate release and enhance post-synaptic plasticity by increasing cAMP signaling. The model provides an impressive integration across several fields, but will require empirical investigation to test its validity. Furthermore, although the model is presented as universally applicable throughout brain, NE actually has very heterogeneous actions in different brain circuits. In particular, although the GANE model addresses the effects of normal arousal mechanisms in sensory cortex and hippocampus, it is important to discuss how this model may relate to NE actions in higher cognitive circuits, and to conditions of psychopathology.

The noradrenergic system plays an essential role in the pathophysiology and treatment of psychiatric disorders. For example, noradrenergic dysregulation is associated with posttraumatic stress disorder (PTSD), and $\alpha 1$ antagonists can reduce these symptoms (A. F. T. Arnsten, Raskind, Taylor, & Connor, 2015; Southwick et al., 1999). Many antidepressants

*Corresponding Author: Steven Southwick, MD, Clinical Neurosciences Division, VA National Center for PTSD, VA Connecticut Healthcare System, 950 Campbell Avenue, 151E, West Haven, CT, USA, 06516, steven.southwick@yale.edu; 203-983-5711, ext. 2464.

target the noradrenergic system (Klimek et al., 1997), and $\alpha 2A$ agonists enhance cognition in patients with attention deficit hyperactivity disorder (ADHD) (A. F. T. Arnsten & Wang, 2016). Similarly, accumulating evidence implicates glutamate in the etiology and treatment of mental disorders (Chambers et al., 1999; Krystal, Sanacora, & Duman, 2013). It is not clear whether the GANE model applies to traumatic stress conditions, as the research cited by Mather et al. utilized subtle arousing conditions, e.g. an emotional word. However, it is likely to explain several aspects of PTSD, e.g. enhancing the consolidation of traumatic events that may contribute to flashbacks and intrusive memories. However, additional, higher brain changes during trauma may not be captured by this model, as NE actions in brain are more heterogeneous than described.

Most important for human cognition, the newly evolved circuits in layer III of the dorsolateral prefrontal cortex (dlPFC) that underlie higher cognitive operations are modulated in a unique manner, that is often opposite to classic synapses in sensory cortex, amygdala and hippocampus (A. F. T. Arnsten, Wang, & Paspalas, 2012). Indeed, these newly evolved “Delay cell” circuits in the dlPFC are even regulated differently than sensory/response-related neurons within the dlPFC. For example, Delay cell persistent firing is mediated by NMDAR with NR2B subunits that are exclusively in the post-synaptic density, not extra-synaptic they are in classic synapses (Wang et al., 2013). Furthermore, Delay cells are only subtly influenced by AMPA receptors, and show reduced, rather than increased, neuronal firing following systemic ketamine (Wang et al., 2013). In contrast, Response feedback cells in dlPFC (likely layer V) show a more classic profile, with large AMPA receptor influences and increased firing with systemic ketamine (Wang et al., 2013). These marked differences extend to intracellular cAMP signaling events as well. In classic synapses, activation of cAMP signaling, e.g. arising from β -adrenoceptor stimulation, increases glutamate release from axon terminals and strengthens LTP post-synaptically. However, in layer III dlPFC circuits, increased cAMP signaling *weakens* connections by opening cAMP-PKA regulated potassium channels in dendritic spines (A. F. Arnsten, 2015; A. F. T. Arnsten et al., 2012). Instead, it is inhibition of cAMP signaling via post-synaptic $\alpha 2A$ -adrenoceptors that strengthens network connectivity by closing potassium channels near the synapse (Wang et al., 2007). There is currently no evidence of NE “hot spots” in these circuits; e.g. blockade of β receptors within the primate dlPFC has no effect on working memory performance (Li & Mei, 1994), even though there are likely high levels of glutamate release in dlPFC arising from the persistent firing of these neuronal networks. Thus, the model shown in Figure 6 of Mather et al. is misleading, as it does not differentiate NE actions in classic synapses versus those in more newly evolved dlPFC circuits.

Mather et al. also provide an over-simplified discussion of NE actions at $\alpha 1$ adrenoceptors. Although they focus on $\alpha 1$ mechanisms that weaken plasticity, there are many synapses where $\alpha 1$ promotes synaptic actions, e.g. in somatosensory cortex (Mouradian, Seller, & Waterhouse, 1991; Waterhouse, Moises, & Woodward, 1981; Waterhouse, Mouradian, Sessler, & Lin, 2000). There are also key circuits where $\alpha 1$ receptor activation potentiates beta receptor actions, e.g. in amygdala, where $\alpha 1$ receptors facilitate β -adrenergic enhancement of memory consolidation (Ferry, Roozendaal, & McGaugh, 1999a, 1999b). These effects are opposite to those described by Mather and colleagues. Their model also does not capture the important finding that high levels of NE release in PFC during stress

decrease persistent firing and working memory abilities through stimulation of $\alpha 1$ receptors (Birnbaum et al., 2004). All of these actions likely have a key effect in switching control of behavior from thoughtful, flexible top-down control by PFC under conditions of safety (moderate levels of arousal), to reflexive, unconscious habits mediated by sensorimotor cortex and subcortical structures during uncontrollable stress (very high levels of arousal).

These mechanisms have particular relevance to the symptoms of PTSD, where there is extensive evidence of elevated noradrenergic activity (Southwick et al., 1999). For example, the $\alpha 2$ antagonist, yohimbine, worsens symptoms and induces hypofrontality in subjects with PTSD at doses that have little effect in control subjects (Bremner et al., 1997; Southwick et al., 1993). These drug actions may arise from a combination of neural events, e.g. loss of dlPFC top-down control from blockade of post-synaptic $\alpha 2A$ receptors and increased NE stimulation of $\alpha 1$ receptors in dlPFC, as well as increased NE release in “hot spots” in amygdala, hippocampus and sensory cortex that may exacerbate anxiety and flashbacks (A. F. T. Arnsten et al., 2015). Thus, the GANE model may apply to NE actions in classic brain circuits, but not to higher cortical circuits where circuits are strengthened by $\alpha 2A$ - rather than β -adrenoceptor mechanisms.

References

- Arnsten AF. Stress weakens prefrontal networks: molecular insults to higher cognition. *Nat Neurosci.* 2015; 18:1376–1385. [PubMed: 26404712]
- Arnsten AFT, Raskind M, Taylor FB, Connor DF. The effects of stress exposure on prefrontal cortex: Translating basic research into successful treatments for Post-Traumatic Stress Disorder. *Neurobiology of Stress.* 2015; 1:89–99. [PubMed: 25436222]
- Arnsten AFT, Wang M. Targeting Prefrontal Cortical Systems for Drug Development: Potential Therapies for Cognitive Disorders. *Annual Rev Pharmacology Toxicology.* 2016; 56 epub January 6, 2016.
- Arnsten AFT, Wang M, Paspalas CD. Neuromodulation of thought: Flexibilities and vulnerabilities in prefrontal cortical network synapses. *Neuron.* 2012; 76(1):223–239. [PubMed: 23040817]
- Birnbaum SB, Yuan P, Wang M, Vijayraghavan S, Bloom A, Davis D, Arnsten AFT. Protein kinase C overactivity impairs prefrontal cortical regulation of working memory. *Science.* 2004; 306:882–884. [PubMed: 15514161]
- Bremner JD, Innis RB, Ng CK, Staib LH, Salomon RM, Bronen RA, Charney DS. Positron emission tomography measurement of cerebral metabolic correlates of yohimbine administration in combat-related posttraumatic stress disorder. *Arch Gen Psychiatry.* 1997; 54:246–254. [PubMed: 9075465]
- Chambers RA, Bremner JD, Moghaddam B, Southwick SM, Charney DS, Krystal JH. Glutamate and post-traumatic stress disorder: toward a psychobiology of dissociation. *Semin Clin Neuropsychiatry.* 1999; 4:274–281. [PubMed: 10553033]
- Ferry B, Roozendaal B, McGaugh JL. Basolateral amygdala noradrenergic influences on memory storage are mediated by an interaction between beta- and alpha-1-adrenoceptors. *J Neurosci.* 1999a; 19:5119–5123. [PubMed: 10366644]
- Ferry B, Roozendaal B, McGaugh JL. Involvement of alpha-1-adrenoceptors in the basolateral amygdala in modulation of memory storage. *Eur J Pharmacol.* 1999b; 372:9–16. [PubMed: 10374709]
- Klimek V, Stockmeier C, Overholser J, Meltzer HY, Kalka S, Dille G, Ordway GA. Reduced levels of norepinephrine transporters in the locus coeruleus in major depression. *J Neurosci.* 1997; 17:8451–8458. [PubMed: 9334417]
- Krystal JH, Sanacora G, Duman RS. Rapid-acting glutamatergic antidepressants: the path to ketamine and beyond. *Biol Psychiatry.* 2013; 73:1133–1141. [PubMed: 23726151]

- Li BM, Mei ZT. Delayed response deficit induced by local injection of the alpha-2 adrenergic antagonist yohimbine into the dorsolateral prefrontal cortex in young adult monkeys. *Behav Neural Biol.* 1994; 62:134–139. [PubMed: 7993303]
- Mather M, Clewett D, Sakaki M, Harley CW. Norepinephrine ignites local hot spots of neuronal excitation: How arousal amplifies selectivity in perception and memory. *Behav Brain Sci.* 2015; 1:1–100.
- Mouradian RD, Seller FM, Waterhouse BD. Noradrenergic potentiation of excitatory transmitter action in cerebrocortical slices: evidence of mediation by an alpha1-receptor-linked second messenger pathway. *Brain Res.* 1991; 546:83–95. [PubMed: 1677306]
- Southwick SM, Bremner JD, Rasmusson A, Morgan CA, Arnsten A, Charney DS. Role of norepinephrine in the pathophysiology and treatment of posttraumatic stress disorder. *Biological Psychiatry.* 1999; 46:1192–1204. [PubMed: 10560025]
- Southwick SM, Krystal JH, Morgan CA, Johnson D, Nagy LM, Nicolaou A, Charney DS. Abnormal noradrenergic function in posttraumatic stress disorder. *Arch Gen Psychiatry.* 1993; 50:266–274. [PubMed: 8466387]
- Wang M, Ramos B, Paspalas C, Shu Y, Simen A, Duque A, Arnsten AFT. Alpha2A-adrenoceptor stimulation strengthens working memory networks by inhibiting cAMP-HCN channel signaling in prefrontal cortex. *Cell.* 2007; 129:397–410. [PubMed: 17448997]
- Wang M, Yang Y, Wang CJ, Gamo NJ, Jin LE, Mazer JA, Arnsten AF. NMDA receptors subserve working memory persistent neuronal firing in dorsolateral prefrontal cortex. *Neuron.* 2013; 77(4): 736–749. [PubMed: 23439125]
- Waterhouse BD, Moises HC, Woodward DJ. Alpha-receptor-mediated facilitation of somatosensory cortical neuronal responses to excitatory synaptic inputs and iontophoretically applied acetylcholine. *Neuropharmacology.* 1981; 20:907–920. [PubMed: 6272150]
- Waterhouse BD, Mouradian R, Sessler FM, Lin RC. Differential modulatory effects of norepinephrine on synaptically driven responses of layer V barrel field cortical neurons. *Brain Res.* 2000; 868:39–47. [PubMed: 10841886]