expectancy in Canada in 2001 was 79.5 years (77.0 for men and 82.0 for women). Canada has the seventh highest life expectancy for men and the 11th highest life expectancy for women, similar to other wealthy industrialised countries.

However, residents of Nunavik, the Inuit region of Quebec, have the lowest life expectancy in Canada, at 66.7 years (66.3 for men and 70.2 for women). Life expectancy in Nunavik falls between that of the Dominican Republic (67.0) and Egypt (66.5), ranked 111 and 112 respectively out of 191 countries. In 2001, the population of Nunavik was 9632, including 8760 people of aboriginal origin.

Statistics Canada also reported low life expectancies in various other northern and isolated parts of Canada. Smoking, heavy drinking, and higher mortality rates (partly due to suicides) are also prevalent in these communities.

Barbara Kermode-Scott *Calgary*

See www.statcan.ca/Daily/English/050201/d050201a.htm

Researchers to clone human stem cells to study motor neurone disease

The British research team that created Dolly the sheep has been granted a licence to clone human embryos to study motor neurone disease.

The licence will allow researchers at the Roslin Institute in Edinburgh and King's College in London to generate stem cell lines from embryos created using DNA from patients with motor neurone disease. They will focus on uncovering the cause of the motor neurone disease in patients whose condition cannot be linked to the genes already identified as causing the disease.

It is only the second time that the United Kingdom's Human Fertilisation and Embryology Authority has granted a licence for therapeutic cloning. The first licence was granted last year to the Newcastle Centre for Life,

where researches will use the stem cells they generate to research diabetes, among other diseases (*BMJ* 2004;329:417).

Under the latest licence, researchers will use cell nuclear replacement to grow motor neurone cells, enabling them to investigate what causes the cells to degenerate.

The cultured neurones will also be used to test the ability of new drugs to halt or reverse the disease process. Hundreds of thousands of drugs can be screened in cultured cells in a year for about £100 000 (\$186 000; €145 000), whereas it takes nearly two years and £20m to screen just one drug in patients.

Zosia Kmietowicz *London*

Asbestos related cancer deaths set to rise

Deaths from exposure to asbestos are increasing and will peak in the next decade, according to new research. Annual deaths from mesothelioma among men in Britain will rise to between 1950 and 2450 a year between 2011 and 2015, compared with 153 deaths in 1968, say researchers in the *British Journal of Cancer* (published online on 25 January ahead of print, www.nature.com/bjc; doi:10.1038/sj.bjc.6602307).

They predict that the total number of deaths since 1968 will rise to 90 000 by 2050, with 65 000 of those deaths after 2002.

In the study, researchers from the Health and Safety Executive, the London School of Hygiene and Tropical Medicine, and the Institute of Cancer Research used Poisson regression analysis to model male mesothelioma deaths from 1968 to 2001 and to predict numbers of male deaths in 2002-50.

The annual number of mesothelioma deaths in Britain has risen increasingly rapidly, with deaths 12 times higher in 2001 than in 1968, says the report. It says that men born in about 1940 have higher death rates than any previous or subsequent birth cohort.

Roger Dobson *Abergavenny*

Concerted effort needed to stop polio transmission in Asia by 2006

Ganapati Mudur *New Delhi*

The numbers of poliomyelitis cases in India, Pakistan, and Afghanistan almost halved last year. But stopping transmission of the poliovirus in Asia by the end of 2005 will require intensified immunisation campaigns in all three countries, public health officials said last week.

The three countries—containing Asia's last pockets of polio—reported 186 cases in 2004 compared with 336 in 2003. Health officials from the subcontinent met at the World Health Organization last week and outlined plans for 21 additional immunisation campaigns this year.

"This is the year to rid Asia of polio," said Bill Sargeant, chairman of Rotary International's PolioPlus Committee, a partner in the 16 year old global polio eradication initiative.

Polio is currently at its lowest ever levels in Asia. India reported 132 children paralysed by the poliovirus in 2004; Pakistan detected 50 and Afghanistan reported four.

But officials cautioned that "final success" would hinge on reaching "underserved children" in parts of Uttar Pradesh and Bihar in India. "Up to 10% of children are still being missed in the highest priority districts in these areas," said Steve Stewart,

a spokesman for the US Centers for Disease Control and Prevention in Atlanta, Georgia. He said that the quality of supplementary immunisation activities, which complement routine programmes, also needed to be improved in Pakistan's Sindh province and in southern Afghanistan.

But Thekkekarra Jacob John, former head of virology at the Christian Medical College in Vellore, has warned that persistent transmission of polio in Uttar Pradesh may be the result of a "high force of transmission" of the virus, brought about by low income and literacy, poor sanitation, and inadequate health infrastructure. "Conventional interventions that had worked in other poor communities may not work where the virus has a high force of transmission," he said.

Mr Stewart said that it was technically feasible to stop polio transmission in Asia by 2005 and in Africa by mid-2006. But in 2003-4 polio spread from Nigeria to 13 African countries and has re-established transmission in Burkina Faso, the Central African Republic, Chad, Côte d'Ivoire, and Sudan.

"These importations threaten the goal of ending transmission in Africa this year," he said. □

Children with polio play at the Amar Jyoti Rehabilitation Centre in Delhi, India