Barrier in West Bank threatens residents' health care, says report

Deborah Cohen *London*

Almost a third of West Bank Palestinian villages will be denied free and open access to health-care facilities once the Israeli government has completed constructing the West Bank barrier, says a report by the French non-governmental organisation Médecins du Monde.

In June 2002 the Israeli government began to build a barrier around the West Bank that, according to the Israel Defence Forces, will "lessen the impact and scope of terrorism on the citizens of Israel." Already 185 km long, it will eventually be 622 km long and consist of fences, ditches, patrol roads, and a concrete wall 8-9 m high costing an estimated \$4.7m (£2.5m; €3.6m) per km.

Worst hit will be the villages confined within the "seam zone," the territory between the green line (the 1949 armistice line that delineates Israel's pre-1967 border with the West Bank) and the barrier, according to the report. Villages situated in the enclaves formed by the route of the barrier will also be badly affected, it says. Palestinians wanting to leave these areas to travel to another town in the West Bank will have to apply for a permit

and pass through a checkpoint or gate guarded by a soldier.

People requiring emergency health treatment in Palestinian hospitals in other parts of the West Bank face an even longer journey than at present and may not be able to access emergency care at all during the night, according to Régis Garrigue, head of Médecins du Monde's Palestinian projects. He said: "The people inside the enclaves are in a big jail. Sometimes the passage into the enclaves is shut for a day or a night or even a week."

In Abu Dis and Aizaria, two Palestinian towns where the barrier has already been completed, the average time for an ambulance to travel to the nearest hospitals in Jerusalem has increased from about 10 minutes to over one hour and 50 minutes, according to the report. Mr Garrigue says that once the barrier is completed this problem will affect many more villages.

"If patients don't have access to their medical facilities or if medical staff cannot reach the facilities to provide medical services the healthcare system is bound to deteriorate," the report warns.

The extension of the barrier in the West Bank will lead to further difficulties for residents trying to get health care

Dr Yitzhak Sever, director of the Department of International Relations in the Ministry of Health, said that all the inhabitants on the west of the barrier, located within the seam zone, are given permits to travel to other Palestinian hospitals. More than 45 000 permits for patients were issued over the past year. He also stressed that Israeli doctors treat Palestinian patients and train health professionals, despite the former Palestinian Health Minister's refusal to sit on joint Palestinian-Israeli health committees after the

second intifada in 2000.

"Maybe the barrier will not be continued, we don't know. It's there for security reasons as determined by the Ministry of Defence. I know in the majority of places in the West Bank, building is frozen and the barrier will be under debate in the negotiations about disengagement and peace. We hope the new era will bring peace for the two peoples," he told the *BMJ*. □

The Ultimate Barrier: Impact of the Wall on the Palestinian Health Care System is available at www.healthandwall.org under "More info"

Findings from COX 2 studies are released

Scott Gottlieb *New York*

Results from recent studies that raised concerns about the cardiovascular safety of cyclo-oxygenase-2 (COX 2) inhibitors have been made available early, in advance of hearings on the drug class by the US Food and Drug Administration.

All the studies have been published online in the *New England Journal of Medicine* (<http://nejm.org>), and the journal lifted its normal embargo date for the studies because of what it described as the "possible public health implications."

The results came immediately before a three-day hearing

looking into the benefits and risks of COX 2 inhibitors that is being held by the US Food and Drug Administration's advisory committee, starting on 16 February. Among other things, the advisory committee is expected to evaluate recent data that suggest that, in addition to rofecoxib (Vioxx), other COX 2 inhibitors may be associated with an increased risk of cardiovascular events and to consider changes to the labelling of these medicines.

In one study, Solomon and colleagues reviewed all potentially serious cardiovascular events among 2035 patients with a history of colorectal neoplasia who were enrolled in a trial comparing two doses of celecoxib (200 mg or 400 mg twice daily) with placebo for the prevention of colorectal adenomas.

A cardiovascular event occurred in 7 of 679 (1.0%)

patients in the placebo group, compared with 16 of 685 (2.3%) patients receiving 200 mg of celecoxib twice daily (hazard ratio 2.3; 95% confidence interval 0.9 to 5.5) and with 23 of 671 (3.4%) patients receiving 400 mg of celecoxib twice daily (3.4; 1.4 to 7.8).

On the basis of these observations, the data and safety monitoring board recommended early discontinuation of the study drug.

In a second study, Bresalier and colleagues report the finding of an increased cardiovascular risk with rofecoxib that prompted the removal of that particular drug from the market by its manufacturer last September (*BMJ* 2004;329:816).

That trial—the APPROVe (adenomatous polyp prevention with Vioxx) trial—is being published for the first time, although the drug manufacturers, Merck,

took action on the findings last year. A total of 2586 patients with a history of colorectal adenomas were randomised: 1287 were assigned to receive 25 mg of rofecoxib daily, and 1299 to receive placebo.

A total of 46 patients in the rofecoxib group had a confirmed thrombotic event during 3059 patient years of follow up (1.50 events per 100 patient years), compared with 26 patients in the placebo group during 3327 patient years of follow up (0.78 events per 100 patient years); the corresponding relative risk was 1.92 (95% confidence interval 1.19 to 3.11; $P=0.008$).

A third study, by Nussmeier and colleagues, shows that short term use of valdecoxib and parecoxib in patients who had had coronary artery bypass grafting was associated with increased cardiovascular risk. □