EDITORIAL

Gaps in the Early Infant Diagnosis cascade in a high HIV prevalence setting

Rashida A. Ferrand^{1,2}<http://dx.doi.org/10.5588/pha.17.0049>

The scale-up of Prevention of Mother-To-Child Transmission (PMTCT) programmes has resulted in a substantial decline in the numbers of children born with the human immunodeficiency virus (HIV).¹ However, the PMTCT strategy is aimed not only at eliminating paediatric infections but also at improving the survival of infants born with HIV. A key component of PMTCT programming is therefore to test HIV-exposed infants through HIV-DNA polymerase chain reaction (PCR) testing at 6 weeks after birth (i.e., early infant diagnosis [EID]), and to start antiretroviral therapy (ART) promptly in those who test HIV-positive, an intervention which reduces mortality by 76%.²

The study by Phiri et al. in this issue of *Public Health Action* compared each step of the cascade of EID from registration of an infant into the programme to receipt of results by the child's care giver in two health facilities in Malawi.³ The first facility was a tertiary hospital performing DNA PCR testing on site, while the second was a district hospital that sent samples to a laboratory situated 360 km away.

This study demonstrates the gaps that exist across the EID cascade. First, only half of the samples were collected within 6–8 weeks of birth. Second, at both sites only about half of the care givers received their children's test results. At the tertiary hospital this was largely due to samples not being collected, while at the district hospital the tests were either not processed at the laboratory or the results were not received by health care providers. Third, a median delay of 8 weeks occurred from sample collection at the district hospital to receipt of the sample by the laboratory, largely due to erratic transport facilities, a finding that has been reported from other settings.^{4,5} Notably, at both health facilities, all HIV-positive infants initiated ART later than 12 weeks, which is when peak mortality occurs,⁶ and nearly a third of infants did not complete the EID cascade and were lost to follow-up by 24 months.

The impact of these delays is underscored by the fact that the mortality rate was significantly higher among infants who did not have a blood sample col-

lected or the blood sample was not collected in a timely fashion; furthermore, infants of care givers who received the results within one month of sample collection from their infants were at significantly lower risk of death than infants of care givers who did not receive their child's result.

The World Health Organization recommends ART initiation as early as possible in infants infected with HIV, to avert the substantial risk of mortality.⁷ HIV testing is the requisite first step to starting timely treatment and, as this study demonstrates, delays and gaps across the EID cascade have a direct impact on mortality. The survival gains from ART will only be realised if health systems are strengthened to address the barriers to timely HIV diagnosis of infants. Tracing procedures for HIV-exposed infants who do not present for follow-up, standards for timely delivery of samples and communication of results to health facilities and care givers and audit of these standards are critical to ensure quality of services. Finally, investment in point-of-care tests for HIV DNA testing is needed if universal coverage of EID is to be achieved.

References

- UNAIDS. 2014 progress report on the global plan towards the elimination of new HIV infections among children by 2015 and keeping their mothers alive. Geneva, Switzerland: UNAIDS, 2014.
- Violari A, Cotton M F, Gibb D M, et al. Early antiretroviral therapy and mortality among HIV-infected infants. *N Engl J Med* 2008; 359: 2233–2244.
- Phiri N A, Lee H-Y, Chilenga L, et al. Early infant diagnosis and outcomes in HIV-exposed infants at a central and a district hospital. *Public Health Action* 2017; 7: 6–12.
- Anoje C, Aiyenigba B, Suzuki C, et al. Reducing mother-to-child transmission of HIV: findings from an early infant diagnosis program in south-south region of Nigeria. *BMC Public Health* 2012; 12: 184.
- Mugambi M L, Deo S, Kekitiinwa A, Kiyaga C, Singer M E. Do diagnosis delays impact receipt of test results? Evidence from the HIV early infant diagnosis program in Uganda. *PLOS One* 2013; 8(11): e78891.
- Bourne D E, Thompson M, Brody L L, et al. Emergence of a peak in early infant mortality due to HIV/AIDS in South Africa. *AIDS* 2009; 23: 101–106.
- World Health Organization. Guideline on when to start antiretroviral therapy and on pre-exposure prophylaxis for HIV. Geneva, Switzerland: WHO, 2015.

AFFILIATIONS

- London School of Hygiene & Tropical Medicine, London, UK
- Biomedical Research and Training Institute, Harare, Zimbabwe

CORRESPONDENCE

Rashida A Ferrand
Clinical Research Department
London School of Hygiene & Tropical Medicine
Keppel Street
London WC1E 7HT, UK
e-mail: Rashida.Ferrand@lshtm.ac.uk

Conflicts of interest: none declared.

PHA 2017; 7(2): 78
© 2017 The Union