management strategy for assessing risk and preventing violence. This should include measures to calm down a potentially violent situation and intervene safely when violence occurs.

One member of staff should assume control of a potentially violent situation. They should encourage reasoning and explain to the service user and others nearby what they intend to do.

Only after these measures have failed to calm the service user should rapid tranquillisation, physical restraint and seclusion be considered. Intervention must be reasonable and proportionate. Failure to act in accordance with its recommendations would "not only be a failure to act in accordance with best practice, but in some circumstances may have legal consequences."

The guidelines were developed for the National Institute for Clinical Excellence (NICE) by the National Collaborating Centre for Nursing and Supportive Care, a professionally led group based at the Royal College of Nursing; they also included input from service user groups.

Susan Mayor *London*

The guidelines are at www.nice.org.uk/CG025NICEguideline

Conservatives promise to end waiting lists

Performance targets set by the government for NHS hospitals would be abolished under a new Conservative government, according to the party's health manifesto, released last week. Britain's 28 strategic health authorities would also go, while primary care trusts would have their number and functions reduced, in an effort "to restore the primacy of the doctor-patient relationship."

But the manifesto drew criticism from health unions and the Labour party for its proposal to hand NHS money to private patients as a reward for seeking care outside the NHS. Under the plan, patients who sought a private operation would be entitled to 50% of the NHS cost for that procedure as a contribution towards their bill.

Efforts to stamp out hospital acquired infections take centre stage in the party's programme, entitled "Action on Health." The party promises that matrons will have the power to overrule hospital managers on ward closures in cases of infection.

Reducing the bureaucracy of primary care trusts and encouraging private medicine would eliminate waiting lists under a Tory government, the manifesto claims.

Owen Dyer *London*

Doctor who spoke out on public health issue is sued

A doctor who claims he was doing his job according to the tenets of the Hippocratic oath when he spoke out about risks to health from the operations of a major logging company in Tasmania is being sued for causing alleged damage to the company's business activities.

Dr Frank Nicklason, a staff specialist physician at the Royal Hobart Hospital, is one of 20 defendants, including prominent environmentalists and another doctor, named in the writ by Gunns Ltd, which is seeking almost \$A6.3m (£2.6m; \$5m; €3.8m) in damages.

Dr Nicklason said the case may stop doctors raising legitimate health concerns because of fear of being involved in prolonged and expensive legal action.

"It has succeeded in shutting other people up in Tasmania, mainly small local environmental and community groups," he said of the writ, which was served in December.

Dr Nicklason faces years of legal action and a damages claim of \$A250 000 for calling for an independent risk assessment of large piles of woodchips in the port of Burnie.

He said the Hippocratic oath required him to prevent illness not just in patients but in society as well, and he believed, after talking with experts, that the stockpiles of shredded wood on the wharf posed potential health risks to Burnie's citizens.

Christopher Zinn *Sydney*

NICE issues guidance for diagnosis and treatment of lung cancer

Susan Mayor *London*

Guidelines for achieving earlier diagnosis, greater use of evidence based treatments, and better coordination of care for patients with lung cancer have been published for England and Wales this week. The recommendations, from the National Institute for Clinical Excellence (NICE), are aimed at improving the poor survival rates from lung cancer.

The guidelines recommend urgent referral for chest x ray examination for anyone with haemoptysis or unexplained or persistent symptoms that might indicate lung cancer, including cough, chest pain, dyspnoea, or weight loss. Patients whose chest x ray examination or computed tomogram suggests lung cancer should be referred urgently to a chest physician working as a member of a lung cancer multidisciplinary team.

Every cancer network must have rapid access to 18F-deoxyglucose positron emission tomography (FDG-PET) to improve staging of lung cancer, say the guidelines. They also recommend that all cancer units should have one or more trained lung cancer nurse specialists to see patients before and after diagnosis to provide continuing support and to facilitate communication between all professionals involved in an individual patient's care.

Currently just 5.5% of people diagnosed with lung cancer in England and 6% of those in Wales live for more than five years. This is about 5% lower than the European average and 7-10% lower

than the rate for the United States.

The guidance is based on a review of evidence by NICE and the National Collaborating Centre for Acute Care, a group set up by NICE to include representatives from the medical royal colleges, professional bodies, and patient organisations. For the first time, NICE worked in collaboration with the Scottish Intercollegiate Guidelines Network (SIGN).

Dr Jesme Baird, director of patient care with the Roy Castle Lung Cancer Foundation and chairwoman of the guideline development group, commented: "Despite being the commonest cancer, lung cancer remains a disease associated with much negativity. The recommendations made in this guideline highlight many positive developments."

The guidelines recommend greater use of modern, evidence based treatments. Patients with stages I or II non-small cell lung cancer that is medically inoperable should be treated using continuous hyperfractionated accelerated radiotherapy (CHART)—an intensive form of radiotherapy given three times a day over a relatively short period of just over two weeks—rather than standard radiotherapy. The guidelines also recommend offering chemotherapy to patients with stages III and IV non-small cell lung cancer who are generally well—to improve survival, disease control, and quality of life. □

The NICE guideline is available from <http://www.nice.org.uk/CG024NICEguideline>

Chest x ray film of a 70 year old woman showing lung cancer