Published in final edited form as:

Dev Psychobiol. 2017 July ; 59(5): 658–667. doi:10.1002/dev.21523.

The Feasibility of Salivary Sample Collection in an International Pediatric Cohort: the TEDDY Study

Roth Roswith^{1,2}, Judith Baxter³, Kendra Vehik⁴, Diane Hopkins⁵, Michael Killian⁶, Patricia Gesualdo³, Jessica Melin⁷, Barbara Simell⁸, Elisabeth Strauss¹, Åke Lernmark⁷, and Suzanne Bennett Johnson⁹ TEDDY Study Group

¹Institute of Diabetes Research, Helmholtz Zentrum München, and Klinikum rechts der Isar, Technische Universität München, and Forschergruppe Diabetes e.V, Neuherberg, Germany

²University of Graz, Graz, Austria ³Barbara Davis Center for Childhood Diabetes, University of Colorado Anschutz Medical Campus, Aurora CO, U.S.A ⁴Pediatric Epidemiology Center, University of South Florida, Tampa FL, U.S.A ⁵Georgia Health Sciences University Augusta GA, U.S.A ⁶Pacific Northwest Diabetes Research Institute, Seattle WA, U.S.A ⁷Department of Clinical Sciences, Lund University, CRC, Malmö, Sweden ⁸Department of Pediatrics, Turku University Central Hospital, Turku, Finland ⁹College of Medicine, Florida State University, Tallahassee FL, U.S.A

Abstract

Saliva offers a relatively noninvasive method for measuring analytes such as cortisol, holding particular promise for use in pediatric populations on a large scale if a rigorous collection protocol is feasible in diverse settings. The Environmental Determinants of Diabetes in the Young study protocol, conducted in centers in the US, Sweden, Finland and Germany, used salivary collection to assess cortisol level as a physiologic marker of stress. Saliva was collected using Sorbettes from subjects at 3.5, 4.5, and 5.5 years of age. Parents collected a morning sample, and staff collected pre- and post-blood draw samples. Feasibility was assessed based on protocol completion, adherence with instructions, factors affecting adherence and sufficiency of saliva sample for cortisol determination. Collection of saliva samples in a diverse pediatric population is feasible. Establishing non-invasive and acceptable methods for collecting physiological parameters of stress will allow better exploration of determinants of health in this important population.

Keywords

salivary collection; adherence; methodological feasibility; multi-center international; pediatric cohort study; salivary cortisol

INTRODUCTION

The last decade has seen a marked increase in the use of saliva for determination of many analytes including corticosteroids, alpha-amylase, immunoglobulin, as well as DNA (Chiappin, Antonelli, Gatti, & De Palo, 2007; Nater, Rohleder, Schlotz, Ehlert, & Kirschbaum, 2007; Rohleder & Nater, 2009). Interest and further understanding of the role stress hormones play in disease risk has prompted an increase in the number of studies using salivary cortisol as a biomarker of stress in many populations (Hellhammer, Wust, & Kudielka, 2009). Further, saliva sampling is an inexpensive and noninvasive method offering advantages over serum testing, particularly in pediatric populations. However, there are no reports in the literature that evaluate the feasibility of saliva collection generally, or salivary cortisol collection specifically, in large-scale pediatric, international cohort studies. The Environmental Determinants of Diabetes in the Young (TEDDY) Study has completed salivary collection on about 4000 children ages 3.5–5.5 years old, across four countries in six study centers and provides a unique opportunity to assess multiple aspects of this method.

Collecting saliva is convenient for self-sampling in naturalistic environments and has been used with young children (Gunnar & White, 2001; McCarthy et al., 2009; Neu, Goldstein, Gao, & Laudenslager, 2007). Research has used saliva samples to assess general stress levels using the indicator of Cortisol Awakening Response (CAR) (Bäumler, Kirschbaum, Kliegel, Alexander, & Stalder, 2013; Bright, Granger, & Frick, 2012; Bruce, Fisher, Pears, & Levine, 2009; Freitag et al., 2009; Gribbin, Watamura, Cairns, Harsh, & Lebourgeois, 2012; Saridjan et al., 2010; Stalder et al., 2013; Tegethoff, Knierzinger, Meyer, & Meinlschmidt, 2013). Salivary cortisol can also be used to assess the child's response to a particular stressor (e.g., venipuncture, injection, inoculation) (Davis & Granger, 2009; Felt et al., 2000; McCarthy et al., 2009; McCarthy et al., 2011) and other stress evoking procedures (e.g. Trier Social Stress Test for Children) (Gunnar, Talge, & Herrera, 2009; Kirschbaum, Pirke, & Hellhammer, 1993). However, all of these studies employed relatively small samples of children.

Adherence with sampling procedures, especially the timing of sampling, is crucial for accurate assessment of cortisol levels (Gunnar & Talge, 2007; Hall et al., 2011; Hanrahan, McCarthy, Kleiber, Lutgendorf, & Tsalikian, 2006; Jessop & Turner-Cobb, 2008; Kudielka, Hawkley, Adam, & Cacioppo, 2007; Schwartz, Granger, Susman, Gunnar, & Laird, 1998; Smyth, Clow, Thorn, Hucklebridge & Evans, 2013; Stalder, Kirschbaum, Kudielka et al., 2016). Instructions such as preventing the child from eating or drinking something except water, crying or participating in vigorous activities for 30 minutes prior to collection, as well as managing timing of the child's medication may pose additional challenges (Hanrahan et al., 2006).

The constraints of the sampling protocol in conjunction with child and parent factors (e.g., motivation, ability, work, and family responsibilities) significantly raise the potential for protocol non-adherence. In a small study, Smith et al. (Smith & Dougherty, 2014) found substantial discrepancy between parent self-report and electronically-monitored report of CAR sample collections, with parents reporting higher collection rates (78% – 85% versus

56%–77% of requested samples). Further, to our knowledge there are no reports of staff adherence with salivary sampling protocols. These strict protocol requirements of saliva collection for cortisol measurement combined with the diverse cohort in the TEDDY study provided an ideal opportunity to conduct a rigorous examination of feasibility of a saliva collection protocol among parents and study staff.

Another important consideration is whether the study's saliva collection protocol results in sufficient sample volume for laboratory assay requirements. Very few studies report these data. McCarthy et al. (2009) reported 19 out of 1355 cortisol samples were inadequate, 11 due to questionable sampling.

The aim of this study was to conduct a multifaceted assessment of the feasibility of salivary collection for cortisol determination in a large longitudinal multi-national cohort study of preschoolers. Feasibility was assessed by an examination of protocol acceptance and completion, adherence to instructions by both parents and study staff, factors affecting adherence, and sufficiency of sample for cortisol determination.

METHOD

Study Design and Population

TEDDY is a prospective cohort study funded by the National Institutes of Health (NIH) with the primary goal to identify environmental triggers of type 1 diabetes (T1D) in children genetically at-risk for the disease. It includes six clinical research centers, three in the United States (US) – Colorado, Georgia/Florida, Washington and three in Europe (EU) – Finland, Germany, and Sweden. Detailed study design and methods have been previously published (The TEDDY Study Group, 2007, 2008). The study was approved by local Institutional Review Boards and is monitored by an External Evaluation Committee formed by NIH. Participants in this study are children determined at birth to have high risk genetic markers for T1D whose parents agreed to participate in a 15-year observational study starting at the age of 3–4.5 months. TEDDY's protocol includes 2–4 visits per year with extensive data collection through interviews and biologic sample collection. The salivary collection protocol was approved in 2008 with the European centers being the first to start followed two-three years later by the US centers. The focus of the current report is on the feasibility of collecting salivary samples that would be used to assess cortisol in the morning after awakening and before and after receiving a venous blood draw at TEDDY visits when the child was 42-, 54-, and 66-months of age.

As of December 2014, 12,690 TEDDY visits had been completed where the saliva cortisol protocol was in place (8977 EU and 3713 US TEDDY visits). The experience from these 12,690 study visit saliva collections is the basis for the feasibility assessment.

Method and Procedures of Saliva Collection

Each annual collection consisted of three salivary samples: 1) a morning sample about 30 minutes (time window 25–35 minutes) after awakening taken at home by parents; 2) a sample collected before a blood draw at a TEDDY study visit by study staff (<10 minutes);

and 3) a sample collected about 20–30 minutes (time window 15–35 minutes) after a blood draw by study staff.

All study staff received in-person training in the collection technique. At the clinic visit prior to the first home collection, research staff demonstrated the saliva collection technique to the parent. Some parents were given the written instructions and the saliva kit to take home while other parents were mailed the kit and instructions prior to the next study visit. The instructions ask parents to collect the saliva the morning of their next study visit, after the child has been awake for about 30 minutes and before eating, drinking or brushing the child's teeth.

Each Salimetrics salivary collection kit consisted of three Sorbettes (cotton pads on a stick developed for collection in infants) and a storage tube (Donzella, Talge, Smith, & Gunnar, 2008; Putnam et al., 2012). The Salimetrics kit instructions indicate that up to three Sorbettes should be used in immediate succession to assure sufficient sample volume. A single Sorbette should be placed under the child's tongue until the Sorbette tip is saturated with saliva. Upon removal, the second Sorbette is placed under the child's tongue until saturation, removed and replaced by a third Sorbette. All three Sorbettes are placed in one storage tube. Upon delivery to the study clinic for processing, the tube is placed in a centrifuge and spun at 1500g for 15 minutes. After the spinning is complete, the Sorbette are removed with tweezers and discarded. The saliva sample is then stored in a -70°C freezer.

On the morning of the in-home collection, the parent was instructed to document the time the child woke up and the time the first saliva sample was collected. In-home collection was scheduled on the same day as a TEDDY study visit. If a parent failed to complete the morning in-home collection on the same day as the TEDDY study visit, collection within one week of the study visit was accepted. Home samples were returned to the child's study site either in person or by standard mail. Cortisol concentrations remain stable for days if samples are mailed before freezing (Clements & Parker, 1998).

At the in-clinic saliva collection, the staff documented the time the child last ate, whether caffeine was consumed since waking up and whether the child had participated in any vigorous activity or had been upset or crying in the prior 30 minutes. If the child had some food or drink – except water – in the prior 30 minutes, the collection was either postponed for 30 minutes or the protocol violation was noted. If the child was on steroid medication the salivary cortisol collection was not performed. The protocol instructions included an initial saliva collection < 10 minutes before the child's blood draw and a second saliva collection 20–30 minutes after the blood draw. The time of the pre- and post-blood draw saliva collection, the time of the blood draw attempt, and whether it was a venous or capillary blood draw were documented.

Center Differences in the Implementation of the Protocol

The multi-center, cross-national, longitudinal structure of TEDDY requires a consistent investment in developing standardized protocols that can be implemented successfully in diverse settings. We attempted to minimize the differences in implementing the salivary

cortisol protocol between the clinical sites by use of extensive training and written documentation in the manual of operations.

Although a collection demonstration was given to parents at all TEDDY study sites, there were important center differences in the timing of when the saliva kit was given to the parent. The Swedish and Finnish centers distributed the saliva kits and instructions to the parents at the visit three to six months before the collection visit. The other 4 clinical centers mailed the saliva kit and instructions two to eight weeks before the sample collection visit.

Centers also differed in their inclusion or exclusion of participants who were on a long-distance protocol (LDP), meaning that they do not come into a study center but do the study protocol (including blood draws) at a remote location. This protocol option was developed in part for US TEDDY centers where there is greater population mobility resulting in a portion of the cohort that has moved too far away to continue to be seen at the study site, but who are committed to continuing with the basic study protocol. Due to the increased burden on the family, the US sites made the decision to exclude LDP participants from the saliva protocol. The Swedish and Finnish study sites do not routinely have subjects on a LDP.

The German site is unique in that 80% of the cohort have their study data collected at their local pediatrician's office and not at a study clinic. Therefore, the staff developed remote sampling collection procedures where children have all of their saliva collections (morning-, pre- and post-blood draw) completed by parents. This was accomplished by developing instructions that were more detailed and included pictures of the collection procedure since in-person demonstration was not possible. These instructions were reinforced with labels for each collection tube.

Saliva Cortisol Analysis

Ability to consistently obtain a sufficient sample for analysis is an important feasibility criteria for any assay. To assess the sufficiency of the saliva cortisol sample collected, a subset of 5495 samples collected was sent to the laboratory of the Department of Clinical Chemistry, University Hospital, Linköping, Sweden. The assay used for determination of cortisol in saliva is a commercial enzyme-linked immunoassay from Salimetrics LLC, Carlsbad, CA applied on a Tecan Freedom EvoLyzer (Tecan, Männedorf, Switzerland).

RESULTS

Protocol Acceptance

As of December 31, 2014, TEDDY participants who attended the 42- (N=4307), 54- (N=4545) or 66- (N=3838) month visit were eligible to participate in the salivary cortisol protocol, for a total of 3713 US and 8977 EU TEDDY visits. Table 1 describes the participation and refusal rates for each of these visits for all TEDDY participants and for EU and US centers separately. In the 12,690 TEDDY visits, 814 (6.4%) of potential participants were not offered the protocol by study staff for reasons such as clinic preparedness to start the protocol, difficult or missed visits where data collection was completed over the phone rather than in person, saliva collection packets not received, or clinic discretion that this visit/subject wasn't a good candidate for this protocol. More US (14.9%) compared to EU

(2.9%) participants across all visit intervals were not offered the protocol ($\chi^2=624.6$, $p<0.0001$). The proportion not offered the protocol in the US ranged from 5.9–30.7% and was more common in the Georgia/Florida (30.7%) and Washington (19.3%) centers where a greater percentage of the cohort has challenges completing the core TEDDY visit protocol and staying in the study. For the EU centers the range was 2.7–9.1% (Center specific data not shown).

Study-wide the rate of completing at least one of three samples on any given saliva collection day was 95.9% among those participants offered the protocol. All centers had rates above 90%, ranging from 91.9–98.6%. This included Germany where a majority of participants were on LDP. There was no meaningful difference in the completion rates for US (98.1%) compared to the EU centers (95.1%). Very few participants offered the protocol refused to do it.

Protocol Adherence

The number and percentage of visits that were missing elements of the saliva collection protocol among the 11,390 visits where at least one cortisol sample was obtained is shown in Table 2. Missing samples at each time point (morning, pre- and post-blood draw) were generally uncommon, ranging from 0.8–4.9% across all sites. The morning sample, which relied solely on the parent for collection, was missing more often than saliva samples collected at the study visit ($\chi^2=353.1$, $p<0.0001$). There was considerable site variability in success getting the morning sample, ranging from 1.3–19.8% missing (data not shown). Overall the US centers had greater difficulty obtaining the morning cortisol sample compared to the EU centers (11.2% and 2.5% missing, respectively; $\chi^2=363.1$, $p<0.0001$). When samples were collected in clinic there were very few missed collections and no differences between centers.

Some collected samples were excluded for the following reasons: brushed teeth without rinsing, ate or drank less than 30 minutes before cortisol collection or were on oral steroids during last 30 days. Exclusion of a sample for any of these reasons was uncommon, suggesting that parents were good at following the instructions. There was limited center variability in proportion of samples excluded (1.0–5.2%), though on average this was not different for US (3.7%) compared to EU (2.5%) sites.

Staff adherence with timing of saliva collection was best for both EU and US centers for the sample collected after blood draw, with 95% and 95.6%, respectively, being collected within the 15–35-minute window (data not shown). The pre-blood draw sample had worse adherence, with center differences emerging. Among EU centers 78.7% of samples were collected in the protocol window of <10 min before blood draw, compared to 60.4% ($\chi^2=393.6$, $p<0.0001$) in the US centers. Parent adherence with the timing of the morning saliva sample collection (25–35 minutes after awakening) was 65.8%, with a persistent difference between the EU (70.3%) and US (54.0 %) ($\chi^2=264.3$, $p<0.0001$) centers.

If saliva was collected, a sufficient sample was obtained almost all of the time at all sites, with across site sample sufficiency ranging from 97–100%. The one exception to this was Germany where sample sufficiency was lower (79.5%) but improved over time to 90%.

DISCUSSION

Salivary cortisol collection has emerged in pediatric research as a biomarker of stress in both US (Bright et al., 2012; Bruce et al., 2009; Felt et al., 2000; McCarthy et al., 2009; McCarthy et al., 2011) and European pediatric studies (Bäumler et al., 2013; Freitag et al., 2009; Saridjan et al., 2010; Stalder et al., 2013; Tegethoff et al., 2013) with relatively small sample sizes. However, these reports provide very limited information about collection adherence and quality, information important for any effort to collect saliva samples on a large scale. The TEDDY experience provides an opportunity to examine several aspects of the feasibility of salivary collection in a large multi-center, cross-national longitudinal study of children ages 3.5–5.5 years old that required sample collections from both parents and staff. Following a well developed standardized protocol, the TEDDY experience demonstrates that a salivary cortisol protocol could be implemented successfully in diverse settings. This analysis has also suggested areas where challenges to saliva collection completion and protocol adherence exist and that could be attended to in the protocol design.

Different aspects of adherence were investigated which are associated with saliva collection feasibility. Among all TEDDY children offered the saliva sampling protocol, 95.9% provided one or more saliva samples (US: 98.1%, EU: 95.1%). Among the US and EU parents collecting the morning saliva sample, 95.1% were able to obtain a sample, with a notable range between centers where US parents had lower adherence (88.8%) compared to EU parents (97.5%). Among the staff, 99.2% of samples were collected. These rates are comparable or higher than those reported by studies where adherence was objectively assessed (Broderick, Arnold, Kudielka, & Kirschbaum, 2004; Smith & Dougherty, 2014).

The refusal rate was very low at 4.1% (US: 1.9%, EU: 4.9%). The lower refusal rate as well as the high adherence rate among those offered the protocol in US stands in stark contrast to the proportion of subjects who were selectively not offered the protocol (US 14.9%) because they were experiencing challenges completing the standard TEDDY protocol compared to those not offered it in the EU centers (2.9%).

Center differences in offering the protocol reflected EU/US differences in retention concerns and participant burden, where retention challenges have been more apparent for the US centers (Baxter et al., 2012; Johnson et al., 2011, 2014, 2016; Lernmark et al., 2016). Retention of study participants in a longitudinal cohort is critical to the overall study design and can be particularly challenging for a pediatric cohort enrolled at the age of three months and followed for fifteen years.

Overall completion and adherence to the sample timing collection protocol was lower for the morning sample compared to the pre- and post-blood draw samples. Smith & Dougherty (2014) observed 78–85% self-reported adherence in parents who collected the CAR saliva samples. In the present study, with a more diverse population and longitudinal sample collection, higher completion rates were observed (EU: 97.5%; US: 88.8%). The pre-post blood draw collection rates were even higher, with only 0.8% before and 1.0% after the blood draw samples not collected. These completion rates are especially striking because although they were collected primarily by staff in the clinic setting, they also included pre-

and post-blood draw saliva collections by parents on the German LDP. In addition, very few saliva samples had to be excluded (in total 2.8%, EU: 2.5%, US 3.7%) because of failure to follow the protocol properly.

Saliva can be used to measure many different analytes which may have specific requirements such as time of collection and restrictions on food, drink, and medications (Chiappin et al., 2007; Hellhammer et al., 2009; Nater et al., 2007; Rohleder & Nater, 2009). TEDDY's salivary cortisol collection is useful to assess feasibility as it was done in four countries, by different collectors, in different settings, with detailed instructions regarding timing and restrictions. Though a majority of all samples were collected according to protocol within the timing windows specified, this element had the most variability in compliance across sample collections and centers. The morning collection had the lowest percentage of saliva samples collected in the protocol time window and showed a wide degree of variability across centers. Strict adherence to the instructions may not be as easily accomplished by parents in the child's home environment with its increased demands (child's sleep patterns, morning family schedules, etc.) compared to a clinic setting. Since the timing of the morning saliva collection was provided by parent report, parents may have overestimated adherence with the instructions. Future studies should examine common barriers to in-home saliva collection protocol adherence, so that strategies to address these barriers could be designed. The use of objective devices to measure the child wake up and saliva collection times could be very useful (Stalder et al., 2016).

Failure to follow the protocol in terms of the timing of the saliva collection was also a problem among TEDDY staff. Of the two saliva samples collected by staff at the TEDDY visit, the pre-blood draw collection time adherence was lower than the post-blood draw time adherence. The logistical variables associated with any TEDDY visit are more complex at the beginning of a visit compared to its end, when the post-blood draw saliva sample is done. It is not always easy to predict when the blood draw will occur relative to other data collection elements, with staff making the best effort possible to determine when the pre-draw saliva collection was to be administered. The differences in adherence for this element between the EU and US centers may reflect differences in how the clinics are organized generally around timing and conduct of the blood draws. It is possible that a stricter visit schedule for those visits that included a saliva sample collection, using an objective electronic device to record sample collection time (Stalder et al., 2016) and improved organization might have increased the time adherence for the pre-blood draw saliva collection.

Among the 5495 saliva samples sent to the lab for cortisol assay, only 1.6% of samples were excluded due to an insufficient amount of saliva. Bright et al (Bright et al., 2012) reported 34 invalid samples of 256 (14%) mostly due to insufficient sample sizes. The excellent sufficiency results in the TEDDY study also confirmed the suitability of the Sorbette as collection device for toddlers (Donzella et al., 2008; Putnam et al., 2012), although Salimetrics now offers another saliva collection device for children and infants (<https://www.salimetrics.com/collection-system/childrens-swab>).

The study has several limitations. Smyth et al. (Smyth, Clow, Thorn, Huklebridge & Evans, 2013) reported that delays of 5–15 minutes between awakening and the start of saliva sampling matter in assessment of the CAR. Since both the awakening time of the children the saliva collection time was determined by parent report, we cannot be sure of the accuracy of these reports. Although evidence suggests that parents are reasonably reliable reporters of children's awaking times (Tikotsky & Sadeh, 2001), the use of actigraphy would provide a more objective assessment of children's wake times and cap tracks could be used to record the saliva collection (Smith & Dougherty, 2013). Recently published expert consensus guidelines for the collection of morning cortisol for the assessment of the CAR include both the use of various electronic devices and careful attention to participant instructions (Stalder et al., 2016). Adam & Kumari (2009) described benefits and challenges of assessing saliva cortisol in large-scale epidemiological research in comparison to convenience samples. They suggest using electronic monitoring devices to measure timing adherence for at least a subsample of participants.

We acknowledge that the TEDDY population is a highly-motivated sample of educated parents who have volunteered to participate in a longitudinal study with a demanding protocol. Such parents might be more motivated than most to be highly adherent with the study saliva collection protocol. Or, they might be more motivated to report greater adherence. In either case, our study findings may not replicate in other populations.

The same limitations hold for the time of saliva collection of the pre- and post-blood draw reported by clinical staff. Sampling time details in the clinic were noted by clinical staff without objective device. Though research staff may feel pressure to report greater adherence, Stalder et al. (2016) conclude that clinical staff are reliable in recording. In this study, the highly-trained TEDDY clinical staff reported poorer adherence to some aspects of the pre-post blood draw saliva collection and there were differences in adherence between centers, suggesting biased or better adherence reporting is not likely a limitation in this case.

Nevertheless, we believe the findings of this study are valuable and provide important information about using salivary collection in an international and cross-cultural context of a longitudinal study. This effort demonstrated that saliva collection by both parent and staff according to a standardized protocol was possible and yielded sufficient samples for laboratory analysis. Also, demonstrated here was the importance of creating a well-documented, standardized protocol that was attentive to center specific conditions and was flexible, within reason, for parents and staff to complete. Establishing non-invasive and acceptable methods for collecting physiological parameters of stress in children will allow better exploration of determinants of health in this important population.

Acknowledgments

TEDDY Study Acknowledgements

The TEDDY Study Group (See appendix)

Funded by U01 DK63829, U01 DK63861, U01 DK63821, U01 DK63865, U01 DK63863, U01 DK63836, U01 DK63790, UC4 DK63829, UC4 DK63861, UC4 DK63821, UC4 DK63865, UC4 DK63863, UC4 DK63836, UC4 DK95300, and UC4 DK100238, and Contract No. HHSN267200700014C from the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK), National Institute of Allergy and Infectious Diseases (NIAID),

National Institute of Child Health and Human Development (NICHD), National Institute of Environmental Health Sciences (NIEHS), Juvenile Diabetes Research Foundation (JDRF), and Centers for Disease Control and Prevention (CDC). This work supported in part by the NIH/NCATS Clinical and Translational Science Awards to the University of Florida (UL1 TR000064) and the University of Colorado (UL1 TR001082).

Appendix

The Teddy Study Group

Colorado Clinical Center: Marian Rewers, M.D., Ph.D., PI^{1,4,5,6,10,11}, Kimberly Bautista¹², Judith Baxter^{9,10,12,15}, Ruth Bedoy², Daniel Felipe-Morales, Brigitte Frohnert, M.D., Patricia Gesualdo^{2,6,12,14,15}, Michelle Hoffman^{12,13,14}, Rachel Karban¹², Edwin Liu, M.D.¹³, Jill Norris, Ph.D.^{2,3,12}, Adela Samper-Imaz, Andrea Steck, M.D.^{3,14}, Kathleen Waugh^{6,7,12,15}, Hali Wright¹². University of Colorado, Anschutz Medical Campus, Barbara Davis Center for Childhood Diabetes.

Georgia/Florida Clinical Center: Jin-Xiong She, Ph.D., PI^{1,3,4,11}, Desmond Schatz, M.D.^{*4,5,7,8}, Diane Hopkins¹², Leigh Steed^{12,13,14,15}, Jamie Thomas^{*6,12}, Katherine Silvis², Michael Haller, M.D.^{*14}, Meena Shankar^{*2}, Eleni Sheehan^{*}, Melissa Gardiner, Richard McIndoe, Ph.D., Ashok Sharma, Joshua Williams, Gabriela Foghis, Stephen W. Anderson, M.D.[^], Richard Robinson. Center for Biotechnology and Genomic Medicine, Georgia Regents University. ^{*}University of Florida, [^]Pediatric Endocrine Associates, Atlanta.

Germany Clinical Center: Anette G. Ziegler, M.D., PI^{1,3,4,11}, Andreas Beyerlein Ph.D.², Ezio Bonifacio Ph.D.^{*5}, Michael Hummel, M.D.¹³, Sandra Hummel, Ph.D.², Kristina Foterek^{¥2}, Mathilde Kersting, Ph.D.^{¥2}, Annette Knopff⁷, Sibylle Koletzko, M.D.^{¶13}, Claudia Peplow¹², Roswith Roth, Ph.D.⁹, Joanna Stock^{9,12}, Elisabeth Strauss¹², Katharina Warncke, M.D.¹⁴, Christiane Winkler, Ph.D.^{2,12,15}. Forschergruppe Diabetes e.V. and Institute of Diabetes Research, Helmholtz Zentrum München, and Klinikum rechts der Isar, Technische Universität München. ^{*}Center for Regenerative Therapies, TU Dresden, [¶]Dr. von Hauner Children's Hospital, Department of Gastroenterology, Ludwig Maximilians University Munich, [¥]Research Institute for Child Nutrition, Dortmund.

Finland Clinical Center: Jorma Toppari, M.D., Ph.D., PI^{¥1,4,11,14}, Olli G. Simell, M.D., Ph.D.^{¥1,4,11,13}, Annika Adamsson, Ph.D.^{^12}, Heikki Hyöty, M.D., Ph.D.^{*±6}, Jorma Ilonen, M.D., Ph.D.^{¥¶3}, Sanna Jokipuu[^], Tiina Kallio[^], Miia Kähönen^{μ¶}, Mikael Knip, M.D., Ph.D.^{*±5}, Annika Koivu[¥], Mirva Koreasalo^{*±§2}, Kalle Kurppa, M.D., Ph.D.^{*±13}, Maria Lönnrot, M.D., Ph.D.^{*±6}, Elina Mäntymäki[¥], Katja Multasuo^{μ¶}, Juha Mykkanen, Ph.D.^{¥3}, Tiina Niininen^{±*12}, Mia Nyblom^{*±}, Petra Rajala[^], Jenna Rautanen^{±§}, Anne Riikonen^{*±}, Minna Romo[¥], Satu Simell, M.D., Ph.D.^{^±13}, Tuula Simell, Ph.D., Ville Simell^{¥13}, Maija Sjöberg^{¥12,14}, Aino Stenius^{μ¶12}, Maria Särmä[^], Sini Vainionpää[^], Eeva Varjonen^{¥12}, Riitta Veijola, M.D., Ph.D.^{μ¶14}, Suvi M. Virtanen, M.D., Ph.D.^{*±§2}, Mari Vähä-Mäkilä[^], Mari Åkerlund^{*±§}. [¥]University of Turku, ^{*}University of Tampere, ^μUniversity of Oulu, [^]Turku University Hospital, Hospital District of Southwest Finland, [±]Tampere University Hospital, [¶]Oulu University Hospital, [§]National Institute for Health and Welfare, Finland, [¶]University of Kuopio.

Sweden Clinical Center: Åke Lernmark, Ph.D., PI^{1,3,4,5,6,8,10,11,15}, Daniel Agardh, M.D., Ph.D.¹³, Carin Andrén Aronsson^{2,13}, Maria Ask, Jenny Bremer, Ulla-Marie Carlsson, Corrado Cilio, Ph.D., M.D.⁵, Emelie Ericson-Hallström, Lina Fransson, Thomas Gard, Joanna Gerardsson, Rasmus Bennet, Monica Hansen, Gertie Hansson¹², Susanne Hyberg, Fredrik Johansen, Berglind Jonasdottir M.D., Linda Jonsson, Helena Elding Larsson M.D., Ph.D.^{6,14}, Sigrid Lenrick Forss, Maria Månsson-Martinez, Maria Markan, Jessica Melin¹², Zeliha Mestan, Kobra Rahmati, Anita Ramelius, Anna Rosenquist, Falastin Salami, Sara Sibthorpe, Birgitta Sjöberg, Ulrica Swartling, Ph.D.^{9,12}, Evelyn Tekum Amboh, Erika Trulsson, Carina Törn, Ph.D.^{3,15}, Anne Wallin, Åsa Wimar^{12,14}, Sofie Åberg, Lund University.

Washington Clinical Center: William A. Hagopian, M.D., Ph.D., PI^{1,3,4,5,6,7,11,13,14}, Xiang Yan, M.D., Michael Killian^{6,7,12,13}, Claire Cowen Crouch^{12,14,15}, Jennifer Skidmore², Stephen Ayres, Kayleen Dunson, Diana Heaney, Rachel Hervey, Corbin Johnson, Rachel Lyons, Arlene Meyer, Denise Mulenga, Emma Schulte, Elizabeth Scott, Joshua Stabbert, John Willis. Pacific Northwest Diabetes Research Institute.

Pennsylvania Satellite Center: Dorothy Becker, M.D., Margaret Franciscus, MaryEllen Dalmagro-Elias Smith², Ashi Daftary, M.D., Mary Beth Klein, Chrystal Yates. Children's Hospital of Pittsburgh of UPMC.

Data Coordinating Center: Jeffrey P. Krischer, Ph.D., PI^{1,4,5,10,11}, Michael Abbondandolo, Sarah Austin-Gonzalez, Sandra Baethke, Rasheedah Brown^{12,15}, Brant Burkhardt, Ph.D.^{5,6}, Martha Butterworth², Joanna Clasen, David Cuthbertson, Christopher Eberhard, Steven Fiske⁹, Dena Garcia, Jennifer Garmeson, Veena Gowda, David Hadley, Ph.D., Kathleen Heyman, Hye-Seung Lee, Ph.D.^{1,2,13,15}, Shu Liu, Xiang Liu, Ph.D.^{2,3,9,14}, Kristian Lynch, Ph.D.^{5,6,9,15}, Jamie Malloy, Cristina McCarthy^{12,15}, Wendy McLeod, Steven Meulemans, Chris Shaffer, Laura Smith, Ph.D.^{9,12}, Susan Smith^{12,15}, Roy Tamura, Ph.D.^{1,2,13}, Ulla Uusitalo, Ph.D.^{2,15}, Kendra Vehik, Ph.D.^{4,5,6,14,15}, Ponni Vijayakandipan, Keith Wood, Jimin Yang, Ph.D., R.D.^{2,15}. University of South Florida.

Project scientist: Beena Akolkar, Ph.D.^{1,3,4,5,6,7,10,11}. National Institutes of Diabetes and Digestive and Kidney Diseases.

Other contributors: Kasia Bourcier, Ph.D.⁵, National Institutes of Allergy and Infectious Diseases. Thomas Brieze, Ph.D.^{6,15}, Columbia University. Suzanne Bennett Johnson, Ph.D.^{9,12}, Florida State University. Steve Oberste, Ph.D.⁶, Centers for Disease Control and Prevention. Eric Triplett, Ph.D.⁶, University of Florida.

Autoantibody Reference Laboratories: Liping Yu, M.D.^{^5}, Dongmei Miao, M.D.[^], Polly Bingley, M.D., FRCP^{*5}, Alistair Williams*, Kyla Chandler*, Saba Rokni*, Claire Caygill*, Nicholas Lovis, Claire Williams, Rebecca Wyatt*. ^Barbara Davis Center for Childhood Diabetes, University of Colorado Denver, *School of Clinical Sciences, University of Bristol UK.

Cortisol Laboratory: Elisabeth Aardal Eriksson, M.D., Ph.D., Ing-Marie Lundgren, Ewa Lönn Karlsson, Dzeneta Nezirevic Dernroth, Ph.D. Department of Clinical Chemistry, Linköping University Hospital, Linköping, Sweden.

Dietary Biomarkers Laboratory: Iris Erlund, Ph.D.², Irma Salminen, Jouko Sundvall, Jaana Leiviskä, Nina Kangas. National Institute for Health and Welfare, Helsinki, Finland.

HbA1c Laboratory: Randie R. Little, Ph.D., Alethea L. Tennill. Diabetes Diagnostic Laboratory, Dept. of Pathology, University of Missouri School of Medicine.

HLA Reference Laboratory: Henry Erlich, Ph.D.³, Steven J. Mack, Ph.D., Anna Lisa Fear. Center for Genetics, Children's Hospital Oakland Research Institute.

Metabolomics Laboratory: Oliver Fiehn, Ph.D., Bill Wikoff, Ph.D., Brian Defelice, Dmitry Grapov, Ph.D., Tobias Kind, Ph.D., Mine Palazoglu, Luis Valdiviez, Benjamin Wancewicz, Gert Wohlgemuth, Joyce Wong. UC Davis Metabolomics Center.

Microbiome and Viral Metagenomics Laboratory: Joseph F. Petrosino, Ph.D.⁶. Alkek Center for Metagenomics and Microbiome Research, Department of Molecular Virology and Microbiology, Baylor College of Medicine.

OGTT Laboratory: Santica M. Marcovina, Ph.D., Sc.D., Vinod P. Gaur, Ph.D., Northwest Lipid Metabolism and Diabetes Research Laboratories, University of Washington.

Proteomics Laboratory: Richard D. Smith, Ph.D., Thomas O. Metz, Ph.D., Charles Ansong, Ph.D., Bobbie-Jo Webb-Robertson, Ph.D., and Hugh D. Mitchell, Ph.D. Pacific Northwest National Laboratory.

Repository: Heather Higgins, Sandra Ke. NIDDK Biosample Repository at Fisher BioServices.

RNA Laboratory and Gene Expression Laboratory: Jin-Xiong She, Ph.D., PI^{1,3,4,11}, Richard McIndoe, Ph.D., Haitao Liu, M.D., John Nechtman, Yansheng Zhao, Na Jiang, M.D., Yanna Tian, M.S. Guangkuo Dong, M.S. Jinfinity Biosciences, LLC.

SNP Laboratory: Stephen S. Rich, Ph.D.³, Wei-Min Chen, Ph.D.³, Suna Onengut-Gumuscu, Ph.D.³, Emily Farber, Rebecca Roche Pickin, Ph.D., Jordan Davis, Dan Gallo, Jessica Bonnie, Paul Campolieto. Center for Public Health Genomics, University of Virginia.

Committees:

¹Ancillary Studies, ²Diet, ³Genetics, ⁴Human Subjects/Publicity/Publications, ⁵Immune Markers, ⁶Infectious Agents, ⁷Laboratory Implementation, ⁸Maternal Studies, ⁹Psychosocial, ¹⁰Quality Assurance, ¹¹Steering, ¹²Study Coordinators, ¹³Celiac Disease, ¹⁴Clinical Implementation, ¹⁵Quality Assurance Subcommittee on Data Quality.

References

- Adam EK, Kumari M. Assessing salivary cortisol in large-scale, epidemiological research. *Psychoneuroendocrinology*. 2009; 34(10):1423–1436. [PubMed: 19647372]
- Bäumler D, Kirschbaum C, Kliegel M, Alexander N, Stalder T. The cortisol awakening response in toddlers and young children. *Psychoneuroendocrinology*. 2013; 38(11):2485–2492. DOI: 10.1016/j.psyneuen.2013.05.008 [PubMed: 23768972]
- Baxter J, Vehik K, Johnson SB, Lernmark B, Roth R, Simell T, Teddy Study Group. Differences in recruitment and early retention among ethnic minority participants in a large pediatric cohort: the TEDDY Study. *Contemporary Clinical Trials*. 2012; 33(4):633–640. DOI: 10.1016/j.cct.2012.03.009 [PubMed: 22484339]
- Bright MA, Granger DA, Frick JE. Do infants show a cortisol awakening response? *Developmental Psychobiology*. 2012; 54(7):736–743. DOI: 10.1002/dev.20617 [PubMed: 22006547]
- Broderick JE, Arnold D, Kudielka BM, Kirschbaum C. Salivary cortisol sampling compliance: comparison of patients and healthy volunteers. *Psychoneuroendocrinology*. 2004; 29(5):636–650. DOI: 10.1016/S0306-4530(03)00093-3 [PubMed: 15041086]
- Bruce J, Fisher PA, Pears KC, Levine S. Morning cortisol Levels in preschool-aged foster children: differential effects of maltreatment type. *Developmental Psychobiology*. 2009; 51(1):14–23. DOI: 10.1002/dev.20333 [PubMed: 18720365]
- Chiappin S, Antonelli G, Gatti R, De Palo EF. Saliva specimen: a new laboratory tool for diagnostic and basic investigation. *Clinica Chimica Acta*. 2007; 383(1–2):30–40. DOI: 10.1016/j.cca.2007.04.011
- Clements AD, Parker CR. The relationship between salivary cortisol concentrations in frozen versus mailed samples. *Psychoneuroendocrinology*. 1998; 23(6):613–616. [PubMed: 9802131]
- Davis EP, Granger DA. Developmental differences in infant salivary alpha-amylase and cortisol responses to stress. *Psychoneuroendocrinology*. 2009; 34(6):795–804. DOI: 10.1016/j.psyneuen.2009.02.001 [PubMed: 19268476]
- Donzella B, Talge NM, Smith TL, Gunnar MR. To spear or not to spear: comparison of saliva collection methods. *Developmental Psychobiology*. 2008; 50(7):714–717. DOI: 10.1002/dev.20340 [PubMed: 18767029]
- Felt BT, Mollen E, Diaz S, Renaud E, Zeglis M, Wheatcroft G, Mendelow D. Behavioral interventions reduce infant distress at immunization. *Archives of Pediatric and Adolescent Medicine*. 2000; 154(7):719–724.
- Freitag CM, Hänig S, Palmason H, Meyer J, Wüst S, Seitz C. Cortisol awakening response in healthy children and children with ADHD: impact of comorbid disorders and psychosocial risk factors. *Psychoneuroendocrinology*. 2009; 34(7):1019–1028. DOI: 10.1016/j.psyneuen.2009.01.018 [PubMed: 19278790]
- Gribbin CE, Watamura SE, Cairns A, Harsh JR, Lebourgeois MK. The cortisol awakening response (CAR) in 2- to 4-year-old children: effects of acute nighttime sleep restriction, wake time, and daytime napping. *Developmental Psychobiology*. 2012; 54(4):412–422. DOI: 10.1002/dev.20599 [PubMed: 21953381]
- Gunnar, MR., Talge, NM. Neuroendocrine Measures in Developmental Research. In: Schmidt, LA., Segalowitz, SJ., editors. *Developmental Psychophysiology: Theory, Systems, and Methods*. Cambridge: Cambridge University Press; 2007. p. 343–366.
- Gunnar MR, Talge NM, Herrera A. Stressor paradigms in developmental studies: what does and does not work to produce mean increases in salivary cortisol. *Psychoneuroendocrinology*. 2009; 34(7): 953–967. DOI: 10.1016/j.psyneuen.2009.02.010 [PubMed: 19321267]
- Gunnar, MR., White, BP. Salivary Cortisol Measures in Infant and Child Assessment. In: Singer, LT., Zeskind, PS., editors. *Biobehavioral Assessment of the Infant*. New York: Guilford Press; 2001. p. 167–189.
- Hall DL, Blyler D, Allen D, Mishel MH, Crandell J, Germino BB, Porter LS. Predictors and patterns of participant adherence to a cortisol collection protocol. *Psychoneuroendocrinology*. 2011; 36(4): 540–546. DOI: 10.1016/j.psyneuen.2010.08.008 [PubMed: 20863625]

- Hanrahan K, McCarthy AM, Kleiber C, Lutgendorf S, Tsalikian E. Strategies for salivary cortisol collection and analysis in research with children. *Applied Nursing Research*. 2006; 19(2):95–101. DOI: 10.1016/j.apnr.2006.02.001 [PubMed: 16728293]
- Hellhammer DH, Wust S, Kudielka BM. Salivary cortisol as a biomarker in stress research. *Psychoneuroendocrinology*. 2009; 34(2):163–171. DOI: 10.1016/j.psyneuen.2008.10.026 [PubMed: 19095358]
- Jessop DS, Turner-Cobb JM. Measurement and meaning of salivary cortisol: a focus on health and disease in children. *Stress*. 2008; 11(1):1–14. DOI: 10.1080/10253890701365527 [PubMed: 17853059]
- Johnson SB, Lee HS, Baxter J, Lernmark B, Roth R, Simell T, Teddy Study Group. The Environmental Determinants of Diabetes in the Young (TEDDY) study: predictors of early study withdrawal among participants with no family history of type 1 diabetes. *Pediatric Diabetes*. 2011; 12(3 Pt 1): 165–171. DOI: 10.1111/j.1399-5448.2010.00686.x [PubMed: 21029290]
- Johnson SB, Lynch K, Lee H, Smith L, Baxter J, Lernmark B, Roth R, Simell T, TEDDY Study Group. At high risk for early withdrawal: Using a cumulative risk model to increase retention in the first year of the TEDDY study. *Journal of Clinical Epidemiology*. 2014; 67:609–611. [PubMed: 24725640]
- Johnson SB, Lynch K, Baxter J, Lernmark B, Roth R, Simell T, Smith L, The TEDDY Study Group. Predicting Later Study Withdrawal in Participants Active in a Longitudinal Birth Cohort Study for 1 Year. *The TEDDY Study Journal of Pediatric Psychology*. 2016; 2016:373–383. DOI: 10.1093/jpepsy/jsv092
- Kirschbaum C, Pirke KM, Hellhammer DH. The ‘Trier Social Stress Test’—a tool for investigating psychobiological stress responses in a laboratory setting. *Neuropsychobiology*. 1993; 28(1–2):76–81. doi:119004. [PubMed: 8255414]
- Kudielka BM, Hawkey LC, Adam EK, Cacioppo JT. Compliance with ambulatory saliva sampling in the Chicago Health, Aging, and Social Relations Study and associations with social support. *Annals of Behavioral Medicine*. 2007; 34(2):209–216. DOI: 10.1080/08836610701566951 [PubMed: 17927559]
- Lernmark B, Lynch K, Baxter J, Roth R, Simell T, Smith L, Swartling U, Johnson SB, TEDDY Study Group. Participant Experiences in the Environmental Determinants of Diabetes in the Young Study: Common Reasons for Withdrawing. *Journal of Diabetes Research*. 2016. Article ID 2720650, 13 pages <http://dx.doi.org/10.1155/2016/2720650>
- McCarthy AM, Hanrahan K, Kleiber C, Zimmerman MB, Lutgendorf S, Tsalikian E. Normative salivary cortisol values and responsivity in children. *Applied Nursing Research*. 2009; 22(1):54–62. DOI: 10.1016/j.apnr.2007.04.009 [PubMed: 19171296]
- McCarthy AM, Hanrahan K, Scott LM, Zemblidge N, Kleiber C, Zimmerman MB. Salivary cortisol responsivity to an intravenous catheter insertion in children with attention-deficit/hyperactivity disorder. *Journal of Pediatric Psychology*. 2011; 36(8):902–910. DOI: 10.1093/jpepsy/jsr012 [PubMed: 21389037]
- Nater UM, Rohleder N, Schlotz W, Ehler U, Kirschbaum C. Determinants of the diurnal course of salivary alpha-amylase. *Psychoneuroendocrinology*. 2007; 32(4):392–401. DOI: 10.1016/j.psyneuen.2007.02.007 [PubMed: 17418498]
- Neu M, Goldstein M, Gao D, Laudenslager ML. Salivary cortisol in preterm infants: Validation of a simple method for collecting saliva for cortisol determination. *Early Human Development*. 2007; 83(1):47–54. DOI: 10.1016/j.earlhumdev.2006.04.003 [PubMed: 16766144]
- Putnam SK, Lopata C, Fox JD, Thomeer ML, Rodgers JD, Volker MA, Lee GK, Neilans EG, Werth J. Comparison of saliva collection methods in children with high-functioning autism spectrum disorders: acceptability and recovery of cortisol. *Child Psychiatry and Human Development*. 2012; 43(4):560–573. DOI: 10.1007/s10578-012-0284-3 [PubMed: 22331443]
- Rohleder N, Nater UM. Determinants of salivary alpha-amylase in humans and methodological considerations. *Psychoneuroendocrinology*. 2009; 34(4):469–485. DOI: 10.1016/j.psyneuen.2008.12.004 [PubMed: 19155141]
- Saridjan NS, Huizink AC, Koetsier JA, Jaddoe VW, Mackenbach JP, Hofman A, Kirschbaum C, Verhulst FC, Tiemeier H. Do social disadvantage and early family adversity affect the diurnal

- cortisol rhythm in infants? The Generation R Study. *Hormones and Behavior*. 2010; 57(2):247–254. DOI: 10.1016/j.yhbeh.2009.12.001 [PubMed: 20006614]
- Schwartz EB, Granger DA, Susman EJ, Gunnar MR, Laird B. Assessing salivary cortisol in studies of child development. *Child Development*. 1998; 69(6):1503–1513. [PubMed: 9914636]
- Smith VC, Dougherty LR. Noisy spit: parental noncompliance with child salivary cortisol sampling. *Developmental Psychobiology*. 2014; 56(4):647–656. DOI: 10.1002/dev.21133 [PubMed: 23754778]
- Smyth N, Clow A, Thorn L, Hucklebridge F, Evans P. Delays of 5–15 min between awakening and the start of saliva sampling matter in assessment of the cortisol awakening response. *Psychoneuroendocrinology*. 2013; 38:1476–1483. [PubMed: 23312064]
- Stalder T, Bäumler D, Miller R, Alexander N, Kliegel M, Kirschbaum C. The cortisol awakening response in infants: ontogeny and associations with development-related variables. *Psychoneuroendocrinology*. 2013; 38(4):552–559. DOI: 10.1016/j.psyneuen.2012.07.015 [PubMed: 22920911]
- Stalder T, Kirschbaum C, Kudielka BM, Adam EK, Pruessner JC, Wüst S, Dockray S, Smyth N, Evans P, Hellhammer DH, Miller R, Wetherell MA, Lupien SJ, Clow A. Assessment of the cortisol awakening response: Expert consensus guidelines. *Psychoneuroendocrinology*. 2016; 63:414–432. [PubMed: 26563991]
- The TEDDY Study Group. The Environmental Determinants of Diabetes in the Young (TEDDY) study: study design. *Pediatric Diabetes*. 2007; 8(5):286–298. DOI: 10.1111/j.1399-5448.2007.00269.x [PubMed: 17850472]
- The TEDDY Study Group. The Environmental Determinants of Diabetes in the Young (TEDDY) Study. *Annals of the New York Academy of Sciences*. 2008; 1150:1–13. DOI: 10.1196/annals.1447.062
- Tegethoff M, Knierzinger N, Meyer AH, Meinlschmidt G. Cortisol awakening response in infants during the first six postnatal months and its relation to birth outcome. *Psychoneuroendocrinology*. 2013; 38(5):629–637. DOI: 10.1016/j.psyneuen.2012.08.002 [PubMed: 22951380]
- Tikotzki L, Sadeh A. Sleep patterns and sleep disruptions in kindergarten children. *Journal of Clinical Child Psychology*. 2001; 30:581–591. [PubMed: 11708244]

Table 1
Saliva Sample Collection Among Eligible TEDDY Subjects at United States (US) and European (EU) Centers

	42 month visit		54 month visit		66 month visit		Total	
	N	%	N	%	N	%	N	%
Total								
Completed visit	4307		4545		3838		12690	
Not offered SSP*	207	4.8%	345	7.6%	262	6.8%	814	6.4%
Total eligible for SSP	4100	95.2%	4200	92.4%	3576	93.2%	11876	93.6%
1 sample	3948	96.3%	4016	95.6%	3426	95.8%	11390	95.9%
Refused	152	3.7%	184	4.4%	150	4.2%	486	4.1%
Completed visits	1090		1402		1221		3713	
US								
Not offered SSP	141	12.9%	247	17.6%	164	13.4%	552	14.9%
Total eligible for SSP	949	87.1%	1155	82.4%	1057	86.6%	3161	85.1%
1 sample	935	98.5%	1128	97.7%	1038	98.2%	3101	98.1%
Refused	14	1.5%	27	2.3%	19	1.8%	60	1.9%
Completed visits	3217		3143		2617		8977	
EU								
Not offered SSP	66	2.1%	98	3.1%	98	3.7%	262	2.9%
Total eligible for SSP	3151	97.9%	3045	96.9%	2519	96.3%	8715	97.1%
1 sample	3013	95.6%	2888	94.8%	2388	94.8%	8289	95.1%
Refused	138	4.4%	157	5.2%	131	5.2%	426	4.9%

* Salivary Sample Protocol – SSP; United States (US), Europe (EU)

Table 2

Completion of Salivary Sample Collection for TEDDY Study Overall, United States (US) and European (EU) Centers

		42 month visit		54 month visit		66 month visit		Total	
		N	%	N	%	N	%	N	%
Total	1 cortisol samples	3948		4016		3426		11390	
	No morning sample	134	3.4%	241	6.0%	180	5.3%	555	4.9%
	No clinic sample	21	0.5%	32	0.8%	33	1.0%	86	0.8%
	Sample excluded *	129	3.3%	106	2.6%	85	2.5%	320	2.8%
US	1 cortisol samples	935		1128		1038		3101	
	No morning sample	87	9.3%	142	12.6%	117	11.3%	346	11.2%
	No clinic sample	9	1.0%	11	1.0%	15	1.4%	35	1.1%
	Sample excluded *	42	4.5%	40	3.5%	33	3.2%	115	3.7%
EU	1 cortisol samples	3013		2888		2388		8289	
	No morning sample	47	1.6%	99	3.4%	63	2.6%	209	2.5%
	No clinic sample	12	0.4%	21	0.7%	18	0.8%	51	0.6%
	Sample excluded *	87	2.9%	66	2.3%	52	2.2%	205	2.5%

* Sample exclusion reasons: child brushed teeth without rinsing, ate or drank less than 30 minutes before cortisol sample collection or were on oral steroids in last 30 days; United States (US), Europe (EU)