Published in final edited form as:

Nature. 2017 February 02; 542(7639): 43–48. doi:10.1038/nature20818.

IL-17 is a neuromodulator of *Caenorhabditis elegans*' sensory responses

Changchun Chen¹, Eisuke Itakura^{1,2}, Geoffrey M. Nelson¹, Ming Sheng¹, Patrick Laurent^{1,3}, Lorenz A. Fenk¹, Rebecca A. Butcher⁴, Ramanujan S. Hegde¹, and Mario de Bono^{1,*}

¹MRC Laboratory of Molecular Biology, Francis Crick Avenue, Cambridge CB2 0QH, UK

⁴Department of Chemistry, University of Florida, Gainesville, FL, 32611, USA

Abstract

Interleukin 17 (IL-17) is a major pro-inflammatory cytokine: it mediates responses to pathogens or tissue damage, and drives autoimmune diseases. Little is known about its role in the nervous system. Here we show that IL-17 has neuromodulator-like properties in *C. elegans*. IL-17 can act directly on neurons to alter their response properties and contribution to behavior. Using unbiased genetic screens we delineate an IL-17 signaling pathway and show that it acts in the RMG hub interneurons. Disrupting IL-17 signaling reduces RMG responsiveness to input from O₂-sensors, and renders sustained escape from 21% O₂ transient and contingent on additional stimuli. Over-activating IL-17 receptors abnormally heightens responses to 21% O₂ in RMG neurons and whole animals. IL-17 deficiency can be bypassed by optogenetic stimulation of RMG. Inducing IL-17 expression in adults can rescue mutant defects within 6 hours. These findings reveal a non-immunologic role of IL-17 modulating circuit function and behavior.

Introduction

The contributions of a neuron to circuit activity and behavior depend on its responsiveness to upstream inputs, and its ability to drive downstream outputs. By modifying the input–output relationships between neurons, neuromodulators play crucial roles in information processing

Correspondence and requests for materials should be addressed to M.d.B (debono@mrc-lmb.cam.ac.uk).

²Present address: Department of Nanobiology, Graduate School of Advanced Integration Science, Chiba University, Chiba, 263-8522, Japan.

³Present address: Université Libre de Bruxelles, Lab of Neurophysiology, Campus Erasme, 808 route de lennik, 1070 Bruxelles, Belgium

Author Contributions C.C., E.I., R.S.H. and M.d.B designed experiments. C.C., E.I., M.S., P.L., L.A.F performed experiments. G.N. did the genome sequence data analysis. R.A.B provided reagents. C.C., E.I., R.S.H., and M.d.B analyzed the data; C.C. and M.d.B. wrote the paper.

Author Information Reprints and permissions information is available at www.nature.com/reprints.

The authors declare no competing financial interests.

Code Availability

Zentracker, the custom-written Matlab software used to analyze worm movies is available at: <https://github.com/wormtracker/zentracker>.

Neuron Analyzer, the custom-written Matlab program used to analyze neural image stacks is available at: <https://github.com/neuronanalyser/neuronanalyser>.

Data Availability

The data that support the findings of this study are available from the corresponding author.

by the brain: they are thought to underlie changes in arousal, mood, and global animal states^{1–4}, and to contribute to learning and forgetting^{5,6}. Identifying neuromodulatory signaling pathways and understanding their *in vivo* effects on circuit function are important goals to understand the neural basis for behavior⁷.

The comparatively simple *C. elegans* nervous system allows behaviors to be linked to small numbers of identified neurons connected by few, anatomically defined, chemical and electrical connections^{8,9} (wormwiring.org). Aggregation is an intricate *C. elegans* behavior governed by multiple sensory neurons, including receptors for O₂ (URX, AQR, PQR), noxious stimuli (ASH, ADL), food cues (ADF, ASK), and pheromones (ASJ, ADL, ASK)^{10–13}. Anatomically-defined gap junctions connect the URX, ASH, ADL, and ASK neurons with RMG interneurons⁹, which process sensory information to control aggregation¹³. Here, we use forward genetics to probe molecular mechanisms regulating the properties of these neurons and connections.

IL-17 promotes *C. elegans* aggregation

We mutagenized aggregating *npr-1(null)* (*neuropeptide receptor 1*) animals and identified > 800 mutants that aggregated poorly but were otherwise overtly wild-type. We sequenced the genomes of 583 mutants, and identified the lesions causing their aggregation defects. Six harbored mutations in Y64G10A.6, an *interleukin 17* cytokine receptor ortholog we named *ilcr-1* (Extended data Fig. 1a; Extended data Table 1). An *ilcr-1* deletion, *tm5866*, disrupted aggregation and was rescued by expressing *ilcr-1* cDNA from the *ilcr-1* promoter (Fig. 1a). The *C. elegans* genome encodes a second IL-17-like receptor, F56D1.2, that we named *ilcr-2*. We did not identify *ilcr-2* alleles in our collection but CRISPR-generated *ilcr-2* nulls failed to aggregate (Fig. 1a; Extended data Fig. 1a–d). Expressing *ilcr-2* cDNA from the *ilcr-2* promoter restored aggregation to *ilcr-2* but not *ilcr-1* mutants, and vice-versa (Fig. 1a; data not shown). Like their mammalian homologs¹⁴, ILCR-1 and ILCR-2 have an extracellular domain for ligand binding, a single transmembrane domain, and an intracellular SEF/IL-17R (SEFIR) signaling domain (Extended data Fig. 1f).

Three *C. elegans* genes, T22H6.1, F25D1.3 and C44B12.6, share homology with mammalian IL-17s (Extended data Fig. 1g; data not shown). We did not find F25D1.3 or C44B12.6 alleles in our collection, and disrupting either gene did not perturb aggregation (Extended data Fig. 1d, e). We found four alleles of T22H6.1, including splice-site mutations (Extended data Table 1). Deletions of T22H6.1, *tm5124* and *tm5218*, disrupted aggregation, and were rescued by expressing T22H6.1 cDNA from its promoter (Fig. 1a). T22H6.1, which we named *ilc-17.1* (*interleukin 17* cytokine related gene 1), could therefore encode a ligand for ILCR-1/ILCR-2 receptors. Double and triple mutants of *ilc-17.1*, *ilcr-1* and *ilcr-2* behaved like single mutants (Extended data Fig. 1c; see below), suggesting these genes function in a single pathway.

GFP-tagged ILCR-1 co-localized with HA-tagged ILCR-2 at the plasma membrane of HEK293T cells (Fig. 1b), and ILCR-1-GFP specifically co-precipitated with ILCR-2-HA (Fig. 1c). Like mammalian IL-17s, ILC-17.1 formed disulfide-linked dimers¹⁵ (Extended data Fig. 1h). ILC-17.1 specifically interacted with the surface of cells expressing ILCR-2, or

ILCR-2 and ILCR-1, but not control cells, or cells expressing only ILCR-1 (Fig. 1d, e). In mammals, IL-17 binding to one receptor type is thought to increase affinity for the second receptor type, promoting heteromeric complex formation¹⁶. Together, our data suggest ILC-17.1 signals via ILCR-1/ILCR-2 co-receptors.

C. elegans aggregation is stimulated by 21% O₂^{12,17–19}, and many aggregation-defective mutants exhibit O₂ response defects. On bacterial food *npr-1* animals respond to a 7% - 21% rise in O₂ by increasing their average speed from ~40 to ~180 µm/sec. This arousal is sustained for hours if animals are kept at 21% O₂, with only modest attenuation (Fig. 1f)²⁰. *ilc-17.1*, *ilcr-1* and *ilcr-2* mutants responded to 21% O₂ immediately after being transferred to the assay plates, but unusually, their arousal declined to baseline within 1 hour, and they became unresponsive to 21% O₂ thereafter (Fig. 1f–i). These defects could be rescued by expressing the corresponding wild type gene either from extrachromosomal multicopy arrays (Fig. 1f, g) or a single copy insertion (Extended data Fig. 1i)²¹. Knocking out *ilc-17.1* or *ilcr-1* in animals carrying the natural *npr-1* 215F allele recapitulated the defects we observed in *npr-1*(null) mutants (Extended data Fig. 1j).

IL-17 increases RMG interneuron gain

Functional ILCR-1 and ILCR-2 receptors C-terminally-tagged with GFP were expressed in most neurons, including RMG interneurons (Fig. 2a–d). *ilcr-2* was also expressed in pharyngeal muscle. Selectively expressing *ilcr-1* or *ilcr-2* cDNA in the O₂ sensors, using the *gcy-32* promoter, only weakly rescued *ilcr* mutant phenotypes (Extended data Fig. 2a–d). By contrast, expressing receptors in RMG, using the *npr-1* or *flp-5* promoters, robustly restored persistent escape from 21% O₂ and aggregation to *ilcr* mutants (Extended data Fig. 2a–d). The *flp-5* promoter drives expression in ASG (previously reported as ASE)²² and RMG, but expressing *ilcr-1* cDNA in ASG using the *ops-1* promoter did not rescue *ilcr-1* phenotypes (Extended data Fig. 2e); neither did expression in cholinergic neurons (*unc-17* or *acr-2* promoters), GABAergic neurons (*unc-25* promoter), or interneurons controlling forward and reverse movement (*glr-1* or *nmr-1* promoters) (Extended data Fig. 2f). To confirm RMG as the focus of action we inserted a single copy of *ilcr-1* in which a strong terminator of transcription, flanked by LoxP sites, separated the promoter and open reading frame. As expected, this transgene did not rescue the *ilcr-1* mutant phenotype unless we also selectively expressed Cre recombinase in RMG, using the *flp-5* or *npr-1* promoters (Fig. 2e). These data suggest, the ILCR co-receptors act predominantly in the RMG interneurons to promote aggregation and escape from 21% O₂.

Functional mCherry-tagged ILC-17.1 accumulated in coelomocytes, scavenger cells that endocytose macromolecules from *C. elegans*' body cavity²³, confirming that ILC-17.1 is secreted (Fig. 2f, top panels). A functional bicistronic construct co-expressing *ilc-17.1* and free mCherry labeled the RMG, AUA and ASE neurons (Fig. 2f, bottom panels). Targeted expression of *ilc-17.1* in RMG restored the O₂ response, suggesting the ligand could act as an autocrine signal (Extended data Fig. 2g). However, expressing *ilc-17.1* in the AWA olfactory neurons or the ASE gustatory neurons, also rescued the *ilc-17.1* mutant phenotype (Extended data Fig. 2h), suggesting the ligand source is unimportant. Expression and release of ILC-17.1 from RMG appears to be O₂-independent: mCherry-ILC-17.1 accumulation in

coelomocytes was similar in animals cultivated at 7% and 21% O₂ (Fig. 2g), as were RMG levels of cytoplasmic mCherry expressed from the bicistronic *pilc-17.1::ilc-17.1::sl2::mCherry* transgene (data not shown). Constitutive *ilc-17.1* expression from the *ges-1* or *vha-6* gut-specific promoters rescued the *ilc-17.1* defects (Fig. 2h). The O₂-regulated behavior of these animals suggests *ilc-17.1* does not itself drive arousal, and can promote O₂ circuit function without being regulated by O₂. However, this does not exclude that *ilc-17.1* expression, or its effects on RMG, are regulated by interacting inputs, as happens in the mammalian immune system^{24–25}.

To test if ILC-17.1 altered RMG responsiveness to input from O₂ sensors we used YC2.60 Ca²⁺ sensors²⁶ to image RMG and URX (as a control) in freely moving animals under our behavioral assay conditions. URX Ca²⁺ responses to 21% O₂ were not altered in *ilc-17.1* or *ilcr-2* mutants compared to controls, either 10 minutes or 2 hours after animals were picked onto the assay plate (Fig. 3a). By contrast, RMG Ca²⁺ responses to 21% O₂ were reduced at both time points in mutants (Fig. 3b). Expressing *ilcr-2* in RMG rescued these defects in *ilcr-2* mutants (Extended data Fig. 2i). Imaging RMG in an *npr-1 215F* background revealed similar phenotypes to those in *npr-1* null mutants (Extended data Fig. 2j). As an independent assay of RMG activity we measured *pflp-5::gfp* expression (*FMR*amide-like neuropeptide) in RMG. Expression from this neuropeptide reporter increases with RMG activity²⁷. *ilc-17.1*, *ilcr-1* and *ilcr-2* mutants kept at 21% O₂ showed reduced *pflp-5::gfp* fluorescence compared to *npr-1* controls, consistent with reduced RMG activity (Extended data Fig. 2k). Disrupting *ilc-17.1* reduces RMG responsiveness to O₂ input without compromising its ability to signal to downstream neurons. Stimulating RMG using Channelrhodopsin evoked similar behavioral responses in *ilc-17.1* mutants and control animals (Fig. 3c).

RMG interneurons influence *C. elegans*' pheromone responses, and are coupled by gap junctions to ASK and ADL pheromone-responsive neurons^{9,13,28}. Using Ca²⁺ imaging we found that a mixture of C3+C6+C9 ascarosides^{29,30} evoked a response in RMG reminiscent of that evoked by the same ascaroside mix in ASK neurons¹³. This response was strongly reduced in *ilc-17.1* mutants (Fig. 3d), perhaps because RMG neurons have lower Ca²⁺ at 21% O₂. Moreover, unlike *npr-1* animals, which are attracted to this ascaroside mix¹³, *ilc-17.1* mutants were repelled by it (Fig. 3e). Thus, ILC-17.1 signaling influences pheromone responses.

IL-17 signaling components pathway are expressed broadly in the nervous system. Do they affect all behavioral responses? *ilc-17.1* mutants showed normal chemotaxis to NaCl, mediated by the ASE neurons, and benzaldehyde, mediated by AWC, and did not alter benzaldehyde-evoked Ca²⁺ responses in AIB interneurons (Extended data Fig. 2l-n). Thus, IL-17 defects do not cause general circuit dysfunction.

Why do mutants with chronic defects in RMG's responsiveness to 21% O₂ escape this cue 10 minutes but not 2 hours after being picked for assay? We speculated that physically transferring animals, which stimulates mechanoreceptors, transiently makes reduced RMG output sufficient to drive escape from 21% O₂. Mechanosensory stimulation is known to transiently increase locomotor speed through activation of gentle touch neurons (Y. Tanizawa, B. Zhao and W. Schafer, personal communication). Consistent with our

hypothesis, gently tapping the microscope stage 2 hours after transferring animals for assay instantly restored responsiveness to 21% O₂ to *ilc* signaling mutants, as long as their gentle touch receptors were functional (Fig. 3f). Thus, without *ilc-17.1* signaling *C. elegans* requires additional sensory stimulation to escape 21% O₂.

Heat-shock induced expression of *ilc-17.1*, *ilcr-1* or *ilcr-2* cDNA in adults rescued the respective mutant (Fig. 3g, h; Extended data Fig. 3). Experiments in which we changed heat-shock duration while monitoring expression of a *phsp-16.41::ilc-17.1::mCherry* bicistronic transgene and O₂ responses suggested that ILC-17.1 signaling requires < 6 hours to exert its effects (Extended data Fig. 4). Faster induction of ILC-17.1 expression using longer heat shock does not shorten the time required for rescue.

Elevating IL-17 heightens O₂ responses

Animals overexpressing *ilc-17.1* from its own promoter, the *hsp-16.41*, *flp-5*, *odr-10* (AWA), *flp-6* (ASE), or gut promoters all showed abnormally heightened responses to 21% O₂ (Fig. 1f, 4a; Extended data Figs. 2g, h; Extended data Fig. 4). Moreover, overexpressing *ilc-17.1* in the non-aggregating N2 lab strain conferred aggregation behavior and responsiveness to 21% O₂ (Fig. 4b, c). Neural imaging showed that ILC-17.1 overexpression significantly increased RMG Ca²⁺ responses, both in N2 and in *npr-1* animals (Fig. 4d, e). It also increased *pflp-5::gfp* expression in RMG in N2 and *npr-1* animals kept at 21% O₂, consistent with increased RMG output (Extended data Fig. 5). However, animals overexpressing *ilc-17.1* behaved appropriately at 7% O₂, and had unaltered RMG Ca²⁺ levels at 7% O₂, suggesting ILC-17.1 overexpression did not itself depolarize RMG. O₂-evoked responses in RMG depend on URX27. Over-expressing *ilc-17.1* did not alter O₂-evoked Ca²⁺ response in URX (Fig. 4f).

An IL-17 signaling pathway in neurons

A candidate approach did not identify components acting downstream of the ILC-17.1 receptors (Extended data Fig. 6). We therefore screened our collection for mutants exhibiting the unusual *ilc-17.1* O₂-response defect. Eight strains had such defects, highlighting 3 genes. One gene encoded an Act1 homolog we named *actl-1* (*Act1-lke*) (Extended data Fig. 7). Mammalian Act1 binds IL-17 receptors via SEFIR–SEFIR domain interactions and mediates their signaling output³¹. ACTL-1 has the Death domain-linker-SEFIR domain architecture found in non-vertebrate Act1 homologs, and lacks the helix-loop-helix and U-box found in mammalian Act1 (Extended data Fig. 7)³². Pull-downs from HEK293T cells co-expressing ACTL-1, ILCR-1 and ILCR-2 showed that ACTL-1 physically interacts with ILCR-1/ILCR-2, primarily via association with ILCR-2 (Fig. 5a, b) - consistent with ACTL-1 being an adaptor for *C. elegans* IL-17 receptors. The second gene encoded the sole *C. elegans* IRAK (interleukin-1 receptor-associated kinase) *pik-1* (Extended data Fig. 8)³³. PIK-1-FLAG and ACTL-1-HA co-expressed in HEK cells formed a complex (Fig. 5c, d), suggesting these two proteins interact physically. The remaining gene encodes the sole *C. elegans* homolog of the atypical IκB NF-κB inhibitors NFKBID and NFKBIZ/IκBζ^{34–36}, which we called *nfki-1* (Extended data Fig. 9). In mammals IκBζ contributes to immune responses mediated by IL-1β, Toll-like receptors, and IL-17, but its *in vivo* functions are

poorly understood^{37,38}. *actl-1*, *pik-1* or *nfki-1* mutant defects could be rescued by a single copy insertion of the corresponding wild type gene generated by MosSCI (Extended data Fig. 7c). Various double mutants of *ilc-17.1*, *actl-1*, *pik-1* and *nfki-1* did not show enhanced behavioral phenotypes (Extended data Fig. 7–9), suggesting their gene products act in the same pathway. Overexpressing wild type *nfki-1* restored aggregation and sustained escape from 21% O₂ to *ilcr-1*, *ilcr-2*, *actl-1* and *pik-1* null mutants (Fig. 5e), suggesting that NFKI-1 acts downstream of these signaling components.

actl-1, *pik-1*, and *nfki-1* were expressed in many neurons including RMG (Extended data Fig. 8d; Extended data Fig. 9c; data not shown). Expressing their cDNA in RMG using the *flp-5* promoter rescued the mutant phenotype of each gene (Fig. 5f–h), although not completely. Expressing one or more of *actl-1*, *pik-1* or *nfki-1* from the *ops-1* (ASG neurons), *gcy-32* (AQR, PQR and URX), or *flp-6* (ASE) promoters failed to rescue the corresponding mutant phenotypes, whereas expressing *nfki-1* from the *npr-1* promoter, which drives expression in RMG and other neurons, rescued *nfki-1* phenotypes (Fig. 5f–h). Incomplete rescue of the *flp-5* driven transgenes suggests these genes may act in other neurons besides RMG to regulate O₂ responses, especially since the more broadly expressed *npr-1* promoter conferred significantly better rescue of the *nfki-1* phenotype (Fig. 5h). However, we cannot exclude other explanations, such as variability across transgenes.

Ca²⁺ imaging experiments showed that *actl-1*, *pik-1*, and *nfki-1* mutants recapitulate *ilc-17.1* phenotypes: reduced RMG responsiveness to 21% O₂ (Fig. 5i–k). The RMG Ca²⁺ response defects were non-additive in various double mutant combinations of *ilc-17.1*, *actl-1*, *pik-1* and *nfki-1* (Fig. 5i–k), as expected if these genes' products function together in a signaling pathway. The ILCR-1/2–ACTL-1–PIK-1/IRAK relationship is reminiscent of Toll-like and IL1- β receptor signaling pathways, in which the adaptor MyD88 recruits IRAKs via homotypic Death domains interactions (Fig. 5l). TIR and SEFIR domains are related domains³⁹. Consistent with this evolutionary relationship, in the mammalian immune system IL-17 and TLR receptor pathways activate similar downstream genes¹⁴. *C. elegans* aggregation and arousal at 21% O₂ is influenced by the presence of bacteria, and modulates pathogen responses⁴⁰. Plausibly, IL-17 signaling may have played a role in ancient nervous systems regulating behavioral responses to bacteria.

Conclusions

IL-17 is a potent pro-inflammatory cytokine. Its production in mammals is induced by infection, tissue damage, and stress, and is thought to drive tissue inflammation and autoimmune disease. By contrast, its roles in regulating neural circuit functions are elusive, although members of the IL-17 and IL-17 receptor family are expressed in brain¹⁴. Our data show that physiological levels of IL-17 can act directly on neurons to increase their responsiveness to pre-synaptic input. Elevating IL-17 levels can abnormally increase circuit sensitivity to O₂ input. By potentiating circuit function, IL-17 signaling transforms an escape response to a noxious cue from one that is transient and contingent on other sensory inputs, to one that is unconditional and sustained.

Online Methods

Strains

C. elegans were maintained under standard conditions⁴¹. A detailed description of the forward genetic screens will be published elsewhere. The molecular lesions in alleles identified by forward genetics are listed in Extended Data Table 1. Frameshift mutations were generated using CRISPR-Cas9, as described⁴². Strains used are listed in Supplementary Table 1.

Behavioral assays

Aggregation and locomotion were assayed as described previously^{27,43}, with slight alterations. L4 animals were picked to a fresh plate 24 hours before assay. 50 animals were re-picked to the assay plate (an NGM plate seeded 2 days earlier with a 1 cm diameter OP50 lawn), and bordering and aggregation scored 2 and 24 hours later. The scorer was blind to genotype. To assay locomotion we tracked 30 animals 10 minutes and 2 hours after picking. We took care not to accidentally tap plates at the 2-hour time point. For behavioral assays lasting 9 minutes, statistical comparisons used a 1-minute window that starts 40 seconds after the switch from 7% to 21% O₂ and stops 20 seconds before the switch from 21% to 7% O₂. For the 2-hour long experiments, statistical comparisons used two 5-minutes windows, which start 5 minutes after the switch from 7% to 21% O₂, and 10 minutes before the assay ended. These intervals are sufficiently long to smooth out short-term fluctuations and highlight longer-term trends, while avoiding time periods when animals are rapidly changing their behavior in response to the O₂ stimulus. Movies were analyzed using Zentracker, a custom-written Matlab software (code availability: <https://github.com/wormtracker/zentracker>). All worms in the field of view were analyzed except animals those in contact with other animals. Speed was calculated as instantaneous worm centroid displacement between successive frames, and plotted as Mean $\pm$ S.E.M. Behavioral responses were robust: 3 - 4 biological replicates (each including 50 animals) for aggregation assays, and 30 animals for locomotion assays, provided sufficient statistical power to distinguish phenotypes. Statistical comparisons for speed assays used a two-tailed Mann-Whitney U test.

Quadrant chemotaxis assays: Chemotaxis to a mixture of C3, C6 and C9 ascarosides, each at 10 nM, was assayed as described previously¹³. Briefly, about 2 hours before the assay 200 adult animals were picked a fresh plate seeded with OP50. For the assay, animals were washed 3 times in chemotaxis buffer (1mM CaCl₂, 1mM MgSO₄ and 5mM KH₂PO₄/K₂HPO₄ pH 6.0) then placed in the center of the quadrant assay plate. Assay plates were freshly made with ascarosides added to alternate quadrants. The number of animals in each quadrant was scored after 20 mins, and the chemotaxis index was calculated as (the number of animals on ascaroside quadrants - the number of animals on control quadrants)/ (the number of animals on ascaroside quadrants + the number of animals on control quadrants).

Optogenetics: L4 animals expressing ChR2 in RMG27 were picked 24 hours before the assay and divided into two groups. One group was transferred to plates that had 100 μ l of 5 mM all-trans retinal (Sigma) in 100% ethanol added to the bacterial lawn. The other group

was put on plates with 100 μ l of ethanol as controls. Test and control plates were kept in the dark until animals were picked to assay plates 2 hours before recording. Videos were recorded using a Grasshopper camera (Point Grey) mounted on an M165FC Stereomicroscope (Leica) fitted with GFP excitation filters (ET 480/40x). Excitation light attenuated to 11.6mW/cm² was from a Leica EL6000 mercury lamp. Animals were recorded for 4 minutes without the blue light stimulus, followed by 4 minutes of blue light stimulation and a further 4 minutes without blue light. 7% O₂ was pumped into the microfluidic chamber throughout the assay. To record behavior we used trans-illumination, with light filtered through a 595 nm long-pass optical cast plastic filter (Edmund Optics).

Molecular biology

Expression plasmids were constructed using multisite Gateway (Life Technologies). Promoters used include: *ilc-17.1* (6kb), *ilcr-1* (6.5kb), *ilcr-2* (2.4kb), *npr-1*, *flp-5*, *flp-6*, *vha-6*, *ges-1*, *hsp-16.41*, *acr-2*, *glr-1*, *nmr-1*, *unc-17*, *unc-25*, *pik-1* (2.5kb), *actl-1* (7kb) and *nfki-1* (5.7kb). For rescue experiments, cDNA for each gene was amplified using the Phusion RT-PCR kit (Thermo Scientific) and cloned into pDonor 221.

Primers for *ilc-17.1* cDNA amplification were:

GGGGACAAGTTTGTACAAAAAAGCAGGCTTTTCAGAAAAATGCCAAAATCACCA CACAGGAC and

GGGGACCACTTTGTACAAGAAAGCTGGGTATTAAATTGGAATTTCTGCTTTGATTG TGTAGATAAAG; for *ilcr-1* cDNA amplification

GGGGACAAGTTTGTACAAAAAAGCAGGCTTTTCAGAAAAATGTTTCTTCATTCCG CTGCTCTTC and

GGGGACCACTTTGTACAAGAAAGCTGGGTATCAGTGTATACCAATGACTTCATCAT GGAAC; for *ilcr-2* cDNA amplification:

GGGGACAAGTTTGTACAAAAAAGCAGGCTTTTCAGAAAAATGGTCCGCCCTTTCG ACTACCA and

GGGGACCACTTTGTACAAGAAAGCTGGGTATCATGACTTCTTCACAACAATCGTC TTCC; for *pik-1* cDNA amplification:

GGGGACAAGTTTGTACAAAAAAGCAGGCTTTTCAGAAAAATGGATGATTCTCTTG GAGTATCAGAAGTTG and

GGGGACCACTTTGTACAAGAAAGCTGGGTATTAGACAAGTGGCGGAATTGAATTT TTACA; for *actl-1* cDNA amplification

GGGGACAAGTTTGTACAAAAAAGCAGGCTTTTCAGAAAAATGACTAAGATGAAA ATGGACGTAACAATTGAG and

GGGGACCACTTTGTACAAGAAAGCTGGGTATTATTGTGTAATACTGTAGTTCATGG AATCCTCG. And for *nfki-1* cDNA amplification

GGGGACAAGTTTGTACAAAAAAGCAGGCTTTTCAGAAAAATGGCAACCGTTGCC CCCAAG and

GGGGACCACTTTGTACAAGAAAGCTGGGTATCAAGCTCTCGACTTGTTCCGGGACT G. To generate a functional *mcherry-ilc-17.1* transgene mCherry coding sequences were inserted after the signal sequence between codon 31 and 32 of the *ilc-17.1* gene.

To express *C. elegans* proteins in HEK293T cells, cDNAs for each gene were amplified with primers having a 25 base overlap with sequences in pcDNA vectors and assembled with the vector using the Gibson Assembly kit (New England Biolabs). *ilc-17.1* cDNA without a signal sequence (i.e. after codon 31) was cloned into a pcDNA5 vector that had the prolactin signal sequence, 3xFLAG and TEV cleavage site for N-terminal fusion. *ilcr-1* cDNA was assembled in a pcDNA4 plasmid with EGFP sequences inserted between *Nar*I and *Xba*I sites. *ilcr-2* cDNA was assembled in a pcDNA4 vector that had sequences encoding TEV-3xHA right after the *Xba*I site and in a pcDNA4 plasmid with EGFP sequences inserted between *Nar*I and *Xba*I sites. *actl-1* cDNA was assembled to a pcDNA4 vector that had sequences encoding TEV-3xHA right after the *Xba*I site. *pik-1* cDNA was assembled in a pcDNA4 vector with TEV-3xFLAG for C-terminal tagging.

To tag *ilcr-2* with GFP while including long stretches of 5' and 3' sequences we used fosmid recombineering⁴⁴. The GFP coding sequence was amplified using primers AATCCGACAATGAGGGTGAAGATCCGAAGACGATTGTTGTGAAGAAGTCAATGA GTAAAGGAGAAGAAGCTTTTCAC and GTATCACACCGTAACAAGCAATAAGCAAATAAAAATAACAAGTCTCTTCATTTGTA TAGTTCATCCATGCCATG). The recombineered site was confirmed by sequencing.

Transgenic animals were generated by injecting 50ng/μl of the expression vector with 50ng/μl of a coelomocyte-specific *punc-122::mcherry* marker. Mos-1 mediated single copy gene insertion was performed as described²¹.

Protein sequence alignment

Alignments used T-coffee with default settings in Jalview. The blue shadow indicates the conservation score of residues.

Microscopy

Animals were immobilized using 50 mM sodium azide on 2% agarose pads in M9 buffer. Image acquisition and analysis were performed as described⁴⁵.

Heat-shock experiments

Strains were grown at 15°C, to minimize leaky expression from the *hsp-16.41* promoter. To heat-shock animals NGM plates were wrapped with parafilm and immersed in a 34°C water-bath for 15 or 30 minutes. Animals were transferred to assay plates immediately after heat-shock. The same groups of animals were assayed every 2 hours for 10 hours for their responses to O₂ switches. mCherry expression was measured using an AZ100 microscope (Nikon) fitted with a 553/25 nm excitation filter and a 641/75 nm emission filter. 24 animals were analyzed for each time point and mCherry signals were captured from the whole animal using a 2X lens. Data were analyzed using Image J.

Ca²⁺ imaging

L4 animals expressing the sensor were picked 24 hours prior to imaging. Freely-moving animals were imaged on plates containing NGM41 modified to minimize background fluorescence by omitting peptone and replacing agar with agarose (Fisher, BP164-100).

Plates were dried on the bench overnight and, just before use, spotted with 3 μ l of M9-washed and concentrated OP50 bacteria (grown in 10ml 2xTY to OD600 = 1, spun and resuspended in 1ml M9) to form a 5mm diameter lawn. Animals were picked to the plates, covered by the same microfluidic device used in our behavioral experiments, and allowed to roam freely while we imaged. The O₂ stimulus regime was identical to that in the behavior analysis. Animals were imaged on an AZ100 microscope (Nikon) bearing a TwinCam adapter (Cairn Research, UK) mounted with two ORCA-FLASH4.0 V2 digital cameras (Hamamatsu, Japan) using an AZ Plan Fluor 2x lens with 2x zoom and an exposure time of 100 ms. The excitation light was passed through a 438/24 filter and an FF458-DiO2 dichroic (Semrock) and attenuated to 4.39mW/cm², a light intensity that did not change worm behavior. The TwinCam adapter cube was mounted with CFP (483/32) and YFP (542/27) emission filters and a DC/T510LPXRXT-Uf2 dichroic. Animals were imaged twice, after 10 minutes and 2 hours on the assay plates. Imaging data were analyzed using Neuron Analyzer, a custom-written Matlab program, to analyze the resulting image stacks (code availability: <https://github.com/neuronanalyser/neuronanalyser>).

Ca²⁺ imaging of glued animals used the same hardware and settings, except that a 500 ms exposure time was used. 5 - 6 animals were glued simultaneously onto a 2% agarose pad in M9 on a glass slide, and a small amount of food placed close to their nose. The animals were quickly covered by the microfluidic chamber and 7% O₂ pumped into the chamber for 1 minute before we started imaging.

To image Ca²⁺ responses to pheromone and benzaldehyde we used olfactory chips (Mikrokosmos LLC) as previously described⁴⁶. 100 nM ascaroside and 10⁻⁴ diluted benzaldehyde (diluted in M13 buffer) were used to assess RMG and AIB responses respectively.

Immunofluorescence for mammalian cells

HEK293T cells used in this study were obtained from American Type Culture Collection (ATCC CRL-3216) and were tested for Mycoplasma infection. Cells grown on coverslips were washed with PBS and fixed in 3.7% formaldehyde in PBS for 15 min. Fixed cells were permeabilized with 0.1% Triton X100 in PBS for 5 min, blocked with 10% fetal bovine serum in PBS for 30 min, and incubated with primary antibodies for 1 h. After washing, cells were incubated with AlexaFluor 546-conjugated goat anti-mouse IgG (Invitrogen) for 60 min. Images were acquired on an LSM 780 (Zeiss) using a 63 $\times$ NA 1.42 oil-immersion objective lens.

Immunoprecipitation and immunoblotting

Cell lysates were prepared in lysis buffer [50 mM Tris-HCl pH 7.5, 150 mM NaCl, 1% CHAPS, 1 mM phenylmethanesulfonyl fluoride (PMSF) and protease inhibitor cocktail (complete EDTA-free protease inhibitor, Roche)], clarified by centrifugation at 15,000 rpm, and subjected to immunoprecipitation using anti-HA agarose (Sigma A2095), anti-FLAG M2 affinity gel (Sigma A2220) or GFP-Trap A (Chromotek). Precipitated immunocomplexes were washed five times in a washing buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 1 mM EDTA, and 1% CHAPS) and boiled in sample buffer. Samples were

separated by SDS-PAGE and transferred to Nitrocellulose membranes (Biorad). Immunoblot analysis was performed with anti-Flag M2 (F3165), anti-GFP47 and anti-HA (clone 16B12), and visualized with Super Signal West Pico Chemiluminescent substrate (Pierce).

Purification of FLAG-ILC-17.1

Conditioned medium from Flp-In T-Rex HEK293 cells stably expressing FLAG-ILC-17.1 was collected and concentrated in Amicon Ultra filtration units (Millipore). FLAG-ILC-17.1 was immunoprecipitated using anti-Flag M2 agarose (Sigma), washed with wash buffer (110 mM KAc, 20 mM HEPES pH 7.4 and 2 mM MgCl₂) and eluted using 0.25mg/ml 3xFlag-peptide in PBS.

Flow cytometry analysis

Transfected HEK293T cells were detached with EDTA and kept on ice during the whole staining procedure. Cells were incubated in 10% FBS/PBS containing 0.05ng/μl purified FLAG-ILC-17.1 for 30 min at 4°C. The FLAG tag was detected with the anti-FLAG M2 antibody and AlexaFluor 546-conjugated goat anti-mouse IgG. Stained cells were washed in PBS, re-suspended in 10% FBS/PBS with 1μg/ml DAPI, and analyzed using a flow cytometer (LSRII, BD Biosciences).

Extended Data

Extended data Figure 1. Interleukin-17 related proteins and receptors in *C. elegans*.

a, Schematic of the exon/intron structures of the Y64G10A.6 (*ilcr-1*), F56D1.2 (*ilcr-2*) and T22H6.1 (*ilc-17.1*) genes, with the location of mutations used in assays indicated. -1 FS and -7 FS denote frameshift mutations with 1 and 7 base pair deletions. Other alleles are shown in Extended data Table 1. **b**, Schematic showing the bordering and aggregation assay. The

number of animals on the edge of the food lawn, or in groups, was counted 24 hours after animals were transferred to the assay plates.

c, Bordering and aggregation phenotypes of single, double and triple null mutants of *ilc-17.1*, *ilcr-1* and *ilcr-2*. $n = 4$ assays. ***, $p < 0.001$ (ANOVA with Tukey correction). **d**, CRISPR induced mutations in the F56D1.2 (*ilcr-2*), and the F25D1.3 and C44B12.6 genes, which show homology to mammalian IL-17s. **e**, Bordering and aggregation assays for the F25D1.3 and C44B12.6 mutants. NS, not significant (ANOVA with Tukey correction). **f**, Alignment of the SEFIR domains of IL-17 receptors, including *C. elegans* ILCR-1 and ILCR-2. **g**, Alignment of IL-17D proteins with ILC-17.1. Arrowheads indicate conserved cysteine residues. **h**, ILC-17.1 forms disulfide-linked dimers. SDS-PAGE of affinity-purified FLAG-ILC-17.1 boiled in sample buffer +/- 100 mM DTT. **i**, The O₂ response defects of *ilc-17.1* and *ilcr-1* mutants are rescued by Mos1-mediated single copy insertion (MosSCI) of *ilc-17.1* and *ilcr-1* transgenes. ***, $p < 0.001$ (Mann-Whitney U test). **j**, The O₂ response defects of *ilc-17.1 ilcr-1* and *nfki-1* double mutants with the natural *npr-1 215F* allele. ***, $p < 0.001$ (Mann-Whitney U test).

Extended data Figure 2. ILC-17.1 signaling in RMG promotes aggregation and escape from 21% O₂.

a, b, The bordering and aggregation defects of *ilcr-1* and *ilcr-2* mutants can be rescued by selectively expressing their cDNAs in RMG interneurons (*pnpr-1* and *pflp-5*). Bordering but not aggregation is partially rescued by expression in the O₂-sensing neurons URX, AQR and PQR (*pgcy-32*). *, $p < 0.05$; **, $p < 0.01$; ****, $p < 0.0001$; NS, not significant (ANOVA with Tukey correction). **c, d**, O₂ responses of *ilcr-1* (**c**) and *ilcr-2* (**d**) mutants expressing the corresponding cDNA selectively in RMG (*pnpr-1* and *pflp-5*), the O₂ sensing neurons URX, AQR and PQR (*pgcy-32*), or the ASE gustatory neurons (*pflp-6*). ***, $p < 0.001$. Mann-Whitney U test. **e**, Expressing *ilcr-1* or *pik-1* cDNA in ASG neurons, using the *ops-1* promoter, fails to rescue the O₂ response defects of the corresponding mutants. NS, not significant; ***, $p < 0.001$ (Mann-Whitney U test). **f**, Expressing *ilcr-1* cDNA in cholinergic

neurons (*punc-17* or *pacr-2*), GABAergic neurons (*punc-25*), or interneurons controlling forward and reverse movement (*pglr-1*, *pnmr-1*) fails to restore sustained locomotory arousal to *ilcr-1* mutants kept at 21% O₂. NS, not significant; ***, $p < 0.001$ (Mann-Whitney U test). **g**, Expressing *ilc-17.1* cDNA in RMG using the *flp-5* promoter rescues defective responses to 21% O₂ in *ilc-17.1* mutants. Speed assays were performed 2 hours after animals were transferred to the assay plates. ***, $p < 0.001$ (Mann-Whitney U test). **h**, Expressing *ilc-17.1* cDNA from the *odr-10* promoter (AWA) or the *flp-6* promoter (ASE) not only rescues the *ilc-17.1* defect but confers abnormally heightened escape from 21% O₂. **, $p < 0.01$; ***, $p < 0.001$ (Mann-Whitney U test). **i**, Expressing *ilcr-2* cDNA using the *flp-5* promoter rescues the RMG O₂-evoked Ca²⁺ responses defects of *ilcr-2* mutants. ***, $p < 0.001$ (Mann-Whitney U test). The *npr-1* control is the same as in Figure 4e. **j**, *ilc-17.1; npr-1 215F* and *nfki-1; npr-1 215F* mutants show defective O₂-evoked Ca²⁺ response in RMG. *npr-1 215F* is the allele found in natural isolates of *C. elegans*. ***, $p < 0.001$ (Mann-Whitney U test). **k**, Expression of *pflp-5::GFP* in RMG interneurons, a reporter of RMG neurosecretory activity, is reduced in *ilc-17.1*, *ilcr-1* and *ilcr-2* mutants. ***, $p < 0.001$ (ANOVA with Tukey correction). **l**, **m**, *ilc-17.1*, *pik-1*, and *nfki-1* mutants display normal chemotaxis to NaCl (**l**) and benzaldehyde (**m**). NS, not significant (ANOVA with Tukey correction). **n**, Ca²⁺ responses evoked in AIB interneurons by benzaldehyde (1:10000 dilution) are normal in *ilc-17.1* mutants. bz, benzaldehyde. NS, not significant (Mann-Whitney U test).

Extended data Figure 3. Heat-shock induced expression of ILCR-1 and ILCR-2 in adults rescues behavioral defects of corresponding mutants.

a–d, Transgenic adults expressing *ilcr-1* cDNA (**a, b**) or *ilcr-2* cDNA (**c, d**) from the *hsp-16.41* promoter were assayed either without heat-shock (**a, c**) or after 30 minutes of heat-shock (**b, d**). The assays in (**b**) and (**d**) were performed 8 hours after the heat-shock. **, $p < 0.01$; ***, $p < 0.001$; NS, not significant (Mann-Whitney U test).

Extended data Figure 4. Timeline of heat-shock induced rescue of *ilc-17.1* following 15 and 30-minute heat-shocks.

a, *ilc-17.1* mutants expressing *ilc-17.1* cDNA from the *hsp-16.41* promoter assayed without heat-shock. Assays were performed 2 hours after animals were picked to assay plates. NS, not significant, ***, $p < 0.001$ (Mann-Whitney U test). **b**, **c**, Timelines showing mCherry expression (**b**) and speed at 21% O₂ (**c**) of *ilc-17.1* mutants bearing a *phsp-16.41::ilc-17.1::mCherry* bicistronic transgene and heat shocked for 15 or 30 minutes. $n=24$ for each mCherry measurement; speed in (**c**) is the average of a one-minute time

window, 40s after switching to 21% O₂ and 20s before switching back to 7% O₂, indicated with a red bar in (d, e). The dashed lines in (c) indicate the average speed of *npr-1* (black) and *ilc-17.1 npr-1* (blue) animals for comparison. NS, not significant, **, $p < 0.01$, ***, $p < 0.001$ (Mann-Whitney U test); p values in black are for comparisons to *npr-1* control; p values in blue are for comparisons to *ilc-17.1 npr-1* controls. d, e, Transgenic animals expressing *ilc-17.1* cDNA from the *hsp-16.41* promoter were exposed to 34 °C for 15 (d) or 30 minutes (e), and then assayed every 2 hours. Statistical comparisons between *npr-1* and transgenic animals with heat-shock constructs are indicated in orange. Comparisons between *npr-1 ilc-17.1* and transgenic animals with heat-shock constructs are indicated in black. ***, $p < 0.001$; **, $p < 0.01$; *, $p < 0.05$; NS, not significant (Mann-Whitney U test).

Extended data Figure 5. Overexpressing ILC-17.1 stimulates *pflp-5* neuropeptide expression in RMG.

a, b, Overexpressing *ilc-17.1* stimulates expression from the *pflp-5::GFP* reporter in RMG both in N2 (a) and *npr-1* (b) animals. Conversely, disrupting *ilc-17.1* reduces *pflp-5::gfp* expression. **, $p < 0.01$; ***, $p < 0.001$ (Mann-Whitney U test).

Extended data Figure 6. Phenotypes of *C. elegans* homologs of mammalian genes involved in inflammatory responses.

a, b, Speed was assayed 10 minutes (**a**) and 2 hours (**b**) after animals were transferred to the assay plates. NS, not significant (Mann-Whitney U test).

Extended data Figure 7. The *C. elegans* Act1-like gene.

a, b, *actl-1* alleles. *db789* was isolated in strain AX3544 following mutagenesis using ethylmethanesulfonate (EMS); *db1202* and *db1203* are frameshift mutations generated by CRISPR (**b**). **c**, Mos1 mediated single copy insertion (MosSCI) of the *actl-1*, *pik-1* and *nfki-1* genes rescues the O₂ response defects of the corresponding mutants. ***, $p < 0.001$; **, $p < 0.01$ (Mann-Whitney U test). **d**, A null mutation in *actl-1* confers *ilc-17.1*-like responses to 21% O₂, and does not show additive phenotypes with *ilc-17.1* or *pik-1* null mutations. ***, $p < 0.001$ (Mann-Whitney U test). **e**, The domain architecture of ACTL-1,

showing the locations of the Death and SEFIR domains. **f**, Alignment of the SEFIR domains of ACT1 and ACT1-like proteins. *C. elegans* ACTL-1 is shown at the top.

Extended data Figure 8. Disrupting *pik-1*/IRAK causes *ilc-17.1*-like phenotypes.

a, Schematic drawing of the exon/intron structure of *pik-1*, highlighting the *db842* and *tm2167* mutations used in this study. The allele *db842* was found in strain AX3604; *tm2167* was obtained from the Japanese knockout consortium. **b**, *pik-1* mutants exhibit an *ilc-17.1*-like defect and fail to sustain rapid movement at 21% O₂. The *pik-1* and *ilc-17.1* phenotypes are non-additive, suggesting the proteins encoded by these genes act in the same pathway. **, $p < 0.01$ (Mann-Whitney U test). **c**, The bordering and aggregation defects of *pik-1* mutants can be rescued by expressing *pik-1* cDNA from the *pik-1* or *flp-5* promoters (RMG). ***, $p < 0.001$, ****, $p < 0.0001$ (Mann-Whitney U test). **d**, A functional *ppik-1::pik-1::SL2::mCherry* polycistronic transgene is expressed broadly in the nervous system. RMG expression was confirmed using a *pflp-5::gfp* fiduciary marker. **e**, **f**, Sustained rapid movement of *ilc-17.1* overexpressing animals at 21% O₂ is blocked by mutations in the *pik-1* gene. Speed assays were performed 10 minutes (**e**) and 2 hours (**f**) after picking animals to the assay plates. NS, not significant, ***, $p < 0.001$ (Mann-Whitney U test).

Extended data Figure 9. Mutations in the *nfki-1* gene cause *ilc-17.1*-like phenotypes.

a, b, *nfki-1* alleles. *db923* in strain AX3677 was obtained in an EMS screen; the *db1198* frameshift mutation was generated using CRISPR (**b**). **c**, A functional *pnfki-1::nfki-1::SL2::mCherry* polycistronic transgene is expressed broadly in the nervous system. RMG expression was confirmed using a *pflp-5::gfp* fiduciary marker. **d, e**, Phenotypes of *nfki-1*, *pik-1* and *ilc-17.1* molecular null alleles are not additive. NS, not significant, ***, $p < 0.001$ (Mann-Whitney U test). **f**, Schematic diagram of the human IkBζ b isoform and NFκI-1, highlighting ankyrin repeats (ANK). **g**, Alignment of amino acid

sequences for I κ B ζ orthologs from different species. NFKI-1 is shown at the bottom. Conserved residues are highlighted.

Extended Data Table 1

IL-17 signaling mutants identified by forward genetics

Gene	Strain	Position	Reference	Alteration	Effect	aa change
<i>ilcr-1</i>	AX3344	14278292	C	T	STOP_GAINED	Q159*
	AX3378	14283135	C	T	STOP_GAINED	R548*
	AX3582	14280894	C	T	NON_SYNONYMOUS_CODING	S487F
	AX3667	14279516	TCC	TC	FRAME_SHIFT	-277
	AX3696	14279564	C	T	NON_SYNONYMOUS_CODING	P292L
	AX3836	14279479	C	T	STOP_GAINED	Q264*
<i>ilc-17.1</i>	AX3589	12768673	C	T	SPLICE_SITE_DONOR	/
	AX3617	12767526	C	T	NON_SYNONYMOUS_CODING	G207R
	AX3801	12767612	G	A	NON_SYNONYMOUS_CODING	A194V
	AX3699	12767680	C	T	SPLICE_SITE_ACCEPTOR	/
<i>actl-1</i>	AX3295	6497190	C	T	NON_SYNONYMOUS_CODING	S324N
	AX3544	6499781	C	T	STOP_GAINED	W19*
<i>pik-1</i>	AX3273	13417889	G	A	STOP_GAINED	W30*, W70*
	AX3596	13418117	C	T	STOP_GAINED	Q129*, Q89*
	AX3604	13419353	GAAATGATACATCACAGAA	GA	FRAME_SHIFT	-220, -180
<i>nfki-1</i>	AX3301	15357365	C	T	STOP_GAINED	Q179*
	AX3792	15357721	A	G	NON_SYNONYMOUS_CODING	D247G
	AX3677	15359285	C	T	STOP_GAINED	R286*

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We thank the *Caenorhabditis* Genetics Center (funded by NIH Infrastructure Program P40 OD010440) and the Japanese knockout consortium for strains, Dr. James Hadfield and the Cambridge Research Institute Genomics Core for whole genome sequencing, Zoltan Soltesz for software, Sean Flynn for help with mapping, and de Bono lab members for comments on the manuscript. This work was supported by a European Molecular Biology Organization Fellowship (C.C.), a Japan Society for the Promotion of Science Post-doctoral Fellowship (E.I.), the Medical Research Council, United Kingdom, and the European Research Council (Advanced Grant 269058) to M.d.B.

References

1. Marder E. Neuromodulation of neuronal circuits: back to the future. *Neuron*. 2012; 76:1–11. [PubMed: 23040802]
2. Root CM, Ko KI, Jafari A, Wang JW. Presynaptic facilitation by neuropeptide signaling mediates odor-driven food search. *Cell*. 2011; 145:133–144. [PubMed: 21458672]
3. Lee SH, Dan Y. Neuromodulation of brain states. *Neuron*. 2012; 76:209–222. [PubMed: 23040816]

4. Su CY, Wang JW. Modulation of neural circuits: how stimulus context shapes innate behavior in *Drosophila*. *Curr Opin Neurobiol*. 2014; 29:9–16. [PubMed: 24801064]
5. Oswald D, Waddell S. Olfactory learning skews mushroom body output pathways to steer behavioral choice in *Drosophila*. *Curr Opin Neurobiol*. 2015; 35:178–184. [PubMed: 26496148]
6. Cohn R, Morante I, Ruta V. Coordinated and Compartmentalized Neuromodulation Shapes Sensory Processing in *Drosophila*. *Cell*. 2015; 163:1742–1755. [PubMed: 26687359]
7. McCormick DA, Nusbaum MP. Editorial overview: neuromodulation: tuning the properties of neurons, networks and behavior. *Curr Opin Neurobiol*. 2014; 29:iv–vii. [PubMed: 25457725]
8. Chalfie M, et al. The neural circuit for touch sensitivity in *Caenorhabditis elegans*. *J Neurosci*. 1985; 5:956–964. [PubMed: 3981252]
9. White JG, Southgate E, Thomson JN, Brenner S. The structure of the nervous system of the nematode *Caenorhabditis elegans*. *Philosophical Transactions of the Royal Society of London*. 1986; B:1–340.
10. de Bono M, Tobin DM, Davis MW, Avery L, Bargmann CI. Social feeding in *Caenorhabditis elegans* is induced by neurons that detect aversive stimuli. *Nature*. 2002; 419:899–903. [PubMed: 12410303]
11. Coates JC, de Bono M. Antagonistic pathways in neurons exposed to body fluid regulate social feeding in *Caenorhabditis elegans*. *Nature*. 2002; 419:925–929. [PubMed: 12410311]
12. Gray JM, et al. Oxygen sensation and social feeding mediated by a *C. elegans* guanylate cyclase homologue. *Nature*. 2004; 430:317–322. [PubMed: 15220933]
13. Macosko EZ, et al. A hub-and-spoke circuit drives pheromone attraction and social behaviour in *C. elegans*. *Nature*. 2009; 458:1171–1175. [PubMed: 19349961]
14. Gaffen SL. Structure and signalling in the IL-17 receptor family. *Nat Rev Immunol*. 2009; 9:556–567. [PubMed: 19575028]
15. Hymowitz SG, et al. IL-17s adopt a cystine knot fold: structure and activity of a novel cytokine, IL-17F, and implications for receptor binding. *EMBO J*. 2001; 20:5332–5341. [PubMed: 11574464]
16. Ely LK, Fischer S, Garcia KC. Structural basis of receptor sharing by interleukin 17 cytokines. *Nat Immunol*. 2009; 10:1245–1251. [PubMed: 19838198]
17. Cheung BH, Cohen M, Rogers C, Albayram O, de Bono M. Experience-dependent modulation of *C. elegans* behavior by ambient oxygen. *Curr Biol*. 2005; 15:905–917. [PubMed: 15916947]
18. Rogers C, Persson A, Cheung B, de Bono M. Behavioral motifs and neural pathways coordinating O₂ responses and aggregation in *C. elegans*. *Curr Biol*. 2006; 16:649–659. [PubMed: 16581509]
19. Persson A, et al. Natural variation in a neural globin tunes oxygen sensing in wild *Caenorhabditis elegans*. *Nature*. 2009; 458:1030–1033. [PubMed: 19262507]
20. Busch KE, et al. Tonic signaling from O₂ sensors sets neural circuit activity and behavioral state. *Nat Neurosci*. 2012; 15:581–591. [PubMed: 22388961]
21. Frøkjær-Jensen C, et al. Random and targeted transgene insertion in *Caenorhabditis elegans* using a modified Mos1 transposon. *Nat Methods*. 2014; 11:529–534. [PubMed: 24820376]
22. Kim K, Li C. Expression and regulation of an FMRFamide-related neuropeptide gene family in *Caenorhabditis elegans*. *J Comp Neurol*. 2004; 475:540–550. [PubMed: 15236235]
23. Fares H, Greenwald I. Genetic analysis of endocytosis in *Caenorhabditis elegans*: coelomocyte uptake defective mutants. *Genetics*. 2001; 159:133–145. [PubMed: 11560892]
24. Gaffen SL. Recent advances in the IL-17 cytokine family. *Curr Opin Immunol*. 2011; 23:613–619. [PubMed: 21852080]
25. Iwakura Y, Ishigame H, Saijo S, Nakae S. Functional specialization of interleukin-17 family members. *Immunity*. 2011; 34:149–162. [PubMed: 21349428]
26. Nagai T, Yamada S, Tominaga T, Ichikawa M, Miyawaki A. Expanded dynamic range of fluorescent indicators for Ca²⁺ by circularly permuted yellow fluorescent proteins. *Proc Natl Acad Sci U S A*. 2004; 101:10554–10559. [PubMed: 15247428]
27. Laurent P, et al. Decoding a neural circuit controlling global animal state in *C. elegans*. *Elife*. 2015; 4

28. Jang H, et al. Neuromodulatory state and sex specify alternative behaviors through antagonistic synaptic pathways in *C. elegans*. *Neuron*. 2012; 75:585–592. [PubMed: 22920251]
29. Butcher RA, Fujita M, Schroeder FC, Clardy J. Small-molecule pheromones that control dauer development in *Caenorhabditis elegans*. *Nat Chem Biol*. 2007; 3:420–422. [PubMed: 17558398]
30. Butcher RA, Ragains JR, Kim E, Clardy J. A potent dauer pheromone component in *Caenorhabditis elegans* that acts synergistically with other components. *Proc Natl Acad Sci U S A*. 2008; 105:14288–14292. [PubMed: 18791072]
31. Qian Y, et al. The adaptor Act1 is required for interleukin 17-dependent signaling associated with autoimmune and inflammatory disease. *Nat Immunol*. 2007; 8:247–256. [PubMed: 17277779]
32. Ryzhakov G, Blazek K, Udalova IA. Evolution of vertebrate immunity: sequence and functional analysis of the SEFIR domain family member Act1. *J Mol Evol*. 2011; 72:521–530. [PubMed: 21643828]
33. Pujol N, et al. A reverse genetic analysis of components of the Toll signaling pathway in *Caenorhabditis elegans*. *Curr Biol*. 2001; 11:809–821. [PubMed: 11516642]
34. Kitamura H, Kanehira K, Okita K, Morimatsu M, Saito M. MAIL, a novel nuclear I kappa B protein that potentiates LPS-induced IL-6 production. *FEBS Lett*. 2000; 485:53–56. [PubMed: 11086164]
35. Haruta H, Kato A, Todokoro K. Isolation of a novel interleukin-1-inducible nuclear protein bearing ankyrin-repeat motifs. *J Biol Chem*. 2001; 276:12485–12488. [PubMed: 11278262]
36. Yamazaki S, Muta T, Takeshige K. A novel IkappaB protein, IkappaB-zeta, induced by proinflammatory stimuli, negatively regulates nuclear factor-kappaB in the nuclei. *J Biol Chem*. 2001; 276:27657–27662. [PubMed: 11356851]
37. Kao CY, Kim C, Huang F, Wu R. Requirements for two proximal NF-kappaB binding sites and IkappaB-zeta in IL-17A-induced human beta-defensin 2 expression by conducting airway epithelium. *J Biol Chem*. 2008; 283:15309–15318. [PubMed: 18362142]
38. Karlsen JR, Borregaard N, Cowland JB. Induction of neutrophil gelatinase-associated lipocalin expression by co-stimulation with interleukin-17 and tumor necrosis factor-alpha is controlled by IkappaB-zeta but neither by C/EBP-beta nor C/EBP-delta. *J Biol Chem*. 2010; 285:14088–14100. [PubMed: 20220144]
39. Novatchkova M, Leibbrandt A, Werzowa J, Neubuser A, Eisenhaber F. The STIR-domain superfamily in signal transduction, development and immunity. *Trends Biochem Sci*. 2003; 28:226–229. [PubMed: 12765832]
40. Meisel JD, Kim DH. Behavioral avoidance of pathogenic bacteria by *Caenorhabditis elegans*. *Trends Immunol*. 2014; 35:465–470. [PubMed: 25240986]
41. Sulston, J., Hodgkin, J. The nematode *Caenorhabditis elegans*. Wood, WB., editor. CSHL Press; Cold Spring Harbor: 1988. p. 587-606.
42. Chen C, Fenk LA, de Bono M. Efficient genome editing in *Caenorhabditis elegans* by CRISPR-targeted homologous recombination. *Nucleic Acids Res*. 2013
43. de Bono M, Bargmann CI. Natural variation in a neuropeptide Y receptor homolog modifies social behavior and food response in *C. elegans*. *Cell*. 1998; 94:679–689. [PubMed: 9741632]
44. Tursun B, Cochella L, Carrera I, Hobert O. A toolkit and robust pipeline for the generation of fosmid-based reporter genes in *C. elegans*. *PLoS One*. 2009; 4:e4625. [PubMed: 19259264]
45. Chen C, Itakura E, Weber KP, Hegde RS, de Bono M. An ER complex of ODR-4 and ODR-8/Ufm1 specific protease 2 promotes GPCR maturation by a Ufm1-independent mechanism. *PLoS Genet*. 2014; 10:e1004082. [PubMed: 24603482]
46. Chronis N, Zimmer M, Bargmann CI. Microfluidics for in vivo imaging of neuronal and behavioral activity in *Caenorhabditis elegans*. *Nat Methods*. 2007; 4:727–731. [PubMed: 17704783]
47. Stefanovic S, Hegde RS. Identification of a targeting factor for posttranslational membrane protein insertion into the ER. *Cell*. 2007; 128:1147–1159. [PubMed: 17382883]

Figure 1. IL-17 promotes aggregation and escape from 21% O_2 .

a, Bordering and aggregation phenotypes. n=4 assays. ***, $p < 0.001$; **, $p < 0.01$ (ANOVA with Tukey's correction). Data here and in all Figures show Mean $\pm$ Standard Error of the Mean. **b**, Cell surface expression of ILCR-1-GFP and ILCR-2-HA in HEK293T cells. **c**, Co-immunoprecipitation of ILCR-1 and ILCR-2. **d**, FLAG-ILC-17.1 binding to receptor transfected cells. **e**, FACS of cells transfected with the indicated vector/s and exposed to FLAG-ILC-17.1 or buffer. **f, g**, Average speed of animals kept 2 minutes at 7% O_2 then 2 hours at 21% O_2 . Rescue lines are as in (a). Here and hereafter time intervals at 7% O_2 are open, shading indicates 21% O_2 , and green and red bars on the X-axis indicate intervals used for statistics. n= 4 assays, 120 animals. *, $p < 0.05$; **, $p < 0.01$, and ***, $p < 0.001$. Mann-

Whitney U test. **h, i**, Speed at 7% and 21% O₂ assayed 10 minutes (**h**) or 2 hours (**i**) after animals were transferred to assay plates. n = 3 assays, 90 animals. NS, not significant, ***, $p < 0.001$. Mann-Whitney U test.

Figure 2. Anatomical focus of ILC-17.1 signaling.

a–d *ilcr-1-gfp* and *ilcr-2-gfp* are expressed in many neurons (**a**, **b**), including RMG, identified using *pflp-5::gfp* (**c**, **d**). **e**, *ilcr-1* O₂ response defects rescued by selective expression in RMG (*pnpr-1* and *pflp-5*) ***, *p*<0.001. Mann-Whitney U test. **f**, ILC-17.1 expression. Top left: Functional mCherry-ILC-17.1 accumulates in coelomocytes (labeled with GFP, ccGFP). Top right: a functional *pilc-17.1::ilc-17.1::SL2::mCherry* polycistronic transgene does not highlight coelomocytes but (Bottom) labels the RMG, AUA and ASE neurons. DiO-labeled neurons provide reference points. A, anterior; P, posterior. **g**, mCherry-ILC-17.1 accumulation in coelomocytes is not O₂-dependent. NS, not significant. Student's *t*-test. **h**, Gut-specific expression of *ilc-17.1* cDNA rescues *ilc-17.1* defects. ***, *p*<0.001. Mann-Whitney U test.

Figure 3. ILC-17.1 signaling alters RMG physiology.

a, b, O₂-evoked Ca²⁺ responses in URX (**a**) and RMG (**b**). NS, not significant, **, $p < 0.01$. Mann-Whitney U test. **c,** Effect of stimulating RMG using Channelrhodopsin in animals kept at 7% O₂. ***, $p < 0.001$, Mann-Whitney U test. **d, e,** Disrupting *ilc-17.1* reduces ascaroside-evoked (asc) responses in RMG (**d**) and switches valence of ascaroside pheromones (**e**). **f,** Tapping restores responsiveness to 21% O₂ to *ilc-17.1* mutants. The arrow and purple line indicate taps to the microscope stage at 7 minutes. *mec-4* mutants are touch insensitive. **g, h,** Heat-shock induced expression in adults rescues *ilc-17.1* O₂ responses. Assays were performed 2 hours after animals were picked to the assay plates. The assays in (**h**) were done 8 hours after heat-shock. NS, not significant, ***, $p < 0.001$ (Mann-Whitney U test).

Figure 4. Overexpressing ILC-17.1 heightens responsiveness to 21% O_2

a, Animals overexpressing ILC-17.1 respond more strongly to 21% O_2 than *npr-1* control animals. **, $p < 0.01$ (Mann-Whitney U test). **b**, Overexpressing ILC-17.1 in *N2* animals promotes bordering and aggregation. *, $p < 0.05$ and ****, $p < 0.0001$ (t test). **c**, *N2* animals overexpressing ILC-17.1 show increased responsiveness to 21% O_2 . **, $p < 0.01$ (Mann-Whitney U test). **d–f**, ILC-17.1 overexpression increases O_2 -evoked Ca^{2+} responses in RMG in *N2* (**d**) and *npr-1* (**e**) animals, but does not alter O_2 -evoked Ca^{2+} responses in URX (**f**). NS, not significant; *, $p < 0.05$; **, $p < 0.01$ (Mann-Whitney U test).

Figure 5. ACTL-1, PIK-1/IRAK and NF-κB mediate ILC-17.1 signaling in RMG.

a, b, IP of ILCR-1-GFP and ILCR-2-GFP co-immunoprecipitated ACTL-1-HA (**a**). Conversely, IP of ACTL-1-HA pulled down ILCR-2-GFP and, more weakly, ILCR-1-GFP (**b**). **c, d**, IP of PIK-1-FLAG pulled down ACTL-1-HA (**c**). IP of ACTL-1-HA pulled down PIK-1-FLAG (**d**). **e**, Overexpressing NF-κB rescues the O₂ response defects of *ilcr-1*, *ilcr-2*, *actl-1* and *pik-1* mutants. NS, not significant, ***, $p < 0.001$ (Mann-Whitney U test). **f–h**, The *actl-1* (**f**), *pik-1* (**g**) and *nfki-1* (**h**) O₂ response defects can be rescued by selectively expressing the corresponding cDNA in RMG interneurons (*pnpr-1* and *pflp-5*) but not URX,

AQR and PQR (*pgcy-32*) or ASE (*pflp-6*) sensory neurons. *, $p < 0.05$, ***, $p < 0.001$ (Mann-Whitney U test). **i–k**, O₂-evoked Ca²⁺ responses in RMG of single and double null mutants of *ilc-17.1*, *actl-1*, *pik-1* and *nfki-1*. **, $p < 0.01$, ***, $p < 0.001$ (Mann-Whitney U test). **l**, ILC-17.1 signaling in *C. elegans* compared to IL-17 and Toll-Like Receptor (TLR) pathways in mammals. TF, transcription factor. **m**, Models. ILC-17.1 increases RMG responsiveness to input from the URX O₂ sensors, enabling *C. elegans* arousal at 21% O₂. Without ILC-17.1 *C. elegans* responds poorly to 21% O₂ unless also mechanically stimulated.